

RADEMACHER SUMS, MOONSHINE AND GRAVITY

John F. R. Duncan*

Igor B. Frenkel†

October 21, 2009

Abstract

In 1939 Rademacher derived a conditionally convergent series expression for the elliptic modular invariant, and used this expression—the first Rademacher sum—to verify its modular invariance. By generalizing Rademacher’s approach we construct bases for the spaces of automorphic integrals of arbitrary even integer weight, for groups commensurable with the modular group. Our methods provide explicit expressions for the Fourier expansions of the Rademacher sums we construct at arbitrary cusps, and illuminate various aspects of the structure of the spaces of automorphic integrals, including the actions of Hecke operators.

We give a moduli interpretation for a class of groups commensurable with the modular group which includes all those that are associated to the Monster via monstrous moonshine. We show that within this class the monstrous groups can be characterized just in terms of the behavior of their Rademacher sums. In particular, the genus zero property of monstrous moonshine is encoded naturally in the properties of Rademacher sums.

Just as the elliptic modular invariant gives the graded dimension of the moonshine module, the exponential generating function of the Rademacher sums associated to the modular group furnishes the bi-graded dimension of the Verma module for the Monster Lie algebra. This result generalizes naturally to all the groups of monstrous moonshine, and recovers a certain family of monstrous Lie algebras recently introduced by Carnahan.

Our constructions suggest conjectures relating monstrous moonshine to a distinguished family of chiral three dimensional quantum gravities, and relating monstrous Lie algebras and their Verma modules to the second quantization of this family of chiral three dimensional quantum gravities.

*Department of Pure Mathematics and Mathematical Statistics, Centre for Mathematical Sciences, University of Cambridge, Wilberforce Road, Cambridge, CB3 0WB, U.K.

†Yale University, Department of Mathematics, 10 Hillhouse Avenue, New Haven, CT 06520, U.S.A.
The research of J.D. was conducted during an Assistant Professorship at Harvard University.
The research of I.F. was supported by NSF grant DMS-0457444.

Contents

1	Introduction	3
1.1	Monstrous moonshine	3
1.2	Rademacher sums	4
1.3	Conjugate Rademacher sums	6
1.4	Solid tori	7
1.5	Moonshine via Rademacher sums	8
1.6	Modified Rademacher sums	9
1.7	Hecke operators	12
1.8	Monstrous Lie algebras	14
1.9	Chiral gravity	16
2	Conventions	17
2.1	Functions	17
2.2	Divisors	18
2.3	Isometries	19
2.4	Cosets	21
2.5	Groups	23
2.6	Scalings	24
2.7	Integrals	27
3	Rademacher sums	31
3.1	Construction	31
3.2	Coefficients	35
3.3	Convergence	39
3.4	Variance	41
4	Modified Rademacher sums	48
4.1	Construction	48
4.2	Coefficients	51
4.3	Dirichlet series	52
4.4	Convergence	53
4.5	Variance	59
5	Structural applications	61
5.1	Constants	62
5.2	Inner products	63
5.3	Branching	64

5.4	Fractional orders	68
5.5	Hecke operators	72
6	Moonshine	77
6.1	Genera	77
6.2	Moduli	79
6.3	Exponents	83
6.4	Cusps	84
6.5	Rademacher sums and the Monster	85
7	Gravity	87
7.1	First conjecture	87
7.2	Monstrous Lie algebras	89
7.3	Second conjecture	95
	References	99

1 Introduction

1.1 Monstrous moonshine

A remarkable coincidence between the coefficients of the elliptic modular invariant

$$J(z) = e(-z) + \sum_{n>0} c(n) e(nz), \quad e(nz) = \exp(2\pi i n z), \quad (1.1.1)$$

and the dimensions of the irreducible representations of the *Monster group*, denoted $\mathbb{M}$, led McKay and Thompson [Tho79b] to conjecture the existence of a naturally defined infinite dimensional representation

$$V = V_{-1} \oplus V_1 \oplus V_2 \oplus \cdots \quad (1.1.2)$$

for the Monster group, with the property that $\dim V_n = c(n)$. Thompson [Tho79a] also proposed to consider the modular properties of the series

$$T_g(z) = e(-z) + \sum_{n>0} (\mathrm{tr}|_{V_n} g) e(nz), \quad (1.1.3)$$

for any $g \in \mathbb{M}$, the function $J(z)$ of (1.1.1) being recovered upon taking $g = e$ in (1.1.3). It is clear from the presentation (1.1.3) that $T_g(z)$ is invariant under the *translation group*, which we denote $B(\mathbb{Z})$ and which is generated by the translation $z \mapsto z + 1$. In the case (1.1.1) that g is

the identity element, the invariance extends to the full modular group $PSL_2(\mathbb{Z})$. This leads us to the question: what is special about the invariance groups Γ_g of the McKay–Thompson series (1.1.3)?

In their remarkable paper [CN79], Conway and Norton collected an overwhelming number of coincidences and, in particular, formulated the *moonshine conjecture*:

The series $T_g(z)$ is the normalized hauptmodul of a genus zero group Γ_g lying between $\Gamma_0(N)$ and its normalizer in $PSL_2(\mathbb{R})$, for each $g \in \mathbb{M}$.

The conjecture of McKay and Thompson was proven in [FLM88] by the explicit construction of a vertex operator algebra $V^\natural$ invariant under the Monster group. It was also shown in [FLM88] that the McKay–Thompson series $T_g(z)$ satisfy the moonshine conjecture of Conway and Norton for all elements g in a subgroup of $\mathbb{M}$ arising as an involution centralizer. Finally, a complete proof of the moonshine conjecture was given by Borcherds in [Bor92]. However, in spite of all the subsequent spectacular developments related to the Monster, the consensus of the experts (cf. e.g. [CMS04]), is that “the real nature of moonshine is still remote”.

1.2 Rademacher sums

In this paper we shed new light on the properties of the McKay–Thompson series associated to the Monster, including the mysterious genus zero phenomena. It turns out that the McKay–Thompson series admit presentations as sums over the coset space $B(\mathbb{Z}) \backslash \Gamma_g$. The prototype for this is the following expression for the modular invariant $J(z)$ which was originally derived by Rademacher in [Rad39].

$$J(z) + 12 = e(-z) + \lim_{K \rightarrow \infty} \sum_{\substack{0 < c < K \\ -K^2 < d < K^2}} e\left(-\frac{az+b}{cz+d}\right) - e\left(-\frac{a}{c}\right) \quad (1.2.1)$$

The integers a and b occurring in each summand of (1.2.1) are chosen so that $ad - bc = 1$. This expression (1.2.1) is remarkably simple, but the convergence is rather subtle. (See [Kno90] for a nice exposition.) The subtraction of the constant $e(-a/c)$ in (1.2.1) ensures the existence of the limit in what is otherwise a highly divergent series. The Rademacher sum (1.2.1) has been generalized to various discrete subgroups of $PSL_2(\mathbb{R})$ in a series of papers by Knopp (cf. [Kno61a], [Kno61b], [Kno62a], [Kno62b]).

Given $\Gamma < PSL_2(\mathbb{R})$ we write Γ_∞ for the subgroup of Γ that fixes ∞ . If Γ is commensurable with the modular group $PSL_2(\mathbb{Z})$ and has the property that $\Gamma_\infty = B(\mathbb{Z})$, then we may naturally associate a Rademacher sum to Γ by setting

$$R_\Gamma^{(1)}(z) = e(-z) + \lim_{K \rightarrow \infty} \sum_{\gamma \in (B(\mathbb{Z}) \backslash \Gamma)_{\leq K}^\times} e(-\gamma \cdot z) - e(-\gamma \cdot \infty), \quad (1.2.2)$$

where the sum here is taken over the *rectangle*

$$(B(\mathbb{Z}) \backslash \Gamma)_{\leq K}^{\times} = \left\{ \gamma = B(\mathbb{Z}) \begin{bmatrix} * & * \\ c & d \end{bmatrix} \in B(\mathbb{Z}) \backslash \Gamma \mid 0 < c \leq K, -K^2 \leq d \leq K^2 \right\}. \quad (1.2.3)$$

In (1.2.3) we write $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$ for the image in $PSL_2(\mathbb{R})$ of a matrix $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ in $SL_2(\mathbb{R})$, and in (1.2.2) we write $\gamma \cdot z$ for $(az + b)/(cz + d)$ and $\gamma \cdot \infty$ for a/c when $\gamma = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$.

In general the function $R_{\Gamma}^{(1)}(z)$ defines an abelian integral for Γ , meaning that we have

$$R_{\Gamma}^{(1)}(\gamma \cdot z) = R_{\Gamma}^{(1)}(z) + \omega(\gamma) \quad (1.2.4)$$

for some function $\omega : \Gamma \rightarrow \mathbb{C}$. We show (cf. §6.1) that the Rademacher sum $R_{\Gamma}^{(1)}(z)$ is Γ -invariant if and only if Γ has genus zero. Even more than this, we show that when Γ has genus zero the function

$$R_{\Gamma}^{(1)}(z) - \frac{1}{2}c_{\Gamma}(1, 0) \quad (1.2.5)$$

is the normalized hauptmodul for Γ , for a certain constant $c_{\Gamma}(1, 0)$. This constant $c_{\Gamma}(1, 0)$ turns out to be the *Rademacher constant associated to* Γ as defined in [Nor84] in the case that Γ has genus zero.

Rademacher's proof of the validity of the presentation (1.2.1) relies upon explicit formulas for the Fourier coefficients of $J(z)$. These formulas may be given in terms of Kloosterman sums and Bessel functions, and are recovered from the expression

$$c_{\Gamma}(m, n) = \sum_{\gamma \in B(\mathbb{Z}) \backslash \Gamma / B(\mathbb{Z})} \text{Kl}_{\gamma}(m, n) \text{Bl}_{\gamma}(m, n) \quad (1.2.6)$$

upon taking $\Gamma = PSL_2(\mathbb{Z})$ and $m = 1$, and allowing n to range over $\mathbb{Z}$. Writing $c(\gamma)$ for $|c|$ when $\gamma = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, the functions $\text{Kl}_{\gamma}(m, n)$ and $\text{Bl}_{\gamma}(m, n)$ are defined as follows (cf. §3.2), for $\gamma \in \Gamma$.

$$\text{Kl}_{\gamma}(m, n) = e(-m\gamma \cdot \infty) e(-n\gamma^{-1} \cdot \infty) \quad (1.2.7)$$

$$\text{Bl}_{\gamma}(m, n) = 2\pi i \text{Res}_{\xi=0} e(-mc(\gamma)^{-2}\xi^{-1}) e(n\xi) d\xi \quad (1.2.8)$$

Note that $\text{Kl}_{\gamma}(m, n)$ and $\text{Bl}_{\gamma}(m, n)$ are well-defined complex numbers only when $c(\gamma) \neq 0$, and in this case

$$\text{Kl}_{\gamma}(m, n) = e\left(\frac{-ma + nd}{c}\right), \quad \text{Bl}_{\gamma}(m, n) = \sum_{k \in \mathbb{N}} (4\pi^2)^{k+1} c^{-2k-2} \frac{m^{k+1}}{(k+1)!} \frac{n^k}{k!}, \quad (1.2.9)$$

for $\gamma = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ and $c > 0$. The expression (1.2.6) with $\Gamma = PSL_2(\mathbb{Z})$ and $m = 1$ is, up to

elementary transformations, the formula for the n -th coefficient of the elliptic modular invariant given originally by Rademacher in [Rad38]. More generally, the formula (1.2.6) recovers the Fourier coefficients (other than the constant term) of the Rademacher sum $R_\Gamma^{(1)}(z)$, for Γ an arbitrary group commensurable with $PSL_2(\mathbb{Z})$ and satisfying $\Gamma_\infty = B(\mathbb{Z})$. The following formula encodes the relationship precisely.

$$R_\Gamma^{(1)}(z) = e(-z) + \frac{1}{2}c_\Gamma(1, 0) + \sum_{n \in \mathbb{Z}^+} c_\Gamma(1, n) e(nz) \quad (1.2.10)$$

1.3 Conjugate Rademacher sums

It is very striking that the constant term in the Fourier expansion (1.2.10) of the Rademacher sum $R_\Gamma^{(1)}(z)$ is $c_\Gamma(1, 0)/2$ and not $c_\Gamma(1, 0)$, given that the coefficient of $e(nz)$ in (1.2.10) is exactly $c_\Gamma(1, n)$ for $n > 0$. One may also observe that the formula (1.2.6) defining the values $c_\Gamma(1, n)$ makes sense for arbitrary integers $n \in \mathbb{Z}$, and so it is natural to ask what rôle is played by the $c_\Gamma(1, n)$ for $n < 0$? An answer to this question is obtained when we consider the *conjugate Rademacher sum associated to Γ* , denoted $\bar{R}_\Gamma^{(1)}(z)$, and defined by setting

$$\bar{R}_\Gamma^{(1)}(z) = e(-\bar{z}) + \lim_{K \rightarrow \infty} \sum_{\gamma \in (B(\mathbb{Z}) \backslash \Gamma)_{\leq K}^\times} e(-\gamma \cdot \bar{z}) - e(-\gamma \cdot \infty) \quad (1.3.1)$$

for $z \in \mathbb{H}$. The conjugate Rademacher sum $\bar{R}_\Gamma^{(1)}(z)$ defines an anti-holomorphic function on $\mathbb{H}$. At first glance it appears that we should recover the classical Rademacher sum $R_\Gamma^{(1)}(z)$ after substituting $\bar{z}$ for z in $\bar{R}_\Gamma^{(1)}(z)$, since the expressions defining $R_\Gamma^{(1)}(z)$ and $\bar{R}_\Gamma^{(1)}(\bar{z})$ would appear to coincide, but in fact the delicate limit defining the Rademacher sums behaves very differently depending on which half plane the variable z lies in, and the difference $R_\Gamma^{(1)}(z) - \bar{R}_\Gamma^{(1)}(\bar{z})$ can be rather far from vanishing. As an illustration of this, we show in §3.3 that the Fourier expansion of the conjugate Rademacher sum $\bar{R}_\Gamma^{(1)}(z)$ in terms of the anti-holomorphic exponential $e(\bar{z})$ recovers the values $c_\Gamma(1, n)$ for $n < 0$ as Fourier coefficients. More precisely, we establish the following counterpart to (1.2.10).

$$\bar{R}_\Gamma^{(1)}(z) = e(-\bar{z}) - \frac{1}{2}c_\Gamma(1, 0) - \sum_{n \in \mathbb{Z}^+} c_\Gamma(1, -n) e(-n\bar{z}) \quad (1.3.2)$$

We demonstrate in §3.4 that the holomorphic function $\bar{R}_\Gamma^{(1)}(\bar{z})$ is also an abelian integral. More than this, the difference

$$R_\Gamma^{(1)}(z) - \bar{R}_\Gamma^{(1)}(z) \quad (1.3.3)$$

is a Γ -invariant harmonic function on $\mathbb{H}$.

The Γ -invariance of the function (1.3.3) entails a remarkable formula for $\bar{R}_\Gamma^{(1)}(z)$ in the case that Γ has genus zero. For since the holomorphic Rademacher sum $R_\Gamma^{(1)}(z)$ is Γ -invariant in the genus zero case, the anti-holomorphic function $\bar{R}_\Gamma^{(1)}(z)$ must also be Γ -invariant. Given our knowledge (1.3.2) of the Fourier expansion of $\bar{R}_\Gamma^{(1)}(z)$ it follows that $\bar{R}_\Gamma^{(1)}(z)$ must be identically constant, and furthermore, this constant must be $-c_\Gamma(1, 0)/2$. In the case that $\Gamma = PSL_2(\mathbb{Z})$ we have $c_\Gamma(1, 0) = 24$, and we thus obtain the remarkable identity

$$e(-\bar{z}) + \lim_{K \rightarrow \infty} \sum_{\substack{0 < c < K \\ -K^2 < d < K^2}} e\left(-\frac{a\bar{z} + b}{c\bar{z} + d}\right) - e\left(-\frac{a}{c}\right) = -12, \quad (1.3.4)$$

where in each summand a and b are integers chosen so that $ad - bc = 1$. This implies that the function $R_\Gamma^{(1)}(z) - \bar{R}_\Gamma^{(1)}(z)$ is a (non-normalized) hauptmodul for Γ when Γ has genus zero, and in particular, $c_\Gamma(1, n) = \delta_{1+n, 0}$ for $n < 0$. We recover the normalized hauptmodul for Γ by considering the sum $R_\Gamma^{(1)}(z) + \bar{R}_\Gamma^{(1)}(z)$.

1.4 Solid tori

It was observed in [CN79] that each group attached to the Monster via monstrous moonshine may be described as a group of $n||h$ -type, where a discrete group $\Gamma < PSL_2(\mathbb{R})$ is said to be of $n||h$ -type if it is of the form $\Gamma = \Gamma_0(n||h) + S$ for some $n, h \in \mathbb{Z}^+$ with $h|(n, 24)$, and S a subgroup of the group of exact divisors of n/h . For such n, h , and S , the group $\Gamma_0(n||h) + S$ contains and normalizes $\Gamma_0(nh)$, and has the property that $(\Gamma_0(n||h) + S)_\infty = B(\mathbb{Z})$. (We recall the precise definition of $\Gamma_0(n||h) + S$ in §2.5.)

We shed new light on the groups of $n||h$ -type by demonstrating in §6.2 that they solve a natural family of moduli problems for elliptic curves equipped with a certain kind of extra structure. As we will explain presently, a common feature of all the members of this family is a choice of filling of the elliptic curve in question; that is, a solid torus whose boundary is the given elliptic curve. Hence, we interpret the $n||h$ -type groups as defining moduli of (decorated) solid tori.

The fact that each group Γ of $n||h$ -type satisfies $\Gamma_\infty = B(\mathbb{Z})$ suggests the importance of the quotient $B(\mathbb{Z}) \backslash \mathbb{H}$. Regarding $\mathbb{H}$ as a moduli space for triples (E, γ, γ') where E is an elliptic curve and (γ, γ') is an oriented basis for the first homology group $H_1(E, \mathbb{Z})$ of E , we see that $B(\mathbb{Z}) \backslash \mathbb{H}$ parameterizes pairs (E, γ) where E is an elliptic curve and γ is a primitive element of $H_1(E, \mathbb{Z})$. To each such pair (E, γ) is naturally associated an infinite volume hyperbolic 3-manifold with boundary an elliptic curve, for the cycle γ specifies a way to fill in the surface defined by E , thus yielding a solid torus. Every complete, smooth, infinite volume hyperbolic 3-manifold with boundary an elliptic curve arises in this fashion, so we consider pairs (E, C) where E is an elliptic curve over $\mathbb{C}$ and C is an oriented subgroup of E isomorphic to S^1 . We

call such a pair a *solid torus*; according to our discussion the quotient space $B(\mathbb{Z}) \backslash \mathbb{H}$ naturally parameterizes solid tori.

For (E, C) a solid torus and $n \in \mathbb{Z}^+$ we write $C[n]$ for the group of n -division points of C , and we define an *n -isogeny of solid tori* $(E', C') \rightarrow (E, C)$ to be an isogeny $E' \rightarrow E$ of elliptic curves that maps $C'[n]$ to a subgroup of $C[n]$, and we define an *isogeny of solid tori* to be an isogeny $E' \rightarrow E$ of elliptic curves that restricts to an orientation preserving map $C' \rightarrow C$ on the underlying primitive cycles. For (E, C) a solid torus, the canonical map $E \rightarrow E/C[n]$ may be viewed as defining an isogeny $(E, C) \rightarrow (E, C)/C[n]$ of solid tori, where $(E, C)/C[n]$ is a shorthand for the solid torus whose underlying elliptic curve is $E/C[n]$, and whose primitive cycle is the image of C under the natural map $E \rightarrow E/C[n]$. It may happen that an n -isogeny induces invertible n/e -isogenies (i.e. n/e -isomorphisms) $(E', C')/C'[e] \rightarrow (E, C)$ and $(E', C') \rightarrow (E, C)/C[e]$ for some divisor e of n . In this case we say that (E', C') and (E, C) are *$n + e$ -related*. By considering the existence or otherwise of such morphisms of solid tori we arrive at moduli interpretations for all the $n||h$ -type groups. For example, given a subgroup S of the group of exact divisors of n , say solid tori (E, C) and (E', C') are *$n + S$ -related* if (E, C) and (E', C') are $n + e$ -related for some $e \in S$. Then the $n + S$ -relation is an equivalence relation on solid tori, and the quotient $(\Gamma_0(n) + S) \backslash \mathbb{H}$ is a moduli space for $n + S$ -equivalence classes of solid tori.

In this way we demonstrate that the groups of $n||h$ -type may be explained as being the groups that solve a natural family of moduli problems for decorated solid tori with conformal structure on the boundary.

1.5 Moonshine via Rademacher sums

The fact that the function (1.2.5) recovers the normalized hauptmodul of the group Γ when Γ has genus zero is a strong indication that the Rademacher sums may be useful for the purpose of understanding the functions of monstrous moonshine. A group theoretic characterization of the functions arising as McKay–Thompson series was given in [CMS04]. There the authors showed that a holomorphic function on $\mathbb{H}$ coincides with $T_g(z)$ for some $g \in \mathbb{M}$ if and only if it is the normalized hauptmodul for a group Γ satisfying each of the following properties.

1. The Riemann surface $\Gamma \backslash \mathbb{H}$ has genus zero.
2. The group Γ is of $n||h$ -type.
3. The group Γ is an extension of $\Gamma_0(nh)$ by a group of exponent 2.
4. Each cusp of Γ can be mapped to the infinite cusp by an element $\sigma \in \text{PSL}_2(\mathbb{R})$ with the property that $(\sigma\Gamma\sigma^{-1})_\infty = B(\mathbb{Z})$ and the intersection $\Gamma \cap \sigma\Gamma\sigma^{-1}$ contains $\Gamma_0(nh)$.

As we saw in §1.2 the variance of the Rademacher sum $R_\Gamma^{(1)}(z)$ with respect to the action of Γ detects whether or not Γ has genus zero, and we have discussed in §1.4 the significance of the

groups appearing in condition 2. We show in §§6.3,6.4 that the conditions 3 and 4 also admit natural reformulations in terms of the Rademacher sums.

Our reformulation comes into view when we generalize the construction (1.2.2) so as to associate a Rademacher sum $R_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(1)}(z)$ to each triple $(\Gamma, \mathfrak{p}, \mathfrak{q})$ where Γ is a group commensurable with $PSL_2(\mathbb{Z})$, and $\mathfrak{p}$ and $\mathfrak{q}$ are cusps of Γ . The function $R_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(1)}(z)$ is to be regarded as encoding the *expansion at $\mathfrak{q}$* of the *Rademacher sum $R_{\Gamma, \mathfrak{p}}^{(1)}(z)$ associated to Γ at $\mathfrak{p}$* , although this point of view is accurate only up to certain constant functions (cf. §4.5). Here we write $R_{\Gamma, \mathfrak{p}}^{(1)}(z)$ as a shorthand for $R_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(1)}(z)$ when $\mathfrak{q}$ is the *infinite cusp* $\Gamma \cdot \infty$. It will develop in §3.4 that $R_{\Gamma, \mathfrak{p}}^{(1)}(z)$ is an abelian integral for Γ with a simple pole at $\mathfrak{p}$, and no other singularities, and the behavior of $R_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(1)}(z)$ as $z \rightarrow i\infty$ encodes the behavior of $R_{\Gamma, \mathfrak{p}}^{(1)}(z)$ as $z \rightarrow \mathfrak{q}$, up to a certain constant function. Considering all the functions $R_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(1)}(z)$ for varying cusps $\mathfrak{p}$ and $\mathfrak{q}$ we are able to access more subtle properties of the curve $X_\Gamma = \Gamma \backslash \mathbb{H} \cup \Gamma \backslash \hat{\mathbb{Q}}$ (cf. §2.5) beyond its genus. We find in §6.3 that the condition 3 translates into a certain condition of symmetry in the Rademacher sums associated to the Hecke congruence group $\Gamma_0(nh)$, where n and h are as in condition 2, and we find in §6.4 that the condition 4 translates into the statement that the function $R_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(1)}$ be invariant for this subgroup $\Gamma_0(nh)$.

Our moduli interpretation of the $n||h$ -type groups, together with the invariance and symmetry conditions on Rademacher sums just described, facilitate a reformulation of the conditions of [CMS04]. We thus arrive at a new characterization of the functions of monstrous moonshine in terms of Rademacher sums, and moduli of solid tori.

1.6 Modified Rademacher sums

In order to understand better the behavior of the Rademacher sums $R_\Gamma^{(1)}(z)$, and, in particular, the subtraction of the constant $e(-\gamma \cdot \infty)$ in (1.2.2), we consider in §3 a generalization of the Rademacher sum adapted to the problem of constructing modular forms of arbitrary even integer weight. Our approach to Rademacher sums of negative weight is inspired by the work of Niebur in [Nie74].

For $\kappa, m \in \mathbb{Z}$ we define the *Rademacher sum of weight 2κ and order m associated to Γ* by setting

$$R_\Gamma^{\kappa(m)}(z) = e(-mz) + \lim_{K \rightarrow \infty} \sum_{\gamma \in (B(\mathbb{Z}) \backslash \Gamma)_{\leq K}^\times} e(-m\gamma \cdot z) r^\kappa(m, \gamma, z) (cz + d)^{-2\kappa}, \quad (1.6.1)$$

where $c, d \in \mathbb{R}$ are chosen (for each summand) so that γ is the image of a matrix $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbb{R})$ in $PSL_2(\mathbb{R})$, and $r^\kappa(m, \gamma, z)$ is the *Rademacher regularization factor of weight 2κ* , satisfying

$$r^\kappa(m, \gamma, z) = 1 - e(m\gamma \cdot z - m\gamma \cdot \infty) e(m\gamma \cdot \infty - m\gamma \cdot z)_{<1-2\kappa}, \quad (1.6.2)$$

where $e(z)_{<K}$ denotes the *partial exponential function* $e(z)_{<K} = \sum_{k \in \mathbb{N}, k < K} (2\pi i z)^k / k!$. Observe that we recover the Rademacher sum $R_\Gamma^{(1)}(z)$ of (1.2.2) when $\kappa = 0$. Also, we have $r^\kappa(m, \gamma, z) = 1$ when κ is positive, so that the Rademacher sum $R_\Gamma^{\kappa(m)}(z)$ is a holomorphic Poincaré series (cf. §3.1) when $\kappa \in \mathbb{Z}^+$ (and absolutely convergent for $\kappa > 1$).

In §3.1 we furnish generalizations of the functions $\text{Kl}_\gamma(m, n)$ and $\text{Bl}_\gamma(m, n)$ for arbitrary $\kappa \in \mathbb{Z}$, obtaining a generalization of the formula (1.2.6), and leading to the following analogue of (1.2.10).

$$R_\Gamma^{\kappa(m)}(z) = e(-mz) + \frac{1}{2}c_\Gamma^\kappa(m, 0) + \sum_{n \in \mathbb{Z}^+} c_\Gamma^\kappa(m, n) e(nz) \quad (1.6.3)$$

The *conjugate Rademacher sum of weight 2κ and order m associated to Γ* is defined in direct analogy with the definition (1.3.1) of the conjugate Rademacher sum $\bar{R}_\Gamma^{(1)}(z)$, which we presently recognize as the conjugate Rademacher sum of weight 0 and order 1 associated to Γ .

$$\bar{R}_\Gamma^{\kappa(m)}(z) = e(-m\bar{z}) + \lim_{K \rightarrow \infty} \sum_{\gamma \in (B(\mathbb{Z}) \backslash \Gamma)_{\leq K}^\times} e(-m\gamma \cdot \bar{z}) r^\kappa(m, \gamma, \bar{z}) (c\bar{z} + d)^{-2\kappa}. \quad (1.6.4)$$

We have the following analogue of (1.3.2), and counterpart to (1.6.3), which leads naturally to analogues of (1.3.3) for arbitrary $\kappa \in \mathbb{Z}$.

$$\bar{R}_\Gamma^{\kappa(m)}(z) = e(-m\bar{z}) - \frac{1}{2}c_\Gamma^\kappa(m, 0) - \sum_{n \in \mathbb{Z}^+} c_\Gamma^\kappa(m, -n) e(-n\bar{z}) \quad (1.6.5)$$

The variance of the functions $R_\Gamma^{\kappa(m)}(z)$ with respect to the natural (weight 2κ) action of the group Γ was described by Niebur in [Nie74] for the case that Γ has a single cusp. We verify that for Γ an arbitrary group commensurable with the modular group $PSL_2(\mathbb{Z})$, the function $R_\Gamma^{\kappa(m)}(z) + c_\Gamma^\kappa(m, 0)/2$ defines an *automorphic integral of weight 2κ for Γ* (cf. §2.7) when $m \in \mathbb{Z}^+$, meaning that $R_\Gamma^{\kappa(m)}(z)$ is holomorphic on $\mathbb{H}$, possibly having poles at cusps of Γ , and we have

$$\left(R_\Gamma^{\kappa(m)}(\gamma \cdot z) + \frac{1}{2}c_\Gamma^\kappa(m, 0) \right) (cz + d)^{-2\kappa} = R_\Gamma^{\kappa(m)}(z) + \frac{1}{2}c_\Gamma^\kappa(m, 0) + \omega(\gamma)(z) \quad (1.6.6)$$

for some function $\omega : \Gamma \rightarrow \mathbb{C}[z]$, where the polynomial $\omega(\gamma)(z)$ has degree at most -2κ in z . An identification of the function ω may be given in terms of a certain canonically defined map $I_\kappa(\Gamma) \rightarrow S_{1-\kappa}(\Gamma)$, where $I_\kappa(\Gamma)$ denotes the space of automorphic integrals of weight 2κ for Γ , and $S_\kappa(\Gamma)$ denotes the space of cusp forms of weight 2κ for Γ . We show in §4.5 that the sequence

$$0 \rightarrow M_\kappa(\Gamma) \rightarrow I_\kappa(\Gamma) \rightarrow S_{1-\kappa}(\Gamma) \rightarrow 0 \quad (1.6.7)$$

is exact, where $M_\kappa(\Gamma)$ denotes the space of modular forms of weight 2κ for Γ possibly having

poles at cusps (cf. §2.7), and the map $M_\kappa(\Gamma) \rightarrow I_\kappa(\Gamma)$ is the natural inclusion.

Apart from the delicate convergence of the limit defining the Rademacher sum $R_\Gamma^{\kappa(m)}(z)$, the most curious feature of the functions $R_\Gamma^{\kappa(m)}(z)$ is that they transform naturally with respect to the group Γ only after the addition of the constant $c_\Gamma^\kappa(m, 0)/2$, which is typically non-zero. We overcome this feature of the classical Rademacher sum by introducing in §4 the *continued Rademacher sum of weight 2κ and order m associated to Γ* , defined for $1 - \kappa, m \in \mathbb{Z}^+$ by setting

$$T_\Gamma^{\kappa(m)}(z, s) = e(-mz) + \sum_{\gamma \in (B(\mathbb{Z}) \backslash \Gamma)^\times} e(-m\gamma \cdot z) t^\kappa(m, \gamma, z, s) (cz + d)^{-2\kappa} \quad (1.6.8)$$

where $t^\kappa(m, \gamma, z, s)$ is a generalization of the Rademacher regularization factor $r^\kappa(m, \gamma, z)$ of (1.6.2) satisfying $t^\kappa(m, \gamma, z, 1) = r^\kappa(m, \gamma, z)$.

The right hand side of (1.6.8) converges absolutely for $z \in \mathbb{H}$ and $\Re(s) > 1$. We define $T_\Gamma^{\kappa(m)}(z)$ by taking the limit as s tends to 1 in $T_\Gamma^{\kappa(m)}(z, s)$. We find that

$$T_\Gamma^{\kappa(m)}(z) = \lim_{s \rightarrow 1^+} T_\Gamma^{\kappa(m)}(z, s) = R_\Gamma^{\kappa(m)}(z) - \frac{1}{2} c_\Gamma^\kappa(m, 0), \quad (1.6.9)$$

so that the Fourier expansion of $T_\Gamma^{\kappa(m)}(z)$ has vanishing constant term. Then, in the case that Γ has genus zero, the function $T_\Gamma^{(1)}(z) = T_\Gamma^{0(1)}(z)$ is precisely the normalized hauptmodul of Γ (cf. §2.7). We call $T_\Gamma^{\kappa(m)}(z)$ the *normalized Rademacher sum of weight 2κ and order m associated to Γ* . We may regard the disappearance of the constant term in (1.6.9) as a consequence of the noncommutativity of the limits $s \rightarrow 1$ and $K \rightarrow \infty$ when applied to (the expression obtained by replacing $(B(\mathbb{Z}) \backslash \Gamma)^\times$ with $(B(\mathbb{Z}) \backslash \Gamma)_{\leq K}^\times$ in the right hand side of (1.6.8).

To recover functions with natural modular properties at non-positive weights we modify the continued Rademacher sum by subtracting a renormalized value at $z = 0$.

$$Q_\Gamma^{\kappa(m)}(z, s) = T_\Gamma^{\kappa(m)}(z, s) - T_{0, \Gamma}^{\kappa(m)}(s) \quad (1.6.10)$$

The function $T_{0, \Gamma}^{\kappa(m)}(s)$ is defined by an expression analogous to (1.6.8) which also converges absolutely for $\Re(s) > 1$, and would vanish at $s = 1$ if it were not for the noncommutativity of limits as $s \rightarrow 1$ and $K \rightarrow \infty$. We define the *modified Rademacher sum of weight 2κ and order m associated to Γ* , denoted $Q_\Gamma^{\kappa(m)}(z)$, by taking the limit as s tends to 1 in $Q_\Gamma^{\kappa(m)}(z, s)$, and as a counterpart to (1.6.9) we have

$$Q_\Gamma^{\kappa(m)}(z) = \lim_{s \rightarrow 1^+} Q_\Gamma^{\kappa(m)}(z, s) = R_\Gamma^{\kappa(m)}(z) + \frac{1}{2} c_\Gamma^\kappa(m, 0), \quad (1.6.11)$$

so that, in light of (1.6.6), the modified Rademacher sum $Q_\Gamma^{\kappa(m)}(z)$ is an automorphic integral of weight 2κ for Γ .

The definition of the modified Rademacher sum $Q_\Gamma^{\kappa(m)}(z)$ is inspired by Hurwitz's relation for the Hurwitz zeta function, and our continuation procedure may be regarded as identifying the factor $1/2$, appearing in the constant term of the Fourier expansion (1.6.3) of the classical Rademacher sum of non-positive weight, with -1 times the value of the Riemann zeta function $\zeta(s)$ at $s = 0$.

In order to obtain and analyze spanning sets for spaces of automorphic integrals we attach modified Rademacher sums $Q_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z)$ to each triple $(\Gamma, \mathfrak{p}, \mathfrak{q})$ where Γ is a group commensurable with the modular group $PSL_2(\mathbb{Z})$ and $\mathfrak{p}$ and $\mathfrak{q}$ are cusps of Γ . The function $Q_\Gamma^{\kappa(m)}(z)$ is then recovered upon taking both $\mathfrak{p}$ and $\mathfrak{q}$ to be the *infinite cusp* $\Gamma \cdot \infty$. If we write $Q_{\Gamma, \mathfrak{p}}^{\kappa(m)}(z)$ as a shorthand for $Q_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z)$ when $\mathfrak{q} = \Gamma \cdot \infty$, then the modified Rademacher sums $Q_{\Gamma, \mathfrak{p}}^{\kappa(m)}(z)$, for varying cusps $\mathfrak{p} \in \Gamma \backslash \hat{\mathbb{Q}}$ and positive integers $m \in \mathbb{Z}^+$, constitute a basis for the space $I_\kappa(\Gamma)$ when $\kappa < 0$. In the case that $\kappa = 0$ the modified Rademacher sums $Q_{\Gamma, \mathfrak{p}}^{\kappa(m)}(z)$ span a subspace of $I_0(\Gamma)$ of codimension 1, and a full basis is obtained by including a constant function. The function $Q_{\Gamma, \mathfrak{p}}^{\kappa(m)}(z)$ defines an automorphic integral with a single pole at the cusp $\mathfrak{p}$, and the function $Q_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z)$ encodes the Fourier expansion of $Q_{\Gamma, \mathfrak{p}}^{\kappa(m)}(z)$ at the cusp $\mathfrak{q}$, and no correction of constant terms is necessary for the validity of these statements.

1.7 Hecke operators

The original result of Rademacher [Rad39] identifies the first Rademacher sum $R_\Gamma^{(1)}(z)$, for $\Gamma = PSL_2(\mathbb{Z})$, with the function $J(z) + 12$, where $J(z)$ is the normalized hauptmodul for the modular group. In light of this it is natural to try to identify more general Rademacher sums in a similar fashion. In this section we will take $\kappa = 0$, but we will allow m to be an arbitrary non-zero integer. The identity

$$R_\Gamma^{(-m)}(z) = \overline{\bar{R}_\Gamma^{(m)}(z)} \quad (1.7.1)$$

follows immediately from the definitions of the classical and conjugate Rademacher sums, demonstrating that the conjugate Rademacher sums of positive order may be recovered from the classical Rademacher sums with negative order. On the other hand, one can show, as we did for the special case that $m = 1$, that $\bar{R}_\Gamma^{(m)}(z)$ is the constant function with constant value $-c_\Gamma(m, 0)/2$ in case Γ has genus zero. When $\Gamma = PSL_2(\mathbb{Z})$ we have $c_\Gamma(m, 0) = 24\sigma(m, 1)$, where $\sigma(m, 1)$ is the sum of the divisors of m , so we have the following generalization of (1.3.4) for $m \in \mathbb{Z}^+$.

$$e(-m\bar{z}) + \lim_{K \rightarrow \infty} \sum_{\substack{0 < c < K \\ -K^2 < d < K^2}} e\left(-\frac{ma\bar{z} + mb}{c\bar{z} + d}\right) - e\left(-\frac{ma}{c}\right) = -12\sigma(m, 1) \quad (1.7.2)$$

To obtain an expression for the classical Rademacher sums $R_\Gamma^{(m)}(z)$ with positive m we can

analyze again its Fourier coefficients $c_\Gamma(m, n)$. We show in §6.1 that $c_\Gamma(m, n) = \delta_{m+n,0}$ for $n < 0$ in case Γ has genus zero. The resulting function $T_\Gamma^{(m)}(z) = R_\Gamma^{(m)}(z) - c_\Gamma(m, 0)/2$, satisfying the identity

$$T_\Gamma^{(m)}(z) + \frac{1}{2}c_\Gamma(m, 0) = e(-mz) + \lim_{K \rightarrow \infty} \sum_{\gamma \in (B(\mathbb{Z}) \backslash \Gamma)_{\leq K}^\times} e(-m\gamma \cdot z) - e(-m\gamma \cdot \infty), \quad (1.7.3)$$

is therefore holomorphic on the upper half plane, invariant for the action of Γ , and of the form $q^{-m} + o(1)$ for $q = e(z)$. Consequently each normalized Rademacher sum $T_\Gamma^{(m)}(z)$, for $m \in \mathbb{Z}^+$, is expressible as a degree m polynomial in $T_\Gamma^{(1)}(z) = T_\Gamma(z)$. This polynomial is none other than the so-called *m-th Faber polynomial for $T_\Gamma(z)$* .

Proceeding from another direction, we can obtain the functions $T_\Gamma^{(m)}(z)$ from the normalized hauptmodul $T_\Gamma(z)$ by applying Hecke operators. In §5.5 we study, in particular, the case that Γ is the modular group $PSL_2(\mathbb{Z})$. Recall that for $n \in \mathbb{Z}^+$ the action of the Hecke operator $\hat{T}(n)$ on a modular function $f(z)$ may be given by setting

$$(\hat{T}(n)f)(z) = \frac{1}{n} \sum_{\substack{ad=n \\ 0 \leq b < d}} f\left(\frac{az+b}{d}\right). \quad (1.7.4)$$

From this description one may deduce that the difference $T_\Gamma^{(m)}(z) - m(\hat{T}(m)T_\Gamma)(z)$ is holomorphic in $\mathbb{H}$ and vanishes as $z \rightarrow i\infty$. We thus obtain the remarkable expression

$$T_\Gamma^{(m)}(z) = m(\hat{T}(m)T_\Gamma)(z) \quad (1.7.5)$$

for the higher order Rademacher sums which generalizes the original result of Rademacher. Combining the presentation (1.2.1) of $T_\Gamma(z)$ as a Rademacher sum with the definition (1.7.4) of the Hecke operator $\hat{T}(m)$ we obtain the expression

$$m(\hat{T}(m)T_\Gamma)(z) + \frac{1}{2}c_\Gamma(m, 0) = e(-mz) + \lim_{K \rightarrow \infty} \sum_{\gamma \in (B(\mathbb{Z}) \backslash M(m))_{\leq K}^\times} e(-\gamma \cdot z) - e(-\gamma \cdot \infty), \quad (1.7.6)$$

where $M(m)$ denotes the (image in $PGL_2^+(\mathbb{Q})$ of the) set of 2×2 matrices with integral entries and determinant m . Comparison of (1.7.3) with (1.7.6), in view of the identity (1.7.5), suggests an equality of some, if not all, exponential terms in both sums, and in fact there is an injective map

$$\begin{aligned} B(\mathbb{Z}) \backslash M(1) &\hookrightarrow B(\mathbb{Z}) \backslash M(m) \\ \gamma &\mapsto \tilde{\gamma}, \end{aligned} \quad (1.7.7)$$

since $M(1) = \Gamma$ when $\Gamma = PSL_2(\mathbb{Z})$, with the property that $e(-m\gamma \cdot z) = e(-\tilde{\gamma} \cdot z)$. In the case that m is prime, the remaining terms assemble into a fractional power Rademacher sum

$$e\left(-\frac{1}{m}z\right) + \lim_{K \rightarrow \infty} \sum_{\gamma \in (B(m\mathbb{Z}) \backslash \Gamma)_{\leq K}^{\times}} e\left(-\frac{1}{m}\gamma \cdot z\right) - e\left(-\frac{1}{m}\gamma \cdot \infty\right), \quad (1.7.8)$$

where $B(m\mathbb{Z})$ is the m -fold translation group, generated by $z \mapsto z + m$. Thus we obtain another injective correspondence

$$\begin{aligned} B(m\mathbb{Z}) \backslash M(1) &\hookrightarrow B(\mathbb{Z}) \backslash M(m) \\ \gamma &\mapsto \tilde{\gamma} \end{aligned} \quad (1.7.9)$$

with the property that $e(-\gamma \cdot z/m) = e(-\tilde{\gamma} \cdot z)$. The identity (1.7.6) implies the vanishing of the fractional power Rademacher sum (1.7.8). This can also be proven directly, and we give a general vanishing result for fractional Rademacher sums in §5.4. More generally we obtain fractional power sums of the form

$$e\left(-\frac{m}{l^2}z\right) + \lim_{K \rightarrow \infty} \sum_{\gamma \in (B(l\mathbb{Z}) \backslash \Gamma)_{\leq K}^{\times}} e\left(-\frac{m}{l^2}\gamma \cdot z\right) - e\left(-\frac{m}{l^2}\gamma \cdot \infty\right), \quad (1.7.10)$$

for each exact divisor l of m , and each of these fractional power sums vanishes except for the sum corresponding to $l = 1$. Thus our analysis covers generalizations of the Rademacher sums to fractional orders in addition to the higher integral orders. All the results discussed in this section admit generalizations to arbitrary groups commensurable with the modular group.

1.8 Monstrous Lie algebras

The presentation (1.7.5) of the functions $T_{\Gamma}^{(m)}(z)$ by means of Hecke operators immediately implies that their Fourier coefficients are positive integers. This fact points to the existence of further algebraic structures beyond the moonshine module vertex operator algebra $V^{\natural}$. In his proof of the moonshine conjectures [Bor92] Borcherds introduced the Monster Lie algebra $\mathfrak{m}$, which admits a presentation as a bi-graded generalized Kac–Moody algebra. The key to Borcherds’s method is the denominator identity for $\mathfrak{m}$. This in turn can be interpreted as a BGG-type resolution of the trivial $\mathfrak{m}$ -module. Let $\mathcal{V}$ denote the Verma module with highest weight 0 for $\mathfrak{m}$. Then $\mathcal{V}$ is the first term in this BGG-type resolution of the trivial module, and the bi-graded dimension $\text{gdim } \mathcal{V}$ of $\mathcal{V}$ is obtained by computing the coefficients of p and q in the

expression

$$\mathrm{gdim} \mathcal{V} = \exp \left(\sum_{m \in \mathbb{Z}^+} (\hat{T}(m) T_\Gamma)(z) p^m \right) \quad (1.8.1)$$

where $q = e(z)$. Applying the operator $F \mapsto p \partial_p \log F$ to (1.8.1) we obtain a generating function for the higher order normalized Rademacher sums $T_\Gamma^{(m)}(z)$, by (1.7.5). It is well-known (cf. [Mac95]) that this operator relates the complete homogeneous symmetric functions to the power-sum symmetric functions. Thus the bi-graded dimension of the Verma module for the Monster Lie algebra may be viewed as the *complete Rademacher sum* (of weight 0) associated to Γ .

In §7.2 we consider a family of generalized Kac–Moody algebras $\{\mathfrak{m}_g\}$, indexed by elements g in the Monster group $\mathbb{M}$. Just as the Monster Lie algebra may be constructed from the moonshine module $V^\natural$, the monstrous Lie algebras $\mathfrak{m}_g$, studied by Carnahan in [Car07], may be constructed from the g -twisted $V^\natural$ -modules $V_g^\natural$, for $g \in \mathbb{M}$. The graded dimensions of the spaces $V_g^\natural$ are given by functions $J_g(z) = J_{\Gamma_g}(z)$, related to the McKay–Thompson series $T_g(z)$ via the involution $z \mapsto -1/z$.

$$J_g(z) = T_g(-1/z) \quad (1.8.2)$$

One should note that Borcherds set the precedent here, introducing a family $\{\mathfrak{m}'_g\}$ of monstrous Lie superalgebras in [Bor92]. These algebras $\mathfrak{m}'_g$ are more directly related to the functions $T_g(z)$, rather than the $J_g(z)$, and for our purposes the algebras $\mathfrak{m}_g$ appear to be more convenient.

The monstrous Lie algebras $\mathfrak{m}_g$ are constructed in [Car07] using the semi-infinite cohomology version of the no-ghost theorem. We identify the bi-graded subspaces of the algebra $\mathfrak{m}_g$ in terms of the twisted modules $V_h^\natural$, with $h \in \langle g \rangle$. One can show that the $\mathfrak{m}_g$ are generalized Kac–Moody algebras, and one can also deduce formulas for the bi-graded dimensions of their Verma modules $\mathcal{V}_g$, generalizing (1.8.1).

$$\mathrm{gdim} \mathcal{V}_g = \exp \left(\sum_{m \in \mathbb{Z}^+} (\hat{T}(m) T_g)(z) p^m \right) \quad (1.8.3)$$

Carnahan [Car07] also found a remarkable denominator identity which yields an alternative expression for the graded dimension of $\mathcal{V}_g$.

$$\mathrm{gdim} \mathcal{V}_g = \frac{1}{p(T_g(w) - J_g(z))} \quad (1.8.4)$$

This identity (1.8.4) demonstrates that the bi-graded dimension can be viewed as a meromorphic function on $\mathbb{H} \times \mathbb{H}$ with poles at $w \in \Gamma_g \cdot (-1/z)$.

We also show in §7.2 that the bi-graded dimension of $\mathcal{V}_g$ is bounded below, coefficient-wise,

by the generating function

$$\sum_{m \in \mathbb{Z}^+} J_g^{(m)}(z) p^m \quad (1.8.5)$$

when g is of Fricke type. This suggests that the Verma modules $\mathcal{V}_g$ may contain naturally defined subspaces whose bi-graded dimensions coincide with the expressions (1.8.5).

1.9 Chiral gravity

Our results on Rademacher sums, at first glance, add as much to the mystery of monstrous moonshine as they reveal. We have formulated a characterization of the groups of monstrous moonshine in terms of Rademacher sums and solid tori. Having done so we face the new question of where the Rademacher sums themselves appear in relation to the moonshine module vertex operator algebra $V^\natural$? It appears now that the relation to physics should again be useful, as it was in the construction of $V^\natural$ [FLM88], and the proof of the moonshine conjecture [Bor92].

In §7.1 we explain how the Rademacher sums associated to elements of the Monster might be regarded as giving strong evidence for the existence of another construction of the moonshine module $V^\natural$, and its twisted sectors $V_g^\natural$, in which all the features of monstrous moonshine, including the genus zero property, become transparent. In fact the original Rademacher sum (1.2.1) does appear as a saddle point approximation to the partition function of the simplest chiral three dimensional quantum gravity. (See [Wit07], [MW07], [Man07], [MM07], [LSS08] and [MSS09] for the development of this idea in the physics literature.) Our new identity (1.3.4) for the conjugate Rademacher sum must reflect the chiral nature of this three dimensional quantum gravity.

We expect that our analytic continuation of the Rademacher sums, and the explanation of the appearance of the constant term, will also develop from a rigorous analysis of the saddle point approximation using a zeta function regularization of the chiral gravity partition function, as was done for various models of two dimensional CFT. Our analysis of the Rademacher sums corresponding to the discrete groups of monstrous moonshine, and of moduli spaces of decorated solid tori points to a description of all the spaces $V_g^\natural$, for $g \in \mathbb{M}$, via a family of g -twisted versions of the simplest chiral three dimensional quantum gravity. We expect that our reformulation of monstrous moonshine in terms of Rademacher sums and moduli of solid tori will eventually be understood from the properties of this remarkable chiral three dimensional quantum gravity.

In §7.3 we also elucidate the relationship between the higher order Rademacher sums, and the key object of Borcherds' proof of the moonshine conjecture; viz., the Monster Lie algebra, and its twisted counterparts. This leads us to the stringy quantization (cf. [DMVV97]) of $V^\natural$, and we generalize this to all twisted sectors $V_g^\natural$, for $g \in \mathbb{M}$. The resulting second quantized spaces have the same bi-graded dimensions as the Verma modules $\mathcal{V}_g$ (cf. (1.8.3) and may be viewed as

stringy realizations of the representations of the generalized Kac–Moody algebras $\mathfrak{m}_g$. It is an interesting problem to interpret the spaces $\mathcal{V}_g$, and its subspaces with graded dimension given by the higher order normalized Rademacher sums (1.8.5), in terms of the second quantized chiral three dimensional quantum gravities. In particular, the simple description of the singularities of the partition function given by (1.8.4) should have a natural interpretation in terms of quantum gravity.

Thus our present results on the Rademacher sums, and the new conjectures about their origin, clearly indicate that the real nature of moonshine might not be as remote anymore, and more than this, the full examination of its structure might give new insight into the fundamental problem of modern physics.

2 Conventions

2.1 Functions

For $z, s \in \mathbb{C}$ with $z \neq 0$, we write z^s as a shorthand for $\exp(s \log(z))$, where $\log$ denotes the principal branch of the logarithm, so that

$$-\pi < \Im(\log(z)) \leq \pi, \quad (2.1.1)$$

and we write $z^{(s)}$ as a shorthand for $z^s / \Gamma(s+1)$, where $\Gamma(s)$ denotes the Gamma function. We adopt the convention of setting

$$e(z) = \exp(2\pi i z) = \sum_{k \in \mathbb{N}} (2\pi i z)^{(k)} \quad (2.1.2)$$

for $z \in \mathbb{C}$. We write $\Phi(a, b, z)$ for the analytic function on $\mathbb{C}^3$ defined by setting

$$\Phi(a, b, z) = \frac{{}_1F_1(a; b; 2\pi i z)}{\Gamma(b)} = \sum_{k \in \mathbb{N}} \frac{\Gamma(k+a)}{\Gamma(a)\Gamma(k+b)} (2\pi i z)^{(k)} \quad (2.1.3)$$

where ${}_1F_1(a; b; x)$ denotes the confluent hypergeometric function. Kummer's transformations for the confluent hypergeometric function yield the following identity for Φ .

$$\Phi(a, b, z) = e(z) \Phi(b-a, b, -z) \quad (2.1.4)$$

The exponential function $e(z)$ is a solution to the differential equation

$$(z\partial_z + 1 - s)(\partial_z - 2\pi i)u(z) = 0, \quad (2.1.5)$$

which has a regular singular point at 0 and an irregular singularity at ∞ . Another solution is furnished by the function $z \mapsto \Phi(1, 1 + s, z)(2\pi iz)^s$, which we denote also by $e(z, s)$.

$$e(z, s) = \Phi(1, 1 + s, z)(2\pi iz)^s = \sum_{k \in \mathbb{N}} (2\pi iz)^{(k+s)} \quad (2.1.6)$$

Observe that we have $e(z, n) = e(z) - e(z)_{<n}$ for $n \in \mathbb{Z}$, where $e(z)_{<K}$ denotes the *partial exponential* $e(z)_{<K} = \sum_{k \in \mathbb{N}, k < K} (2\pi iz)^{(k)}$. In particular, $e(z, n) = e(z)$ when $1 - n \in \mathbb{Z}^+$.

2.2 Divisors

For $n \in \mathbb{Z}^+$ and $s \in \mathbb{C}$ we write $\sigma(n, s)$ for the *divisor function*

$$\sigma(n, s) = \sum_{d|n} d^s. \quad (2.2.1)$$

For $a, b \in \mathbb{Z}$ we write (a, b) for the greatest common divisor of a and b , and we generalize this notation by writing $((a, b)$ for the greatest positive integer c say, such that $c^2|a$ and $c|b$, and by writing (a^∞, b) for the largest divisor of b that divides some power of a .

$$((a, b) = \max \{c \in \mathbb{Z}^+ \mid c^2|a \text{ and } c|b\} \quad (2.2.2)$$

$$(a^\infty, b) = \lim_{n \rightarrow \infty} (a^n, b) \quad (2.2.3)$$

For $d, n \in \mathbb{Z}^+$ we write $d||n$ in the case that $d|n$ and $(d, n/d) = 1$, and call such a d an *exact divisor* of n . We write $\text{Ex}(n)$ for the set of exact divisors of n . Then $\text{Ex}(n)$ becomes a group of exponent 2 when equipped with the product $(e, f) \mapsto ef/(e, f)^2$. Indeed, if we write $\Pi(n)$ for the set of primes that divide n then there is a naturally defined isomorphism $\text{Ex}(n) \cong (\mathbb{Z}/2)^{\Pi(n)}$ which associates to an exact divisor $e \in \text{Ex}(n)$ the indicator function $\chi_e : \Pi(n) \rightarrow \mathbb{Z}/2$, satisfying $\chi_e(p) = 1$ if $p|e$ and $\chi_e(p) = 0$ otherwise.

Let Π denote the set of all primes. Then the infinite product $(\mathbb{Z}/2)^\Pi$ comes equipped with a naturally defined projection $(\mathbb{Z}/2)^\Pi \rightarrow (\mathbb{Z}/2)^{\Pi(n)}$, and a naturally defined section for it $(\mathbb{Z}/2)^{\Pi(n)} \rightarrow (\mathbb{Z}/2)^\Pi$, for each $n \in \mathbb{Z}^+$. Consequently we have naturally defined morphisms of groups $(\mathbb{Z}/2)^\Pi \rightarrow \text{Ex}(n)$ and $\text{Ex}(n) \rightarrow (\mathbb{Z}/2)^\Pi$ for each n , and for any $m, n \in \mathbb{Z}^+$ we have the natural map $\text{Ex}(m) \rightarrow \text{Ex}(n)$ furnished by the composition $\text{Ex}(m) \rightarrow (\mathbb{Z}/2)^\Pi \rightarrow \text{Ex}(n)$. If m divides n then the map $\text{Ex}(m) \rightarrow \text{Ex}(n)$ is injective, the map $\text{Ex}(n) \rightarrow \text{Ex}(m)$ is surjective, and the composition $\text{Ex}(m) \rightarrow \text{Ex}(n) \rightarrow \text{Ex}(m)$ is the identity on $\text{Ex}(m)$.

2.3 Isometries

Let us write $G(\mathbb{R})$ for the simple real Lie group $PSL_2(\mathbb{R})$. The group $G(\mathbb{R})$ acts naturally, from the left, on the upper-half plane $\mathbb{H} = \{z \in \mathbb{C} \mid \Im(z) > 0\}$, and is just the group of orientation preserving isometries of $\mathbb{H}$ when we equip it with the *hyperbolic measure* $d\mu(z) = dx dy / y^2$, for $z = x + iy$.

It is convenient to enlist matrices in $SL_2(\mathbb{R})$ for the purpose of specifying elements of $G(\mathbb{R})$. We write $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$ for the image of a matrix $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbb{R})$ in $G(\mathbb{R})$. Analogously, we write $A \mapsto [A]$ for the canonical map $SL_2(\mathbb{R}) \rightarrow G(\mathbb{R})$. The action of $G(\mathbb{R})$ on $\mathbb{H}$ is now described explicitly by the formula

$$\gamma \cdot z = \frac{az + b}{cz + d} \quad (2.3.1)$$

for $\gamma = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. The matrix $\theta = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} \in GL_2(\mathbb{R})$ induces an outer-automorphism of $G(\mathbb{R})$, which we call *conjugation on $G(\mathbb{R})$* , and which we denote $\gamma \mapsto \bar{\gamma}$. Explicitly, we have

$$\bar{\gamma} = \begin{bmatrix} -a & b \\ c & -d \end{bmatrix} \Longleftarrow \gamma = \begin{bmatrix} a & b \\ c & d \end{bmatrix}. \quad (2.3.2)$$

For each matrix $A \in GL_2^+(\mathbb{Q})$ there is a unique $\mu \in \mathbb{R}^+$ for which $A' = \mu A$ belongs to $SL_2(\mathbb{R})$. The assignment $GL_2^+(\mathbb{Q}) \rightarrow G(\mathbb{R})$ given by $A \mapsto [A']$ then factors through $PGL_2^+(\mathbb{Q})$, and the resulting map $PGL_2^+(\mathbb{Q}) \rightarrow G(\mathbb{R})$ is an embedding of groups. We write $G(\mathbb{Q})$ for the copy of $PGL_2^+(\mathbb{Q})$ in $G(\mathbb{R})$ obtained in this way. We write $G(\mathbb{Z})$ for the *modular group* $PSL_2(\mathbb{Z})$.

Extending the notation introduced above, we write $[A]$ and $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$ for the image in $G(\mathbb{Q}) < G(\mathbb{R})$ of a matrix $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ in $GL_2^+(\mathbb{Q})$. Given $\mu \in \mathbb{Q}^+$ we write $[\mu]$ as a shorthand for $[A]$ when A is the diagonal matrix $\begin{pmatrix} \mu & 0 \\ 0 & 1 \end{pmatrix}$. Then $[\mu] \cdot z = \mu z$ for $z \in \mathbb{H}$.

$$[\mu] = \begin{bmatrix} \mu & 0 \\ 0 & 1 \end{bmatrix} \quad (2.3.3)$$

Let $B(\mathbb{R})$ denote the Borel subgroup of $G(\mathbb{R})$ consisting of images of upper-triangular matrices in $SL_2(\mathbb{R})$, so that $B(\mathbb{R})$ is just the subgroup of $G(\mathbb{R})$ that fixes the distinguished point ∞ on the *pointed real projective line* $\hat{\mathbb{R}} = \mathbb{R} \cup \{\infty\}$. Let $B(\mathbb{Q})$ and $B(\mathbb{Z})$ denote the intersections $B(\mathbb{R}) \cap G(\mathbb{Q})$ and $B(\mathbb{R}) \cap G(\mathbb{Z})$, respectively. Then $B(\mathbb{Z})$ is the group generated by the *modular translation* $T = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$. Set $\hat{\mathbb{Q}} = \mathbb{Q} \cup \{\infty\} \subset \hat{\mathbb{R}}$. Then $G(\mathbb{Q})$ acts naturally on $\hat{\mathbb{Q}}$, and $B(\mathbb{Q})$ is just the subgroup of $G(\mathbb{Q})$ that fixes the point ∞ . For $\alpha \in \mathbb{Q}$ we write T^α for the element

$$\begin{bmatrix} 1 & \alpha \\ 0 & 1 \end{bmatrix} \in B(\mathbb{Q}).$$

$$T^\alpha = \begin{bmatrix} 1 & \alpha \\ 0 & 1 \end{bmatrix} \quad (2.3.4)$$

Then the elements of $B(\mathbb{Q})$ of the form T^α for $\alpha \in \mathbb{Q}$ constitute the *unipotent subgroup* of $B(\mathbb{Q})$, which we denote $B_u(\mathbb{Q})$. Given $\alpha \in \mathbb{Q}^+$ we write $B(\alpha\mathbb{Z})$ for the subgroup of $B_u(\mathbb{Q})$ generated by T^α .

$$B(\alpha\mathbb{Z}) = \left\{ \begin{bmatrix} 1 & \alpha n \\ 0 & 1 \end{bmatrix} \mid n \in \mathbb{Z} \right\} \quad (2.3.5)$$

The elements of $B(\mathbb{Q})$ of the form $[\mu]$ for $\mu \in \mathbb{Q}^+$ constitute the *diagonal subgroup* of $B(\mathbb{Q})$, which we denote $B_d(\mathbb{Q})$. The group $B(\mathbb{Q})$ is naturally isomorphic to the semidirect product $B_u(\mathbb{Q}) \rtimes B_d(\mathbb{Q})$. In particular, $B_u(\mathbb{Q})$ is a normal subgroup of $B(\mathbb{Q})$ and for any $\chi \in B(\mathbb{Q})$ we have $\chi = T^\alpha[\mu]$ for some uniquely determined $\alpha \in \mathbb{Q}$ and $\mu \in \mathbb{Q}^+$.

An element $\gamma \in G(\mathbb{Q})$ will have many preimages in $GL_2^+(\mathbb{Q})$, but among these there will be exactly two that have integral entries with no common divisor. Either of these two matrices will be called a *preferred representative* for γ . We define functions $Pdet : G(\mathbb{Q}) \rightarrow \mathbb{Z}^+$ and $c : G(\mathbb{Q}) \rightarrow \mathbb{N}$ and $d : G(\mathbb{Q}) \rightarrow \mathbb{N}$ by setting

$$Pdet(\gamma) = ad - bc, \quad c(\gamma) = |c|, \quad d(\gamma) = |d|, \quad (2.3.6)$$

in case $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in GL_2^+(\mathbb{Q})$ is a preferred representative for γ . We call $Pdet$ the *projective determinant*. The group $G(\mathbb{Z})$ is exactly the preimage of 1 with respect to the projective determinant. The projective determinant is not multiplicative, but we have $Pdet(\gamma\sigma) = Pdet(\sigma) = Pdet(\sigma\gamma)$ for all $\sigma \in G(\mathbb{Q})$ in case $Pdet(\gamma) = 1$, so for any $n \in \mathbb{Z}^+$, the preimage of n under $Pdet$ is naturally both a left and right $G(\mathbb{Z})$ -set.

Observe that $c(T\gamma) = c(\gamma T) = c(\gamma)$ and $d(T\gamma) = d(\gamma)$. Given $X \subset G(\mathbb{Q})$ we set $X_\infty = X \cap B(\mathbb{Q})$ and $X^\times = X - X_\infty$. Then $X_\infty = \{\chi \in X \mid c(\chi) = 0\}$. We define $X^{\times\times}$ to be the subset of $X^\times$ consisting of $\chi \in X$ such that both $c(\chi)$ and $d(\chi)$ are non-zero.

$$X_\infty = \{\chi \in X \mid c(\chi) = 0\}, \quad (2.3.7)$$

$$X^\times = \{\chi \in X \mid c(\chi) \neq 0\}, \quad X^{\times\times} = \{\chi \in X \mid c(\chi)d(\chi) \neq 0\} \quad (2.3.8)$$

For $\gamma \in G(\mathbb{Q})$ with $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ a preferred representative, we have

$$\gamma \cdot z - \gamma \cdot z' = \frac{Pdet(A)}{(cz + d)(cz' + d)}(z - z') \quad (2.3.9)$$

for $z, z' \in \mathbb{H}$, so that the derivative of the function $z \mapsto \gamma \cdot z$, which we will denote $\text{jac}(\gamma, z)$, is given by

$$\text{jac}(\gamma, z) = \frac{P\det(A)}{(cz + d)^2}. \quad (2.3.10)$$

Note that the assignment $\text{jac} : G(\mathbb{Q}) \rightarrow \mathcal{O}(\mathbb{H})$, which associates the function $z \mapsto \text{jac}(\gamma, z)$ to an element $\gamma \in G(\mathbb{Q})$, descends naturally to the coset space $B(\mathbb{Z}) \backslash G(\mathbb{Q})$, and even further to $B_u(\mathbb{Q}) \backslash G(\mathbb{Q})$.

From (2.3.9) and (2.3.10) we see that $|\gamma \cdot z - \gamma \cdot z'| = |z - z'|$ just in the case that z and z' belong to the set $\{z \in \mathbb{H} \mid |\text{jac}(\gamma, z)| = 1\}$ which we call the *isometric locus* of γ . The isometric locus of γ is a (Euclidean) semicircle just when $c(\gamma) > 0$, in which case its center is $\gamma^{-1} \cdot \infty$, and its radius is $\sqrt{\varrho(\gamma)}$, where

$$\varrho(\gamma) = \frac{P\det(A)}{c(\gamma)^2} \quad (2.3.11)$$

for A a preferred representative for γ . We call $\varrho(\gamma)$ the *scaling factor* associated to γ .

2.4 Cosets

Given a subset $X \subset G(\mathbb{Q})$, we will write $\llbracket X \rrbracket$ for the set consisting of double cosets of the form $B(\mathbb{Z})\chi B(\mathbb{Z})$, for $\chi \in X$. We will write $\llbracket \chi \rrbracket$ as a shorthand for the particular double coset $B(\mathbb{Z})\chi B(\mathbb{Z})$.

$$\llbracket \chi \rrbracket = B(\mathbb{Z})\chi B(\mathbb{Z}), \quad \llbracket X \rrbracket = \{\llbracket \chi \rrbracket \in B(\mathbb{Z}) \backslash G(\mathbb{Q}) / B(\mathbb{Z}) \mid \chi \in X\} \quad (2.4.1)$$

Similarly, we will write $\llbracket X \rrbracket$ for the set of right cosets of the form $B(\mathbb{Z})\chi$, for $\chi \in X$, and $\chi \mapsto \llbracket \chi \rrbracket$ will denote the natural map $X \rightarrow \llbracket X \rrbracket$, and the notations $\llbracket X \rrbracket$ and $\chi \mapsto \llbracket \chi \rrbracket$ will have the analogous meanings. Observe that $B(\mathbb{Z})$ is stable under the operations of inversion and conjugation (cf. (2.3.2)), and thus these operations descend naturally to the coset spaces $\llbracket G(\mathbb{Q}) \rrbracket$ and $G(\mathbb{Q}) \rrbracket$ and $\llbracket G(\mathbb{Q}) \rrbracket$.

Recall from §2.3 that the assignment $\chi \mapsto \text{jac}(\chi, z)$ descends to a well-defined association of holomorphic functions on $\mathbb{H}$ to right cosets of $B(\mathbb{Z})$ in $G(\mathbb{Q})$. Accordingly, we may safely set $\text{jac}(\llbracket \chi \rrbracket, z) = \text{jac}(\chi, z)$, when given some $\llbracket \chi \rrbracket \in \llbracket G(\mathbb{Q}) \rrbracket$.

$$\begin{aligned} \text{jac}(\cdot, z) : \llbracket G(\mathbb{Q}) \rrbracket &\rightarrow \mathcal{O}(\mathbb{H}) \\ \llbracket \chi \rrbracket &\mapsto \text{jac}(\llbracket \chi \rrbracket, z) \end{aligned} \quad (2.4.2)$$

Given $U \subset \llbracket G(\mathbb{Q}) \rrbracket$ or $S \subset \llbracket G(\mathbb{Q}) \rrbracket$ we set $U_\infty = U \cap \llbracket B(\mathbb{Q}) \rrbracket$ and $U^\times = U - U_\infty$, and similarly, $S_\infty = S \cap \llbracket B(\mathbb{Q}) \rrbracket$ and $S^\times = S - S_\infty$. In particular, $\llbracket G(\mathbb{Q}) \rrbracket^\times$ denotes the set of double cosets

$\llbracket \chi \rrbracket$, for $\chi \in G(\mathbb{Q})$, for which $c(\chi) \neq 0$.

Observe that the isometric loci (cf. §2.3) associated to $\chi, \chi' \in G(\mathbb{Q})$ coincide when $\llbracket \chi \rrbracket = \llbracket \chi' \rrbracket$. Thus we may regard these loci as naturally attached to right cosets of $B(\mathbb{Z})$ in $G(\mathbb{Q})$, and the locus attached to $\llbracket \chi \rrbracket$ is a Euclidean semicircle just when $\llbracket \chi \rrbracket \in \llbracket G(\mathbb{Q}) \rrbracket^\times$. Also, the loci associated to cosets $\llbracket \chi \rrbracket, \llbracket \chi' \rrbracket \in \llbracket G(\mathbb{Q}) \rrbracket^\times$ have the same radii when $\llbracket \chi \rrbracket = \llbracket \chi' \rrbracket$; that is to say, the function ϱ of (2.3.11) descends to a well-defined map $\llbracket G(\mathbb{Q}) \rrbracket^\times \rightarrow \mathbb{Q}^+$. It is worth noting that the function $\chi \mapsto \varrho(\chi)$ actually satisfies an even stronger invariance condition.

Lemma 2.4.1. *The assignment $\llbracket \chi \rrbracket \mapsto \varrho \llbracket \chi \rrbracket$ descends naturally to a well-defined function on the double coset space $B_u(\mathbb{Q}) \backslash G(\mathbb{Q}) / B_u(\mathbb{Q})$. That is, we have $\varrho \llbracket T^\alpha \chi \rrbracket = \varrho \llbracket \chi T^\alpha \rrbracket = \varrho \llbracket \chi \rrbracket$ for any $\alpha \in \mathbb{Q}$ and $\chi \in G(\mathbb{Q})^\times$.*

The function ϱ is also invariant under inversion and conjugation, so that we have $\varrho \llbracket \chi \rrbracket = \varrho \llbracket \bar{\chi} \rrbracket = \varrho \llbracket \chi^{-1} \rrbracket$ for any $\chi \in G(\mathbb{Q})^\times$. For $\chi \in G(\mathbb{Q})^\times$ the identity (2.3.9) may be rewritten

$$\chi \cdot z - \chi \cdot z' = \frac{\varrho \llbracket \chi \rrbracket}{(z - \chi^{-1} \cdot \infty)(z' - \chi^{-1} \cdot \infty)} (z - z'), \quad (2.4.3)$$

and we may consider the limit as $z' \rightarrow \infty$, which yields a useful expression for $\chi \cdot z - \chi \cdot \infty$; viz.,

$$\chi \cdot z - \chi \cdot \infty = -\frac{\varrho \llbracket \chi \rrbracket}{z - \chi^{-1} \cdot \infty}. \quad (2.4.4)$$

Comparing with (2.3.10) and (2.4.4) we have

$$\text{jac}(\llbracket \chi \rrbracket, z) = \frac{\varrho \llbracket \chi \rrbracket}{(z - \chi^{-1} \cdot \infty)^2} \quad (2.4.5)$$

for $\llbracket \chi \rrbracket \in \llbracket G(\mathbb{Q}) \rrbracket^\times$ (and $\text{jac}(\llbracket \chi \rrbracket, z) = 1$ otherwise).

The functions $c, d : G(\mathbb{Q}) \rightarrow \mathbb{N}$, of §2.3, descend to well-defined functions on the coset space $\llbracket G(\mathbb{Q}) \rrbracket = B(\mathbb{Z}) \backslash G(\mathbb{Q})$. The function c descends further, to be well-defined on the double coset space $\llbracket G(\mathbb{Q}) \rrbracket = B(\mathbb{Z}) \backslash G(\mathbb{Q}) / B(\mathbb{Q})$. For $K \in \mathbb{R}^+$ and U a subset of $\llbracket G(\mathbb{Q}) \rrbracket$, define $U_{\leq K}$ to be the subset of U consisting of cosets $\llbracket \chi \rrbracket \in U$ for which $c(\chi) \leq K$ and $d(\chi) \leq K^2$. Analogously, for S a subset of $\llbracket G(\mathbb{Q}) \rrbracket$, define $S_{\leq K}$ to be the subset of S consisting of double cosets $\llbracket \chi \rrbracket \in S$ for which $c(\chi) \leq K$.

$$U_{\leq K} = \{ \llbracket \chi \rrbracket \in U \mid c(\chi) \leq K, d(\chi) \leq K^2 \} \quad (2.4.6)$$

$$S_{\leq K} = \{ \llbracket \chi \rrbracket \in S \mid c(\chi) \leq K \} \quad (2.4.7)$$

2.5 Groups

The group $G(\mathbb{Q})$ is the commensurator of $G(\mathbb{Z})$. Consequently, any group $\Gamma < G(\mathbb{R})$ that is commensurable with the modular group $G(\mathbb{Z})$ is automatically a subgroup of $G(\mathbb{Q})$, and is an example of a Fuchsian group of the first kind. For such a group Γ there is a natural way to equip the orbit space $\Gamma \backslash \mathbb{H}$ with the structure of a Riemann surface; we will denote this object by Y_Γ . Any group Γ commensurable with $G(\mathbb{Z})$ has parabolic elements, and any fixed point of any parabolic element of Γ lies on the (pointed) rational projective line $\hat{\mathbb{Q}} \subset \hat{\mathbb{R}}$. The orbit space $P_\Gamma = \Gamma \backslash \hat{\mathbb{Q}}$ is the set of *cusps* of Γ . For Γ commensurable with $G(\mathbb{Z})$, the Riemann surface Y_Γ is not compact, but can be compactified in a natural way by adjoining a single point for each cusp of Γ . We will write X_Γ for the corresponding compact Riemann surface obtained by the adjunction of the cusps of Γ .

$$X_\Gamma = Y_\Gamma \cup P_\Gamma = \Gamma \backslash \mathbb{H} \cup \Gamma \backslash \hat{\mathbb{Q}} \quad (2.5.1)$$

We will say that Γ is a *group of genus zero* in the case that X_Γ has genus zero as a Riemann surface. We set $\text{genus}(\Gamma) = \text{genus}(X_\Gamma)$ for $\Gamma < G(\mathbb{R})$ commensurable with $G(\mathbb{Z})$.

We write $\hat{\mathbb{C}}$ for the (*pointed*) *complex projective line* $\mathbb{C} \cup \{\infty\}$, which we regard as a compact Riemann surface in the usual way. If Γ is a group for which the compact Riemann surface X_Γ has genus zero, then there must be an isomorphism $\phi : X_\Gamma \xrightarrow{\sim} \hat{\mathbb{C}}$ that witnesses this fact. After applying a Möbius transformation if necessary, we may arrange that ϕ maps the point of X_Γ corresponding to the *infinite cusp* $\Gamma \cdot \infty$ to the distinguished point $\infty \in \hat{\mathbb{C}}$. Such an isomorphism ϕ determines a Γ -invariant holomorphic function, f say, on $\mathbb{H}$, which admits an expression $f(z) = \sum_{n \geq -1} c(n) e(nz)$ with $c(-1) \neq 0$. After multiplying by $1/c(-1)$ we may assume that $c(-1) = 1$. A Γ -invariant function of the form $e(-z) + \sum_{n \geq 0} c(n) e(nz)$ which is holomorphic on $\mathbb{H}$ and induces an isomorphism $X_\Gamma \xrightarrow{\sim} \hat{\mathbb{C}}$ will be called a *hauptmodul* for Γ . In case $c(0) = 0$ we say that f is a *normalized hauptmodul* for Γ .

The *Hecke congruence groups*, denoted $\Gamma_0(n)$ for $n \in \mathbb{Z}^+$, play a special rôle in our analysis.

$$\Gamma_0(n) = \left\{ \begin{bmatrix} a & b \\ cn & d \end{bmatrix} \mid a, b, c, d \in \mathbb{Z}, ad - bcn = 1 \right\} \quad (2.5.2)$$

According to [CN79] the normalizer $N(\Gamma_0(n))$ of $\Gamma_0(n)$ in $G(\mathbb{R})$ is commensurable with $G(\mathbb{Z})$ and admits the description

$$N(\Gamma_0(n)) = \left\{ \begin{bmatrix} ae & b/h \\ cn & de \end{bmatrix} \mid a, b, c, d \in \mathbb{Z}, e \in \mathbb{Z}^+, e \parallel n/h, ade - bcn/eh = 1 \right\} \quad (2.5.3)$$

where $h = (n, 24)$ is the largest divisor of 24 such that h^2 divides n (cf. §2.2). The expression

(2.5.3) tells us, in particular, that $N(\Gamma_0(n))_\infty = \langle T^{1/h} \rangle$ for $h = ((n, 24))$.

Proposition 2.5.1. *Let $n \in \mathbb{Z}^+$. Then $N(\Gamma_0(n))$ acts transitively on $\hat{\mathbb{Q}}$. We have $N(\Gamma_0(n))_\infty = \Gamma_0(n)_\infty$ if and only if n is not divisible by 4 or 9.*

An important family of groups, each one commensurable with the modular group $G(\mathbb{Z})$, and each one containing and normalizing some $\Gamma_0(n)$, was introduced in [CN79]; these are the groups of $n||h$ -type, and we now recall their definition. For $n, h \in \mathbb{Z}^+$ with $h|n$, and for S a subgroup of $\text{Ex}(n/h)$ (cf. §2.2), we define a group $\Gamma_0(n|h) + S$ by setting

$$\Gamma_0(n|h) + S = \left\{ \begin{bmatrix} ae & b/h \\ cn & de \end{bmatrix} \mid a, b, c, d \in \mathbb{Z}, e \in S, ade - bcn/eh = 1 \right\}. \quad (2.5.4)$$

Evidently the group $\Gamma_0(n|h) + S$ contains $\Gamma_0(nh)$. In case $h|24$ it also normalizes $\Gamma_0(nh)$. Assume then that $h|24$. We say that $\Gamma_0(n|h) + S$ is of *Fricke type* if the element $\begin{bmatrix} 0 & -1 \\ nh & 0 \end{bmatrix}$ belongs to it.

The group $\Gamma_0(n||h) + S$ is defined (cf. [CN79]) as the subgroup of $\Gamma_0(n|h) + S$ arising as the kernel of a certain morphism $\Gamma_0(n|h) + S \rightarrow \mathbb{Z}/h$ which factors through the canonical map $\Gamma_0(n|h) + S \rightarrow (\Gamma_0(n|h) + S)/\Gamma_0(nh)$. In order to describe it let S' denote the image of S in $\text{Ex}(nh)$ under the natural injection $\text{Ex}(n/h) \rightarrow \text{Ex}(nh)$ (cf. §2.2). Then as generators for the quotient $(\Gamma_0(n|h) + S)/\Gamma_0(nh)$ we may take the cosets X , Y , and $W_{e'}$ for $e' \in S'$, given by

$$X = \begin{bmatrix} 1 & 1/h \\ 0 & 1 \end{bmatrix} \Gamma_0(nh), \quad Y = \begin{bmatrix} 1 & 0 \\ n & 1 \end{bmatrix} \Gamma_0(nh), \quad W_{e'} = \begin{bmatrix} ae' & b \\ cnh & de' \end{bmatrix} \Gamma_0(nh), \quad (2.5.5)$$

where the a, b, c and d in the definition of $W_{e'}$ are arbitrary integers for which $ade' - bcnh/e' = 1$, and we may define $\Gamma_0(n||h) + S$ to be the kernel of the composition

$$\Gamma_0(n|h) + S \rightarrow (\Gamma_0(n|h) + S)/\Gamma_0(nh) \xrightarrow{\lambda} \mathbb{Z}/h \quad (2.5.6)$$

where the map λ is determined by the requirements that $\lambda(W_{e'}) = 0$ for all $e' \in S'$, $\lambda(X) = 1$, and $\lambda(Y)$ is 1 or -1 according as $\Gamma_0(n|h) + S$ is of Fricke type or not (cf. [Fer93]).

2.6 Scalings

Let Γ be a group commensurable with $G(\mathbb{Z})$ and let $\mathfrak{p} \in \mathbf{P}_\Gamma$ (cf. (2.5.1)) be a cusp of Γ . An element $p \in \mathfrak{p} \subset \hat{\mathbb{Q}}$ will be called a *representative* for $\mathfrak{p}$.

Lemma 2.6.1. *Let Γ be a group commensurable with $G(\mathbb{Z})$ and let $p \in \hat{\mathbb{Q}}$. Then there exists an element $\sigma_p \in G(\mathbb{Q})$ such that*

$$p = \sigma_p \cdot \infty, \quad (\sigma_p^{-1} \Gamma \sigma_p)_\infty = B(\mathbb{Z}). \quad (2.6.1)$$

Proof. In case $p = \infty$ the group Γ_∞ (cf. (2.3.7)) of elements in Γ that fix ∞ is infinite cyclic and generated by some element $\gamma_\infty \in B(\mathbb{Q})$. We claim that in fact γ_∞ lies in $B_u(\mathbb{Q})$, and thus equals T^α (cf. (2.3.4)) for some $\alpha \in \mathbb{Q}^+$. For otherwise $P\det(\gamma_\infty) = e$ for some $e > 1$ (cf. (2.3.6)), so that $P\det(\gamma_\infty^n) = e^n$ for $n \in \mathbb{Z}$, since the restriction $P\det : B(\mathbb{Q}) \rightarrow \mathbb{Z}^+$ is multiplicative. Then each power of γ_∞ lies in a distinct coset of the intersection $\Gamma \cap G(\mathbb{Z})$ in Γ since $P\det$ is invariant under multiplication by elements of $G(\mathbb{Z})$. This contradicts the hypothesis that Γ is commensurable with $G(\mathbb{Z})$, so we conclude that $\Gamma_\infty = \langle T^\alpha \rangle$ for some $\alpha \in \mathbb{Q}^+$. Then we may take $\sigma_\infty = [\alpha]$ (cf. (2.3.3)), for upon calculating $[\mu]T^\alpha[1/\mu] = T^{\mu\alpha}$ we find that

$$([1/\alpha]\Gamma[\alpha])_\infty = [1/\alpha]\Gamma_\infty[\alpha] = [1/\alpha]\langle T^\alpha \rangle[\alpha] = \langle T \rangle, \quad (2.6.2)$$

and the group $\langle T \rangle$ is just $B(\mathbb{Z})$, so the element $\sigma_\infty = [\alpha]$ satisfies the two conditions (2.6.1).

In case $p = a/c$ for coprime integers $a, c \in \mathbb{Z}$ with $c \neq 0$, we may choose $b, d \in \mathbb{Z}$ such that $ad - bc = 1$ and set $\sigma = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Then $p = \sigma \cdot \infty$ and σ satisfies the first condition in (2.6.1). Now we rerun the argument of the paragraph above with $\sigma^{-1}\Gamma\sigma$ in place of Γ to find that $(\sigma^{-1}\Gamma\sigma)_\infty = \langle T^\alpha \rangle$ for some $\alpha \in \mathbb{Q}^+$, and $\sigma_p = \sigma[\alpha]$ satisfies the two conditions (2.6.1). This completes the proof of the lemma. $\square$

An element $\sigma_p \in G(\mathbb{Q})$ satisfying the two conditions (2.6.1) of Lemma 2.6.1 will be called a *scaling element for Γ at the cusp representative p* . It is useful to have a replacement for the notion of scaling element that is independent of a choice of cusp representative.

Lemma 2.6.2. *Let Γ be a group commensurable with $G(\mathbb{Z})$ and let $\mathfrak{p} \in \mathcal{P}_\Gamma$ be a cusp of Γ . Then there exists a coset $\Sigma_{\mathfrak{p}} \in \Gamma \backslash G(\mathbb{Q})$ such that*

$$\mathfrak{p} = \Sigma_{\mathfrak{p}} \cdot \infty, \quad (\Sigma_{\mathfrak{p}}^{-1}\Sigma_{\mathfrak{p}})_\infty = B(\mathbb{Z}). \quad (2.6.3)$$

Proof. Let $p \in \mathfrak{p} \subset \hat{\mathbb{Q}}$ be a representative for $\mathfrak{p}$ and let $\sigma_p \in G(\mathbb{Q})$ satisfy the conditions (2.6.1) of Lemma 2.6.1. Then the coset $\Sigma_{\mathfrak{p}} = \Gamma\sigma_p$ satisfies the conditions (2.6.3). $\square$

A coset $\Sigma_{\mathfrak{p}} \in \Gamma \backslash G(\mathbb{Q})$ satisfying the two conditions (2.6.3) of Lemma 2.6.2 will be called a *scaling coset for Γ at the cusp $\mathfrak{p}$* . Any element $\sigma \in \Sigma_{\mathfrak{p}}$ is a scaling element for Γ at the representative $\sigma \cdot \infty$ for the cusp $\mathfrak{p}$, and if σ_p is a scaling element for Γ at the cusp representative p then the coset $\Gamma\sigma_p$ is a scaling coset for Γ at the cusp $\mathfrak{p} = \Gamma \cdot p$ represented by p .

Scaling cosets are determined only up to right multiplication by elements of $B_u(\mathbb{Q})$. Indeed, both the conditions defining the notion of scaling coset are invariant under the replacement of $\Sigma_{\mathfrak{p}}$ by $\Sigma_{\mathfrak{p}}T^\alpha$ for some $\alpha \in \mathbb{Q}$ (cf. (2.3.4)). A set $\{\Sigma_{\mathfrak{p}} \mid \mathfrak{p} \in \mathcal{P}_\Gamma\} \subset \Gamma \backslash G(\mathbb{Q})$ such that $\Sigma_{\mathfrak{p}}$ is a scaling coset for Γ at $\mathfrak{p}$ for each $\mathfrak{p} \in \mathcal{P}_\Gamma$ will be called a *system of scaling cosets for Γ* .

The problem of constructing scaling cosets for a group Γ may be viewed in the following way. The group $G(\mathbb{Q})$ acts transitively on $\hat{\mathbb{Q}}$, and the subgroup $B(\mathbb{Q})$ is just the stabilizer of

the distinguished point $\infty \in \hat{\mathbb{Q}}$. Thus the map $\Gamma \backslash G(\mathbb{Q}) \rightarrow \mathbf{P}_\Gamma$ given by $\Gamma\sigma \mapsto \Gamma\sigma \cdot \infty$ induces an isomorphism $\Gamma \backslash G(\mathbb{Q})/B(\mathbb{Q}) \cong \mathbf{P}_\Gamma$ which we may regard as identifying the sets $\Gamma \backslash G(\mathbb{Q})/B(\mathbb{Q})$ and $\mathbf{P}_\Gamma$. Now $B(\mathbb{Q})$ is naturally isomorphic to the semidirect product $B_u(\mathbb{Q}) \rtimes B_d(\mathbb{Q})$ (cf. §2.5), so the natural map $\Gamma \backslash G(\mathbb{Q}) \rightarrow \Gamma \backslash G(\mathbb{Q})/B(\mathbb{Q})$ factors through $\Gamma \backslash G(\mathbb{Q})/B_u(\mathbb{Q})$, yielding a naturally defined sequence

$$\Gamma \backslash G(\mathbb{Q}) \rightarrow \Gamma \backslash G(\mathbb{Q})/B_u(\mathbb{Q}) \rightarrow \Gamma \backslash G(\mathbb{Q})/B(\mathbb{Q}) \cong \mathbf{P}_\Gamma \quad (2.6.4)$$

where the fibres of the second map are torsors for the diagonal group $B_d(\mathbb{Q}) \simeq \mathbb{Q}^+$. Given a cusp $\mathfrak{p} \in \mathbf{P}_\Gamma$, any preimage of $\mathfrak{p}$ in $\Gamma \backslash G(\mathbb{Q})$ under the composition (2.6.4) is a coset of Γ satisfying the first condition defining a scaling coset for Γ at $\mathfrak{p}$. In order to satisfy also the second condition we should multiply this coset by $[\mu]$ (cf. (2.3.3)) for some (uniquely defined) $\mu \in \mathbb{Q}^+$. Thus the two conditions defining scaling cosets define a section of the $B_d(\mathbb{Q})$ -bundle $\Gamma \backslash G(\mathbb{Q})/B_u(\mathbb{Q}) \rightarrow \mathbf{P}_\Gamma$. We denote this map $\mathfrak{p} \mapsto \mathfrak{S}_\mathfrak{p}$. We may naturally identify $\mathfrak{S}_\mathfrak{p}$ with the set of scaling cosets for Γ at $\mathfrak{p}$, for these are precisely the preimages of $\mathfrak{S}_\mathfrak{p}$ in $\Gamma \backslash G(\mathbb{Q})$ under the first map of (2.6.4).

$$\begin{aligned} \mathbf{P}_\Gamma &\rightarrow \Gamma \backslash G(\mathbb{Q})/B_u(\mathbb{Q}) \\ \mathfrak{p} &\mapsto \mathfrak{S}_\mathfrak{p} = \{\Sigma_\mathfrak{p} T^\alpha \mid \alpha \in \mathbb{Q}\} \end{aligned} \quad (2.6.5)$$

One says that Γ has *width one at infinity* in the case that $\Gamma_\infty = B(\mathbb{Z})$. Observe that Γ has width one at infinity if and only if Γ is a scaling coset for itself at the infinite cusp $\Gamma \cdot \infty$; that is, if and only if $\Gamma \in \mathfrak{S}_{\Gamma \cdot \infty}$. If Γ does not have width one at infinity then there is a unique $\mu \in \mathbb{Q}^+$ with the property that $\Gamma[\mu]$ is a scaling coset for Γ at the infinite cusp $\Gamma \cdot \infty$, and then the group $\Gamma^{[\mu]} = [1/\mu]\Gamma[\mu]$ is a group with width one at infinity.

When engaged in the task of computing Fourier coefficients of modular forms for a group Γ say, one frequently has use for double coset spaces of the form $\Gamma_p \backslash \Gamma / \Gamma_q$, for some $p, q \in \hat{\mathbb{Q}}$, where Γ_p denotes the stabilizer in Γ of p . If σ_p and σ_q are scaling elements for Γ at p and q , respectively, then we have $\sigma_p^{-1} \Gamma_p \sigma_p = B(\mathbb{Z})$, and similarly with q in place of p , so the set of *translates*

$$\sigma_p^{-1} (\Gamma_p \backslash \Gamma / \Gamma_q) \sigma_q = \{\sigma_p^{-1} \Gamma_p \gamma \Gamma_q \sigma_q \mid \gamma \in \Gamma\} \quad (2.6.6)$$

is in fact a set of double cosets of $B(\mathbb{Z})$. In the notation of §2.4 we have

$$\sigma_p^{-1} (\Gamma_p \backslash \Gamma / \Gamma_q) \sigma_q = \llbracket \sigma_p^{-1} \Gamma \sigma_q \rrbracket = \{ \llbracket \sigma_p^{-1} \gamma \sigma_q \rrbracket \mid \gamma \in \Gamma \}. \quad (2.6.7)$$

Set $\mathfrak{p} = \Gamma \cdot p$ and $\mathfrak{q} = \Gamma \cdot q$, and set $\Sigma_\mathfrak{p} = \Gamma \sigma_p$ and $\Sigma_\mathfrak{q} = \Gamma \sigma_q$, so that $\Sigma_\mathfrak{p}$ and $\Sigma_\mathfrak{q}$ are scaling cosets for Γ at $\mathfrak{p}$ and $\mathfrak{q}$, respectively. Then we have $\sigma_p^{-1} \Gamma \sigma_q = \Sigma_\mathfrak{p}^{-1} \Sigma_\mathfrak{q}$, so that these translates depend only on cusps, and not on cusp representatives. We see from this discussion that for any

pair of cusps $\mathfrak{p}, \mathfrak{q} \in P_\Gamma$, with scaling cosets $\Sigma_{\mathfrak{p}}$ and $\Sigma_{\mathfrak{q}}$, respectively, the set $\Sigma_{\mathfrak{p}}^{-1}\Sigma_{\mathfrak{q}}$ is a union of double cosets of $B(\mathbb{Z})$. The assignment $(\mathfrak{p}, \mathfrak{q}) \mapsto \Sigma_{\mathfrak{p}}^{-1}\Sigma_{\mathfrak{q}}$ is sensitive to the choice of scaling cosets $\Sigma_{\mathfrak{p}}$ and $\Sigma_{\mathfrak{q}}$, to the extent that a different choice will replace $\Sigma_{\mathfrak{p}}^{-1}\Sigma_{\mathfrak{q}}$ with a set of the form $T^\alpha \Sigma_{\mathfrak{p}}^{-1}\Sigma_{\mathfrak{q}} T^\beta$ for some $\alpha, \beta \in \mathbb{Q}$.

In general there may be no canonical choice of scaling coset for a particular group at a particular cusp, but there are situations in which some choices might be preferred over others. For example, it is natural to take $\Sigma_{\Gamma \cdot \infty} = \Gamma$ in case Γ has width one at infinity, for we then have $\Sigma_{\mathfrak{p}}^{-1}\Sigma_{\mathfrak{q}} = \Sigma_{\mathfrak{p}}^{-1}$ when $\mathfrak{q} = \Gamma \cdot \infty$, and $\Sigma_{\mathfrak{p}}^{-1}\Sigma_{\mathfrak{q}} = \Sigma_{\mathfrak{q}}$ when $\mathfrak{p} = \Gamma \cdot \infty$, and $\Sigma_{\mathfrak{p}}^{-1}\Sigma_{\mathfrak{q}} = \Gamma$ when $\mathfrak{p} = \mathfrak{q} = \Gamma \cdot \infty$. We convene to always take the scaling coset $\Sigma_{\Gamma \cdot \infty}$, for Γ at the infinite cusp, to be of the form $\Gamma[\mu]$ for $\mu \in \mathbb{Q}^+$. This value μ is uniquely determined.

Observe that if $\Sigma_{\mathfrak{p}}$ and $\Sigma_{\mathfrak{q}}$ are scaling cosets for Γ at $\mathfrak{p}$ and $\mathfrak{q}$, respectively, then the intersection $\Sigma_{\mathfrak{p}}^{-1}\Sigma_{\mathfrak{q}} \cap B(\mathbb{Q}) = (\Sigma_{\mathfrak{p}}^{-1}\Sigma_{\mathfrak{q}})_\infty$ can be non-empty only in the case that $\mathfrak{p} = \mathfrak{q}$.

Given a system $\{\Sigma_{\mathfrak{p}} \mid \mathfrak{p} \in P_\Gamma\}$ of scaling cosets for Γ , we write $\Gamma^{\mathfrak{p}}$ as a shorthand for the group obtained as the conjugate of Γ by $\Sigma_{\mathfrak{p}}$.

$$\Gamma^{\mathfrak{p}} = \Sigma_{\mathfrak{p}}^{-1}\Sigma_{\mathfrak{p}} \quad (2.6.8)$$

The notation (2.6.8) suppresses the dependence on the choice of scaling coset: a different choice will replace $\Gamma^{\mathfrak{p}}$ with a group of the form $T^{-\alpha}\Gamma^{\mathfrak{p}}T^\alpha$ for some $\alpha \in \mathbb{Q}$.

2.7 Integrals

Write $\mathcal{O}(\mathbb{H})$ for the ring of holomorphic functions on the upper half plane $\mathbb{H}$.

For $\kappa \in \mathbb{Z}$, we define the *weight 2κ (right) action of $G(\mathbb{Q})$ on $\mathcal{O}(\mathbb{H})$* , to be denoted $(f, \chi) \mapsto f|^\kappa \chi$, by setting

$$(f|^\kappa \chi)(z) = f(\chi \cdot z) \text{jac}(\chi, z)^\kappa \quad (2.7.1)$$

for $f \in \mathcal{O}(\mathbb{H})$ and $\chi \in G(\mathbb{Q})$. For Γ a group commensurable with $G(\mathbb{Z})$ (and hence a subgroup of $G(\mathbb{Q})$), we call $f \in \mathcal{O}(\mathbb{H})$ an *unrestricted modular form of weight 2κ for Γ* in case it is a fixed point for the weight 2κ action of Γ . Suppose f is an unrestricted modular form of weight 2κ for Γ . Then for $X \in \Gamma \backslash G(\mathbb{Q})$ a right coset of Γ in $G(\mathbb{Q})$ we may define a function $(f|^\kappa_\Gamma X)(z)$ by setting

$$f|^\kappa_\Gamma X = f|^\kappa \chi \quad (2.7.2)$$

where χ is any representative for the coset $X \in \Gamma \backslash G(\mathbb{Q})$. Let $\mathfrak{p} \in P_\Gamma$ be a cusp of Γ and let $\Sigma_{\mathfrak{p}} \in \Gamma \backslash G(\mathbb{Q})$ a scaling coset for Γ at $\mathfrak{p}$. Then for f an unrestricted modular form of weight 2κ for Γ we define $f|_{\mathfrak{p}} \in \mathcal{O}(\mathbb{H})$ by setting $f|_{\mathfrak{p}} = f|^\kappa_\Gamma \Sigma_{\mathfrak{p}}$. Then $f|_{\mathfrak{p}}(z+1) = f|_{\mathfrak{p}}(z)$ for all $z \in \mathbb{H}$, by

the defining properties of $\Sigma_{\mathfrak{p}}$ (cf. §2.6), so we have

$$f|_{\mathfrak{p}}(z) = \sum_{n \in \mathbb{Z}} c|_{\mathfrak{p}}(n) e(nz) \quad (2.7.3)$$

for some $c|_{\mathfrak{p}}(n) \in \mathbb{Z}$. We call the right hand side of (2.7.3) the *Fourier expansion of f at $\mathfrak{p}$ with respect to $\Sigma_{\mathfrak{p}}$* , and we say that f is *meromorphic at $\mathfrak{p}$* if the right hand side of (2.7.3) is a meromorphic function of $q = e(z)$ in a neighborhood of $q = 0$. A different choice of scaling coset will replace $c|_{\mathfrak{p}}(n)$ with $e(n\alpha)c|_{\mathfrak{p}}(n)$ for some $\alpha \in \mathbb{Q}$, so the notion of being meromorphic at $\mathfrak{p}$ is independent of the choice of scaling coset at $\mathfrak{p}$.

We define $M_{\kappa}(\Gamma)$ to be the vector space consisting of unrestricted modular forms of weight 2κ for Γ that are meromorphic at the cusps of Γ . In a slight departure from standard convention we call $M_{\kappa}(\Gamma)$ the space of *modular forms of weight 2κ for Γ* . We define $S_{\kappa}(\Gamma)$ to be the vector space consisting of modular forms of weight 2κ for Γ that vanish at the cusps of Γ . We call $S_{\kappa}(\Gamma)$ the space of *cuspidal forms of weight 2κ for Γ* . The vector space $S_{\kappa}(\Gamma)$ is the zero vector space when $1 - \kappa \in \mathbb{Z}^+$.

Let $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$, and let $w \in \mathbb{H} \cup \hat{\mathbb{Q}}$. We associate to this data an operator on $\mathcal{O}(\mathbb{H})$, denoted $g \mapsto I_w^{\kappa}g$, by setting $I_{\infty}^{\kappa}g$ to be the zero function, and by setting

$$(I_w^{\kappa}g)(z) = 2\pi i \int_w^{\infty} g(\xi) (2\pi i(\xi - z))^{(-2\kappa)} d\xi \quad (2.7.4)$$

for $w \neq \infty$, where the integral is taken over the vertical line $\{w + it \mid t \in \mathbb{R}^+\}$. We define a closely related operator $g \mapsto J_w^{\kappa}g$ by setting

$$(J_w^{\kappa}g)(z) = \overline{(I_w^{\kappa}g)(\bar{z})}. \quad (2.7.5)$$

Observe that when $1 - \kappa \in \mathbb{Z}^+$, the functions $(I_w^{\kappa}g)(z)$ and $(J_w^{\kappa}g)(z)$ are polynomials of degree at most -2κ in z for any g . For $\chi \in G(\mathbb{Q})$ we have

$$(I_w^{\kappa}g)|^{\kappa}\chi = I_{w \cdot \chi}^{\kappa}(g|^{1-\kappa}\chi) - I_{\infty \cdot \chi}^{\kappa}(g|^{1-\kappa}\chi) \quad (2.7.6)$$

where $w \cdot \chi = \chi^{-1} \cdot w$ denotes the natural right action of $G(\mathbb{Q})$ on $\mathbb{H} \cup \hat{\mathbb{Q}}$. In particular, if $g \in M_{1-\kappa}(\Gamma)$ for some group Γ then we have $(I_w^{\kappa}g)|^{\kappa}\gamma = I_{w \cdot \gamma}^{\kappa}g - I_{\infty \cdot \gamma}^{\kappa}g$ for $\gamma \in \Gamma$.

Lemma 2.7.1. *Let Γ be a group commensurable with $G(\mathbb{Z})$, let $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$, and let $g \in S_{1-\kappa}(\Gamma)$. Then we have $I_p^{\kappa}g = 0$ for all $p \in \hat{\mathbb{Q}}$ if and only if $g = 0$.*

Proof. Certainly $I_p^{\kappa}g = 0$ for all $p \in \hat{\mathbb{Q}}$ if $g = 0$, so let $g \in S_{1-\kappa}(\Gamma)$ such that $I_p^{\kappa}g = 0$ for all

$p \in \hat{\mathbb{Q}}$. Following the proof of Lemma 3.2 in [Nie74] we define a function $f(z)$ by setting

$$f(z) = (I_z^\kappa g)(z) = 2\pi \mathbf{i} \int_z^\infty g(\xi) (2\pi \mathbf{i}(\xi - z))^{(-2\kappa)} d\xi. \quad (2.7.7)$$

The identity (2.7.6) implies $f|^\kappa \gamma = f - I_{\infty, \gamma}^\kappa g$ for $\gamma \in \Gamma$, so that f is an unrestricted modular form of weight 2κ for Γ . We should consider the behavior of f at the cusps of Γ . Let $\mathbf{p} \in \mathbf{P}_\Gamma$, and let $\Sigma_{\mathbf{p}}$ be a scaling coset for Γ at $\mathbf{p}$. Then we have

$$g|_{\mathbf{p}}(\xi) = \sum_{n \in \mathbb{Z}^+} b_{\mathbf{p}}(n) e(n\xi) \quad (2.7.8)$$

for some $b_{\mathbf{p}}(n) \in \mathbb{C}$, where $g|_{\mathbf{p}} = g|_{\Gamma}^{1-\kappa} \Sigma_{\mathbf{p}}$ (cf. (2.7.2)). In order to analyze $f|_{\mathbf{p}} = f|_{\Gamma}^\kappa \Sigma_{\mathbf{p}}$ we choose $\sigma \in \Sigma_{\mathbf{p}}$ and compute

$$f|^\kappa \sigma = I_z^\kappa (g|_{\mathbf{p}}) - I_{\infty, \sigma}^\kappa (g|_{\mathbf{p}}) \quad (2.7.9)$$

by (2.7.6). Now the term $I_{\infty, \sigma}^\kappa (g|_{\mathbf{p}})$ vanishes, since we have

$$\begin{aligned} (I_{\infty, \sigma}^\kappa (g|_{\mathbf{p}}))(z) &= 2\pi \mathbf{i}^{1-2\kappa} \int_{\sigma^{-1} \cdot \infty}^\infty g(\sigma \cdot \xi) \text{jac}(\sigma, \xi)^{1-\kappa} (\xi - z)^{(-2\kappa)} d\xi \\ &= 2\pi \mathbf{i}^{1-2\kappa} \int_{\sigma^{-1} \cdot \infty}^\infty g(\sigma \cdot \xi) (\sigma \cdot \xi - \sigma \cdot z)^{(-2\kappa)} \text{jac}(\sigma, z)^\kappa d(\sigma \cdot \xi) \\ &= -(I_{\sigma, \infty}^\kappa g) \text{jac}(\sigma, z)^\kappa, \end{aligned} \quad (2.7.10)$$

which vanishes by our hypothesis on g . So we have the Fourier expansion

$$f|_{\mathbf{p}}(z) = \sum_{n \in \mathbb{Z}^+} c_{\mathbf{p}}(n) e(nz) \quad (2.7.11)$$

for f at $\mathbf{p}$, where $c_{\mathbf{p}}(n)$ is determined by $b_{\mathbf{p}}(n)$ according to the formula

$$c_{\mathbf{p}}(n) = b_{\mathbf{p}}(n) 2\pi \mathbf{i} (I_0^\kappa g)(0) = -n^{2\kappa-1} b_{\mathbf{p}}(n). \quad (2.7.12)$$

We conclude that f is a cusp form of weight κ . Since $1 - \kappa \in \mathbb{Z}^+$ we have $S_\kappa(\Gamma) = 0$, and so $c_{\mathbf{p}}(n) = 0$ for all $n \in \mathbb{Z}$ and all $\mathbf{p} \in \mathbf{P}_\Gamma$, and so $b_{\mathbf{p}}(n) = 0$ for all $n \in \mathbb{Z}$ and all $\mathbf{p} \in \mathbf{P}_\Gamma$. That is to say, g vanishes identically, which is what we required to show. $\square$

Let $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$. Following Niebur (cf. [Nie74]) we say that a holomorphic function $f \in \mathcal{O}(\mathbb{H})$ is an *unrestricted automorphic integral of weight 2κ* for Γ in case there exists

a cusp form $g \in S_{1-\kappa}(\Gamma)$ such that

$$f|^\kappa \gamma = f - J_{\infty, \gamma}^\kappa g \quad (2.7.13)$$

for each $\gamma \in \Gamma$. According to Lemma 2.7.1 the cusp form $g \in S_{1-\kappa}(\Gamma)$ satisfying (2.7.13) is uniquely determined. We call it the *cusp form associated to the unrestricted automorphic integral* f . The operator $f \mapsto f|^\kappa_\Gamma X$, for X a right coset of Γ , may be extended to unrestricted automorphic integrals as follows. Suppose f is an unrestricted automorphic integral of weight 2κ for Γ with associated cusp form g . Then for $X \in \Gamma \backslash G(\mathbb{Q})$ we may define $f|^\kappa_\Gamma X \in \mathcal{O}(\mathbb{H})$ by setting

$$f|^\kappa_\Gamma X = (f - J_{\chi, \infty}^\kappa g)|^\kappa \chi \quad (2.7.14)$$

where χ is any representative for the coset $X \in \Gamma \backslash G(\mathbb{Q})$. The transformation properties (2.7.6) and (2.7.13) confirm that the function $f|^\kappa_\Gamma X$ is independent of the choice of coset representative χ . Let $\Sigma_{\mathfrak{p}}$ be a scaling coset for Γ at a cusp $\mathfrak{p} \in P_\Gamma$, and define $f|_{\mathfrak{p}}$ by setting

$$f|_{\mathfrak{p}} = f|^\kappa_\Gamma \Sigma_{\mathfrak{p}}. \quad (2.7.15)$$

By the defining properties of scaling cosets we find that $f|_{\mathfrak{p}}(z+1) = f|_{\mathfrak{p}}(z)$ for all $z \in \mathbb{H}$, and so we have

$$f|_{\mathfrak{p}}(z) = \sum_{n \in \mathbb{Z}} c_{|\mathfrak{p}}(n) e(nz) \quad (2.7.16)$$

for some $c_{|\mathfrak{p}}(n) \in \mathbb{C}$. We call (2.7.16) the *Fourier expansion of f at $\mathfrak{p}$ with respect to $\Sigma_{\mathfrak{p}}$* , and we say that f is *meromorphic at $\mathfrak{p}$* if the right hand side of (2.7.16) is a meromorphic function of $q = e(z)$ in a neighborhood of $q = 0$.

We define $I_\kappa(\Gamma)$ to be the vector space consisting of unrestricted automorphic integrals of weight 2κ for Γ that are meromorphic at every cusp of Γ . We call $I_\kappa(\Gamma)$ the space of *automorphic integrals of weight 2κ for Γ* .

The space $M_\kappa(\Gamma)$ of modular forms of weight 2κ for Γ is a subspace of $I_\kappa(\Gamma)$ by definition. Lemma 2.7.1 may be regarded as stating that $M_\kappa(\Gamma)$ is the kernel of the map $I_\kappa(\Gamma) \rightarrow S_{1-\kappa}(\Gamma)$ which sends an automorphic integral f to its associated cusp form g (cf. (2.7.13)). We will see in §4.5 that the map $I_\kappa(\Gamma) \rightarrow S_{1-\kappa}(\Gamma)$ is surjective, so that we have an exact sequence of vector spaces

$$0 \rightarrow M_\kappa(\Gamma) \rightarrow I_\kappa(\Gamma) \rightarrow S_{1-\kappa}(\Gamma) \rightarrow 0. \quad (2.7.17)$$

The sequence (2.7.17) is trivially exact in case $\kappa \in \mathbb{Z}^+$, for then $S_{1-\kappa}(\Gamma) = 0$, so that an automorphic integral of weight 2κ is automatically a modular form. Indeed, and more generally, if $\kappa \in \mathbb{Z}$ is chosen so that $S_{1-\kappa}(\Gamma) = 0$, then the spaces $M_\kappa(\Gamma)$ and $I_\kappa(\Gamma)$ coincide by the definition (cf. (2.7.13)) of an automorphic integral.

3 Rademacher sums

In this section we associate Rademacher sums, and conjugate Rademacher sums (cf. §1.3), of arbitrary even integer weight to triples $(\Gamma, \mathfrak{p}, \mathfrak{q})$ where Γ is a group commensurable with the modular group and $\mathfrak{p}$ and $\mathfrak{q}$ are cusps for Γ . Following the methods of Niebur we derive explicit expressions for the Fourier expansions of these functions, and study how they transform under the action of Γ .

3.1 Construction

Given $\kappa, m \in \mathbb{Z}$ and $[\chi] \in [G(\mathbb{Q})]$, define a function $z \mapsto P_{[\chi]}^{\kappa(m)}(z)$ on $\mathbb{H}$ by setting

$$P_{[\chi]}^{\kappa(m)}(z) = e(-m [\chi \cdot z]) \text{jac}([\chi], z)^\kappa. \quad (3.1.1)$$

Given $U \subset [G(\mathbb{Q})]$, define the *holomorphic Poincaré series of weight 2κ and order m associated to U* , to be denoted $P_U^{\kappa(m)}(z)$, by setting

$$P_U^{\kappa(m)}(z) = \lim_{K \rightarrow \infty} \sum_{[\chi] \in U_{\leq K}} P_{[\chi]}^{\kappa(m)}(z). \quad (3.1.2)$$

If $\kappa > 1$ and $1 - m \in \mathbb{Z}^+$ then for $U = [\Gamma]$ say, for Γ a group commensurable with $G(\mathbb{Z})$, the limiting sum in (3.1.2) converges absolutely and the symbols $\lim_{K \rightarrow \infty}$ and $\leq_K$ are unnecessary. If $1 - \kappa \in \mathbb{Z}^+$ then even with the limit the expression on the right hand side of (3.1.2) fails to converge, and the Poincaré series $P_U^{\kappa(m)}(z)$ is not defined.

As was originally demonstrated by Rademacher in [Rad39], the non-convergence of the Poincaré series $P_U^{\kappa(m)}(z)$ when $\kappa = 0$ can be circumvented, by replacing $P_{[\chi]}^{0(m)}(z)$ with $P_{[\chi]}^{0(m)}(z) - P_{0, [\chi]}^{0(m)}$ in the right-hand side of (3.1.2) where

$$P_{[\chi]}^{0(m)}(z) - P_{0, [\chi]}^{0(m)} = e(-m [\chi \cdot z]) - e(-m [\chi \cdot \infty]) \quad (3.1.3)$$

in case $[\chi] \in [G(\mathbb{Q})]^\times$, and $P_{0, [\chi]}^{0(m)} = 0$ otherwise. Rademacher showed for $m = 1$ (cf. loc. cit.) that the resulting expression is a conditionally convergent series which recovers a function invariant for the modular group $G(\mathbb{Z})$ in the case that $U = [G(\mathbb{Z})]$. Generalizations of Rademacher's construction were given by Knopp (cf. [Kno62a]) and Niebur (cf. [Nie74]) so

as to obtain conditionally convergent Poincaré series for arbitrary real non-positive weights, for various subgroups of $G(\mathbb{R}) = \mathrm{PSL}_2(\mathbb{R})$.

With the constructions of Rademacher, Knopp and Niebur in mind, we introduce the *Rademacher component function of weight 2κ and order m associated to $\llbracket \chi$* , denoted $z \mapsto R_{\llbracket \chi}^{\kappa(m)}(z)$ and defined on $\mathbb{H}$ for $\kappa, m \in \mathbb{Z}$ and $\llbracket \chi \in G(\mathbb{Q})$, by setting

$$R_{\llbracket \chi}^{\kappa(m)}(z) = e(-m \llbracket \chi \cdot z) r^\kappa(m, \llbracket \chi, z) \mathrm{jac}(\llbracket \chi, z)^\kappa, \quad (3.1.4)$$

where $r^\kappa(m, \llbracket \chi, z)$ is the *Rademacher regularization factor of weight 2κ* , given by

$$r^\kappa(m, \llbracket \chi, z) = \Phi(1 - 2\kappa, 2 - 2\kappa, m \llbracket \chi \cdot z - m \llbracket \chi \cdot \infty)(2\pi i(m \llbracket \chi \cdot \infty - m \llbracket \chi \cdot z))^{1-2\kappa} \quad (3.1.5)$$

in case $\chi \in G(\mathbb{Q})^\times$, and $r^\kappa(m, \llbracket \chi, z) = 1$ otherwise (cf. (2.1.3)). For $U \subset G(\mathbb{Q})$ we define the *classical Rademacher sum of weight 2κ and order m associated to U* , to be denoted $R_U^{\kappa(m)}(z)$, by setting

$$R_U^{\kappa(m)}(z) = \lim_{K \rightarrow \infty} \sum_{\llbracket \chi \in U \leq K} R_{\llbracket \chi}^{\kappa(m)}(z). \quad (3.1.6)$$

Employing the Kummer transformation (2.1.4) we may rewrite the Rademacher component function $R_{\llbracket \chi}^{\kappa(m)}(z)$ as

$$R_{\llbracket \chi}^{\kappa(m)}(z) = e(-m \llbracket \chi \cdot \infty) e(m \llbracket \chi \cdot \infty - m \llbracket \chi \cdot z, 1 - 2\kappa) \mathrm{jac}(\llbracket \chi, z)^\kappa \quad (3.1.7)$$

in case $\llbracket \chi \in G(\mathbb{Q})^\times$, where $e(z, s)$ is the generalized exponential function of (2.1.6). Since we restrict attention to the case that κ is an integer (cf. [Nie74] for non-integral weights), the generalized exponential in (3.1.7) may in turn be written in terms of the partial exponential function $e(z)_{<K}$ (cf. §2.1). We thus obtain the identity

$$e(-m \llbracket \chi \cdot z) r^\kappa(m, \llbracket \chi, z) = e(-m \llbracket \chi \cdot z) - e(-m \llbracket \chi \cdot \infty) e(m \llbracket \chi \cdot \infty - m \llbracket \chi \cdot z)_{<1-2\kappa} \quad (3.1.8)$$

when $\chi \in G(\mathbb{Q})^\times$. In particular, we have $r^\kappa(m, \llbracket \chi, z) = 1$ when κ is positive, so that the Rademacher sum $R_U^{\kappa(m)}(z)$ is exactly the Poincaré series $P_U^{\kappa(m)}(z)$ when $\kappa \in \mathbb{Z}^+$.

The case that $\kappa = 0$ is special, for the corresponding Poincaré series, if it were to converge absolutely, would define a Γ -invariant function on $\mathbb{H}$ for any group Γ satisfying $U\Gamma = U$. With $\kappa = 0$ we have $r^0(m, \llbracket \chi, z) = 1 - e(m \llbracket \chi \cdot z - m \llbracket \chi \cdot \infty)$ when $\chi \in G(\mathbb{Q})^\times$, and hence

$$R_{\llbracket \chi}^{0(m)}(z) = P_{\llbracket \chi}^{0(m)}(z) - P_{0, \llbracket \chi}^{0(m)} \quad (3.1.9)$$

for all $\llbracket \chi \in \llbracket G(\mathbb{Q})^\times$. In other words, the Rademacher sum $R_U^{0(m)}(z)$ is given by

$$\begin{aligned} R_U^{0(m)}(z) &= \lim_{K \rightarrow \infty} \sum_{\llbracket \chi \in U_{\leq K}^\times} P_{\llbracket \chi}^{0(m)}(z) - P_{0, \llbracket \chi}^{0(m)} \\ &= \sum_{\llbracket \chi \in U_\infty} e(-m \llbracket \chi \cdot z) + \lim_{K \rightarrow \infty} \sum_{\llbracket \chi \in U_{\leq K}^\times} e(-m \llbracket \chi \cdot z) - e(-m \llbracket \chi \cdot \infty), \end{aligned} \quad (3.1.10)$$

with the latter expression holding at least when there are only finitely many cosets of $B(\mathbb{Z})$ in U_∞ . This expression (3.1.10), with $m = 1$ and $U = \llbracket G(\mathbb{Z})$, is the series originally considered by Rademacher in [Rad39].

It is interesting to consider the function obtained by substituting $\bar{z}$ for z in the right hand side of (3.1.6). We define the *conjugate Rademacher sum of weight 2κ and order m associated to U* , to be denoted $\bar{R}_U^{\kappa(m)}(z)$, by setting

$$\bar{R}_U^{\kappa(m)}(z) = \lim_{K \rightarrow \infty} \sum_{\llbracket \chi \in U_{\leq K}} \bar{R}_{\llbracket \chi}^{\kappa(m)}(z), \quad (3.1.11)$$

where $\bar{R}_{\llbracket \chi}^{\kappa(m)}(z) = R_{\llbracket \chi}^{\kappa(m)}(\bar{z})$ by definition. Evidently, the conjugate Rademacher sum $\bar{R}_U^{\kappa(m)}(z)$ is an anti-holomorphic function on $\mathbb{H}$, whenever the limit defining it exists. At first glance it appears that we should recover the classical Rademacher sum $R_U^{\kappa(m)}(z)$ from the conjugate Rademacher sum according to the identity $R_U^{\kappa(m)}(z) = \bar{R}_U^{\kappa(m)}(\bar{z})$, but as we shall see in §3.4, the relationship between the holomorphic functions $R_U^{\kappa(m)}(z)$ and $\bar{R}_U^{\kappa(m)}(\bar{z})$ is generally more interesting than this.

We typically take U to be of the form $U = \llbracket \Sigma_{\mathbf{p}}^{-1} \Sigma_{\mathbf{q}}$ where $\{\Sigma_{\mathbf{p}} \mid \mathbf{p} \in \mathbf{P}_\Gamma\}$ is a system of scaling cosets (cf. §2.6) for some group Γ commensurable with the modular group, and $\mathbf{p}, \mathbf{q} \in \mathbf{P}_\Gamma$ are cusps of Γ . In this case we write $R_{\Gamma, \mathbf{p}|\mathbf{q}}^{\kappa(m)}(z)$ for $R_U^{\kappa(m)}(z)$, and similarly for the holomorphic Poincaré series, suppressing the choice of scaling cosets from notation. A change in the choice of $\Sigma_{\mathbf{p}}$ and $\Sigma_{\mathbf{q}}$ replaces $R_{\Gamma, \mathbf{p}|\mathbf{q}}^{\kappa(m)}(z)$ with a function of the form $e(\alpha) R_{\Gamma, \mathbf{p}|\mathbf{q}}^{\kappa(m)}(z + \beta)$ for some $\alpha, \beta \in \mathbb{Q}$, and similarly for the holomorphic Poincaré series $P_{\Gamma, \mathbf{p}|\mathbf{q}}^{\kappa(m)}(z)$.

In the case that $\mathbf{p}$ or $\mathbf{q}$ is the *infinite cusp* $\Gamma \cdot \infty$ we omit it from notation, writing $R_{\Gamma, \mathbf{p}}^{\kappa(m)}(z)$ for $R_{\Gamma, \mathbf{p}|\Gamma \cdot \infty}^{\kappa(m)}(z)$, and $R_{\Gamma, \mathbf{q}}^{\kappa(m)}(z)$ for $R_{\Gamma, \Gamma \cdot \infty|\mathbf{q}}^{\kappa(m)}(z)$, and similarly for the holomorphic Poincaré series. The functions $R_{\Gamma, \mathbf{p}}^{\kappa(m)}(z)$ are, in a sense, the most important, for we shall see in §3.4 that $R_{\Gamma, \mathbf{p}}^{\kappa(m)}(z)$ is, up to a certain constant function, an automorphic integral of weight 2κ for Γ with a single pole at the cusp $\mathbf{p}$ in case Γ has width one at infinity (cf. §2.6) and $1 - \kappa, m \in \mathbb{Z}^+$. We shall see also in §3.4 that the function $R_{\Gamma, \mathbf{p}|\mathbf{q}}^{\kappa(m)}(z)$, once corrected by a certain constant function, may be regarded as encoding the behavior of (the above mentioned correction by constant function of) $R_{\Gamma, \mathbf{p}}^{\kappa(m)}(z)$ at the cusp $\mathbf{q}$.

To further emphasize the importance of the functions $R_{\Gamma, \mathbf{p}}^{\kappa(m)}(z)$, we observe that every

Rademacher sum $R_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z)$ is of the form $R_{\Gamma', \mathfrak{p}'}^{\kappa(m)}(z)$ for some group Γ' with width one at infinity, and some cusp $\mathfrak{p}'$ of Γ' . For if $\Sigma_{\mathfrak{p}}$ and $\Sigma_{\mathfrak{q}}$ are the chosen scaling sets for Γ at $\mathfrak{p}$ and $\mathfrak{q}$, respectively, then we have $R_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z) = R_U^{\kappa(m)}(z)$ for $U = \llbracket \Sigma_{\mathfrak{p}}^{-1} \Sigma_{\mathfrak{q}} \rrbracket$. Recall (cf. (2.6.8)) that $\Gamma^{\mathfrak{q}}$ is a shorthand for the group $\Sigma_{\mathfrak{q}}^{-1} \Sigma_{\mathfrak{q}}$, and this group $\Gamma^{\mathfrak{q}}$ has width one at infinity by the defining properties of scaling cosets. If we define $\mathfrak{p}^{\mathfrak{q}} = \Sigma_{\mathfrak{q}}^{-1} \cdot \mathfrak{p} \in \hat{\mathbb{Q}}$ then $\mathfrak{p}^{\mathfrak{q}}$ is a cusp of $\Gamma^{\mathfrak{q}}$, and $\Sigma_{\mathfrak{p}^{\mathfrak{q}}} = \Sigma_{\mathfrak{q}}^{-1} \Sigma_{\mathfrak{p}}$ is a scaling coset for $\Gamma^{\mathfrak{q}}$ at $\mathfrak{p}^{\mathfrak{q}}$. We now have $\Sigma_{\mathfrak{p}}^{-1} \Sigma_{\mathfrak{q}} = \Sigma_{\mathfrak{p}^{\mathfrak{q}}}^{-1}$, so that $U = \llbracket \Sigma_{\mathfrak{p}^{\mathfrak{q}}}^{-1} \rrbracket$. Since $\Gamma^{\mathfrak{q}}$ has width one at infinity we may take it to be a scaling coset for itself at the infinite cusp, and we thus have

$$R_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z) = R_{\Gamma^{\mathfrak{q}}, \mathfrak{p}^{\mathfrak{q}}}^{\kappa(m)}(z) \quad (3.1.12)$$

subject to the understanding that the data defining the right hand side of (3.1.12) is related to the data defining the left hand side of (3.1.12) by $\Gamma^{\mathfrak{q}} = \Sigma_{\mathfrak{q}}^{-1} \Sigma_{\mathfrak{q}}$ and $\mathfrak{p}^{\mathfrak{q}} = \Sigma_{\mathfrak{q}}^{-1} \cdot \mathfrak{p}$ and $\Sigma_{\mathfrak{p}^{\mathfrak{q}}} = \Sigma_{\mathfrak{q}}^{-1} \Sigma_{\mathfrak{p}}$.

Our primary interest in this article is in the distinguished case that $\kappa = 0$. In order to simplify notation, and maintain consistency with the notation of §1, we write $R_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(m)}(z)$ as a shorthand for $R_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{0(m)}(z)$.

We conclude this section with a result which expresses the Rademacher component function $R_{\llbracket \chi \rrbracket}^{\kappa(m)}(z)$ in terms of the functions $P_{\llbracket \chi \rrbracket}^{\kappa(m)}(z)$ of (3.1.1) and the integral operator J_w^{κ} of §2.7.

Lemma 3.1.1. *Let $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$, let $m \in \mathbb{Z}^+$, and let $\llbracket \chi \rrbracket \in \llbracket G(\mathbb{Q}) \rrbracket$. Then we have*

$$R_{\llbracket \chi \rrbracket}^{\kappa(m)}(z) = P_{\llbracket \chi \rrbracket}^{\kappa(m)}(z) + m^{1-2\kappa} \left(J_{\infty, \llbracket \chi \rrbracket}^{\kappa} P_{\llbracket \chi \rrbracket}^{1-\kappa(-m)} \right) (z). \quad (3.1.13)$$

Proof. In case $\llbracket \chi \rrbracket \in \llbracket G(\mathbb{Q}) \rrbracket_{\infty}$ the second term in the right hand side of (3.1.13) vanishes (cf. (3.1.15)), and the identity (3.1.13) then holds by the definition of $R_{\llbracket \chi \rrbracket}^{\kappa(m)}(z)$, so we assume henceforth that $\llbracket \chi \rrbracket \in \llbracket G(\mathbb{Q}) \rrbracket^{\times}$. By (3.1.8) we may write the Rademacher component function $R_{\llbracket \chi \rrbracket}^{\kappa(m)}(z)$ as

$$e(-m \llbracket \chi \rrbracket \cdot z) \text{jac}(\llbracket \chi \rrbracket, z)^{\kappa} - e(-m \llbracket \chi \rrbracket \cdot \infty) e(m \llbracket \chi \rrbracket \cdot \infty - m \llbracket \chi \rrbracket \cdot z)_{<1-2\kappa} \text{jac}(\llbracket \chi \rrbracket, z)^{\kappa}. \quad (3.1.14)$$

The first term in (3.1.14) is just $P_{\llbracket \chi \rrbracket}^{\kappa(m)}(z)$, and a contour integral calculation confirms that the second term in (3.1.14) is $-m^{1-2\kappa}$ times the image of $P_{\llbracket \chi \rrbracket}^{1-\kappa(-m)}(z)$ under $J_{\infty, \llbracket \chi \rrbracket}^{\kappa}$. The details of this contour integral calculation are as follows. We have

$$\overline{\left(J_{\infty, \llbracket \chi \rrbracket}^{\kappa} P_{\llbracket \chi \rrbracket}^{1-\kappa(-m)} \right) (z)} = (2\pi i)^{1-2\kappa} \int_{\chi^{-1} \llbracket \cdot \rrbracket \cdot \infty}^{\infty} e(m \llbracket \chi \rrbracket \cdot \xi) \text{jac}(\llbracket \chi \rrbracket, \xi)^{1-\kappa} (\xi - \bar{z})^{(-2\kappa)} d\xi \quad (3.1.15)$$

by the definition of J_w^κ (cf. §2.7). Choosing a representative χ for the coset $\llbracket \chi \rrbracket$ we compute

$$\begin{aligned}
& \int_{\chi^{-1}\mathbb{I}, \infty}^{\infty} e(m \llbracket \chi \cdot \xi \rrbracket) \text{jac}(\llbracket \chi \rrbracket, \xi)^{1-\kappa} (\xi - \bar{z})^{(-2\kappa)} d\xi \\
&= e(m\chi \cdot \infty) \int_{\infty}^{\chi \cdot \infty} e(m\chi \cdot \xi - m\chi \cdot \infty) (\chi \cdot \xi - \chi \cdot \bar{z})^{(-2\kappa)} d(\chi \cdot \xi) \text{jac}(\chi, \bar{z})^\kappa \\
&= -e(m\chi \cdot \infty) \int_{\chi \cdot \infty}^{\infty} e(m\xi - m\chi \cdot \infty) (\xi - \chi \cdot \bar{z})^{(-2\kappa)} d\xi \text{jac}(\chi, \bar{z})^\kappa \\
&= -e(m\chi \cdot \infty) \int_0^{\infty} e(m\xi) (\xi + \chi \cdot \infty - \chi \cdot \bar{z})^{(-2\kappa)} d\xi \text{jac}(\chi, \bar{z})^\kappa
\end{aligned} \tag{3.1.16}$$

where the last integral in (3.1.16) is taken over the vertical line $\{\mathbf{i}t \mid t \in \mathbb{R}^+\}$. Thus, making the substitution $-2\pi \mathbf{i}m\xi = t$, we find that

$$\begin{aligned}
& \int_{\chi^{-1}\mathbb{I}, \infty}^{\infty} e(m \llbracket \chi \cdot \xi \rrbracket) \text{jac}(\llbracket \chi \rrbracket, \xi)^{1-\kappa} (\xi - \bar{z})^{(-2\kappa)} d\xi \\
&= -e(m\chi \cdot \infty) (-2\pi \mathbf{i}m)^{-1} \int_0^{\infty} e^{-t} ((-2\pi \mathbf{i}m)^{-1}t + \chi \cdot \infty - \chi \cdot \bar{z})^{(-2\kappa)} dt \text{jac}(\chi, \bar{z})^\kappa \\
&= -e(m\chi \cdot \infty) \sum_{k=0}^{-2\kappa} (-2\pi \mathbf{i}m)^{2\kappa-1} (-2\pi \mathbf{i}(m\chi \cdot \infty - m\chi \cdot \bar{z}))^{(k)} \text{jac}(\chi, \bar{z})^\kappa,
\end{aligned} \tag{3.1.17}$$

and from this we deduce that

$$\left(J_{\infty, \llbracket \chi \rrbracket}^\kappa P_{\llbracket \chi \rrbracket}^{1-\kappa(-m)} \right) (z) = -m^{2\kappa-1} e(-m\chi \cdot \infty) e(m\chi \cdot \infty - m\chi \cdot z)_{<1-2\kappa} \text{jac}(\chi, z)^\kappa. \tag{3.1.18}$$

Upon comparison with (3.1.14) we obtain the required identity (3.1.13). This completes the proof. $\square$

3.2 Coefficients

For $\llbracket \chi \rrbracket \in \llbracket G(\mathbb{Q}) \rrbracket^\times$ and $m, n \in \mathbb{Z}$, we define a holomorphic function $s \mapsto \text{Kl}_{\llbracket \chi \rrbracket}(m, n, s)$ on $\mathbb{C}$ by setting

$$\text{Kl}_{\llbracket \chi \rrbracket}(m, n, s) = e(-m \llbracket \chi \rrbracket \cdot \infty) e(-n \llbracket \chi^{-1} \rrbracket \cdot \infty) \varrho[\llbracket \chi \rrbracket]^s. \tag{3.2.1}$$

Here $\varrho[\llbracket \chi \rrbracket]$ denotes the scaling factor of (2.3.11) (cf. Lemma 2.4.1). If $\chi \in G(\mathbb{Q})^\times$ has a preferred representative $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ with $c > 0$, then we have

$$\text{Kl}_{\llbracket \chi \rrbracket}(m, n, s) = e\left(\frac{-ma + nd}{c}\right) \frac{(ad - bc)^s}{c^{2s}}, \tag{3.2.2}$$

so sums of the $\text{Kl}_{[\chi]}(m, n, s)$ over suitable subsets of $[G(\mathbb{Q})]^\times$ recover zeta functions of various kinds. For an example of this, define a function $s \mapsto \text{Kl}_S(m, n, s)$, for S a subset of $[G(\mathbb{Q})]^\times$, by setting

$$\text{Kl}_S(m, n, s) = \sum_{[\chi] \in S} \text{Kl}_{[\chi]}(m, n, s). \quad (3.2.3)$$

Then for $S = [\Gamma]^\times$ for some group $\Gamma < G(\mathbb{Q})$ with $\Gamma_\infty = B(\mathbb{Z})$, the function $\text{Kl}_S(m, n, s)$ is the *Selberg–Kloosterman zeta function associated to Γ* (cf. [GS83]). We write $\text{Kl}_{\Gamma, \mathfrak{p}|\mathfrak{q}}(m, n, s)$ for $\text{Kl}_S(m, n, s)$ in the case that $S = [\Sigma_{\mathfrak{p}}^{-1} \Sigma_{\mathfrak{q}}]^\times$ for Γ a subgroup of $G(\mathbb{Q})$ and $\{\Sigma_{\mathfrak{p}} \mid \mathfrak{p} \in \mathcal{P}_\Gamma\}$ a system of scaling cosets for Γ .

Consider the case that $S = [G(\mathbb{Z})]^\times$. Then the function $\text{Kl}_S(m, n, s)$ is closely related to the Riemann zeta function $\zeta(s)$, for we have

$$\text{Kl}_S(m, 0, s) = \sum_{c \in \mathbb{Z}^+} \sum_{a \in (\mathbb{Z}/c)^\times} \frac{e(-ma/c)}{c^{2s}} = \frac{\sigma(m, 1-2s)}{\zeta(2s)} \quad (3.2.4)$$

in the case that $m > 0$, where $\sigma(m, s)$ is the divisor function of (2.2.1). In particular, the function $s \mapsto \text{Kl}_S(1, 0, s/2)$, for $S = [G(\mathbb{Z})]^\times$, is just the reciprocal of the Riemann zeta function.

The following result is evident from the definition of $\text{Kl}_{[\chi]}(m, n, s)$, and the definition of the conjugation $\chi \mapsto \bar{\chi}$ (cf. §2.3).

Lemma 3.2.1. *For any $\chi \in G(\mathbb{Q})^\times$, $m, n \in \mathbb{Z}$, and $s \in \mathbb{C}$, we have*

$$\text{Kl}_{[\chi]}(m, n, s) = \text{Kl}_{[\chi^{-1}]}(n, m, s) = \text{Kl}_{[\bar{\chi}]}(-m, -n, s), \quad (3.2.5)$$

$$\text{Kl}_{[\chi]}(m, n, s) \varrho[\chi]^{1-s} = \text{Kl}_{[\chi]}(m, n, 1-s) \varrho[\chi]^s. \quad (3.2.6)$$

For a continuous analogue of the function $\text{Kl}_{[\chi]}(m, n, s)$ we define $\text{Bl}_{[\chi]}^\kappa(w, z)$, for $\chi \in G(\mathbb{Q})^\times$, $w, z \in \mathbb{C}$ and $\kappa \in \mathbb{Z}$, by setting

$$\text{Bl}_{[\chi]}^\kappa(w, z) = 2\pi i \text{Res}_{\xi=0} e\left(\frac{-w \varrho[\chi]}{\xi}\right) e(z\xi) \frac{1}{\xi^{2\kappa}} d\xi. \quad (3.2.7)$$

Lemma 3.2.2. *The function $\text{Bl}_{[\chi]}^\kappa(w, z)$ admits the following series representations.*

$$\text{Bl}_{[\chi]}^\kappa(w, z) = (-1)^\kappa \sum_{k \in \mathbb{N}} (4\pi^2)^{k+\kappa} \varrho[\chi]^k w^{(k)} z^{(k+2\kappa-1)} \quad (3.2.8)$$

$$\text{Bl}_{[\chi]}^\kappa(w, z) = (-1)^\kappa \sum_{k \in \mathbb{N}} (4\pi^2)^{k+1-\kappa} \varrho[\chi]^{k+1-2\kappa} w^{(k+1-2\kappa)} z^{(k)} \quad (3.2.9)$$

The first expression (3.2.8) holds for $\kappa \in \mathbb{Z}^+$. The second expression (3.2.9) holds for $1-\kappa \in \mathbb{Z}^+$.

Lemma 3.2.3. *For any $\chi \in G(\mathbb{Q})^\times$ and $w, z \in \mathbb{C}$ and $\kappa \in \mathbb{Z}$ we have*

$$\mathrm{Bl}_{[\chi]}^\kappa(w, z) \varrho[\chi]^\kappa = -\mathrm{Bl}_{[\chi]}^{1-\kappa}(z, w) \varrho[\chi]^{1-\kappa} = -\mathrm{Bl}_{[\chi]}^\kappa(-w, -z) \varrho[\chi]^\kappa. \quad (3.2.10)$$

The main construction of this section is the following. Given a subset $S \subset [G(\mathbb{Q})]^\times$ and $\kappa \in \mathbb{Z}$, we define the *coefficient function*, denoted $(m, n) \mapsto c_S^\kappa(m, n)$, by setting

$$c_S^\kappa(m, n) = \sum_{[\chi] \in S} \mathrm{Kl}_{[\chi]}(m, n, \kappa) \mathrm{Bl}_{[\chi]}^\kappa(m, n). \quad (3.2.11)$$

Lemma 3.2.4. *The coefficient functions $c_S^\kappa(m, n)$ admit the following series representations.*

$$c_S^\kappa(m, n) = (-1)^\kappa \sum_{[\chi] \in S} \sum_{k \in \mathbb{N}} e(-m[\chi] \cdot \infty) e(-n[\chi^{-1}] \cdot \infty) (4\pi^2 \varrho[\chi])^{k+\kappa} m^{(k)} n^{(k+2\kappa-1)} \quad (3.2.12)$$

$$c_S^\kappa(m, n) = (-1)^\kappa \sum_{[\chi] \in S} \sum_{k \in \mathbb{N}} e(-m[\chi] \cdot \infty) e(-n[\chi^{-1}] \cdot \infty) (4\pi^2 \varrho[\chi])^{k+1-\kappa} m^{(k+1-2\kappa)} n^{(k)} \quad (3.2.13)$$

The expression (3.2.12) holds for $\kappa \in \mathbb{Z}^+$. The expression (3.2.13) holds for $1 - \kappa \in \mathbb{Z}^+$.

Proposition 3.2.5. *For $S \subset [G(\mathbb{Q})]^\times$ and $m, n, \kappa \in \mathbb{Z}$ we have*

$$c_S^\kappa(m, n) = -c_{S^{-1}}^{1-\kappa}(n, m) = -c_S^\kappa(-m, -n). \quad (3.2.14)$$

Proof. These identities follow directly from Lemma 3.2.4, and the observation that the values $\varrho[\chi]$, $\varrho[\chi^{-1}]$ and $\varrho[\bar{\chi}]$ all coincide for any $\chi \in G(\mathbb{Q})^\times$. $\square$

In applications, we often take S to be of the form $S = [\Sigma_p^{-1} \Sigma_q]^\times$ where Σ_p and Σ_q are scaling cosets (cf. §2.6) for a group Γ at cusps $p, q \in P_\Gamma$. We write $c_{\Gamma, p|q}^\kappa(m, n)$ for $c_S^\kappa(m, n)$ in the case that $S = [\Sigma_p^{-1} \Sigma_q]^\times$. This notation suppresses the dependence on the choice of scaling cosets; the following lemma describes this dependence explicitly.

Lemma 3.2.6. *Let $p, q \in P_\Gamma$, let Σ_p and Σ'_p be scaling cosets for Γ at p and let Σ_q and Σ'_q be scaling cosets for Γ at q . Set $S = [\Sigma_p^{-1} \Sigma_q]^\times$ and $S' = [(\Sigma'_p)^{-1} \Sigma'_q]^\times$. Then we have $\Sigma'_p = \Sigma_p T^\alpha$ and $\Sigma'_q = \Sigma_q T^\beta$ and $c_{S'}^\kappa(m, n) = e(m\alpha + n\beta) c_S^\kappa(m, n)$ for some $\alpha, \beta \in \mathbb{Q}$.*

With a fixed choice of subset $S \subset [G(\mathbb{Q})]^\times$ and integer $\kappa \in \mathbb{Z}$, we assemble the coefficient functions $c_S^\kappa(m, n)$ into a formal series $\tilde{F}_S^\kappa(p, q)$ by setting

$$\tilde{F}_S^\kappa(p, q) = \sum_{m, n \in \mathbb{Z}} c_S^\kappa(m, n) p^m q^n. \quad (3.2.15)$$

The formal Fourier coefficients of the formal series $\tilde{F}_S^\kappa(p, q)$ with respect to the variable p are of particular importance. For fixed $S \subset \llbracket G(\mathbb{Q}) \rrbracket^\times$ and $\kappa \in \mathbb{Z}$ we define formal series $\tilde{F}_S^{\kappa(m)}(q)$ by requiring that

$$\tilde{F}_S^\kappa(p, q) = \sum_{m \in \mathbb{Z}} \tilde{F}_S^{\kappa(m)}(q) p^m. \quad (3.2.16)$$

We define the *singular* and *regular part* of $\tilde{F}_S^{\kappa(m)}(q)$, to be denoted $\tilde{F}_S^{\kappa(m)}(q)_{\text{sing}}$ and $\tilde{F}_S^{\kappa(m)}(q)_{\text{reg}}$, respectively, by setting

$$\tilde{F}_S^{\kappa(m)}(q)_{\text{sing}} = \sum_{n \in \mathbb{N}} c_S^\kappa(m, -n-1) q^{-n-1}, \quad \tilde{F}_S^{\kappa(m)}(q)_{\text{reg}} = \sum_{n \in \mathbb{N}} c_S^\kappa(m, n) q^n. \quad (3.2.17)$$

We define $\tilde{F}_S^{\kappa(m)}(q)_{\text{van}}$ to be the series obtained by removing the constant term from $\tilde{F}_S^{\kappa(m)}(q)_{\text{reg}}$.

$$\tilde{F}_S^{\kappa(m)}(q)_{\text{van}} = \sum_{n \in \mathbb{Z}^+} c_S^\kappa(m, n) q^n \quad (3.2.18)$$

Given the formal series $\tilde{F}_S^\kappa(p, q)$ for some S and κ , one may ask if the coefficient series $\tilde{F}_S^{\kappa(m)}(q)$ define holomorphic functions on $\mathbb{H}$ upon the substitution of $e(z)$ for q . Supposing this to be the case, we define $F_S^{\kappa(m)}(z)$ by setting

$$F_S^{\kappa(m)}(z) = \tilde{F}_S^{\kappa(m)}(e(z)) = \sum_{n \in \mathbb{Z}} c_S^\kappa(m, n) e(nz), \quad (3.2.19)$$

and we define $F_S^{\kappa(m)}(z)_{\text{sing}}$, $F_S^{\kappa(m)}(z)_{\text{reg}}$ and $F_S^{\kappa(m)}(z)_{\text{van}}$ in the analogous way.

We write $F_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z)$ for $F_S^{\kappa(m)}(z)$ in the case that $S = \llbracket \Sigma_{\mathfrak{p}}^{-1} \Sigma_{\mathfrak{q}} \rrbracket^\times$ for Γ a group commensurable with $G(\mathbb{Z})$ and for $\Sigma_{\mathfrak{p}}, \Sigma_{\mathfrak{q}} \in \Gamma \backslash G(\mathbb{Q})$ scaling cosets for Γ at cusps $\mathfrak{p}, \mathfrak{q} \in \mathcal{P}_\Gamma$. We write as a $F_S^{(m)}(z)$ as a shorthand for $F_S^{0(m)}(z)$, and interpret the notation $F_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(m)}(z)$, &c., similarly.

The regular part of the formal series $\tilde{F}_S^{\kappa(m)}(q)$ indeed defines a holomorphic function on $\mathbb{H}$, for suitable $\kappa, m \in \mathbb{Z}$, in the case that S is of the form $\llbracket \Sigma_{\mathfrak{p}}^{-1} \Sigma_{\mathfrak{q}} \rrbracket^\times$.

Proposition 3.2.7. *Let Γ be a group commensurable with $G(\mathbb{Z})$, and let $\mathfrak{p}, \mathfrak{q} \in \mathcal{P}_\Gamma$ be cusps of Γ . Let $\kappa, m \in \mathbb{Z}$ such that either $\kappa, 1-m \in \mathbb{Z}^+$ or $1-\kappa, m \in \mathbb{Z}^+$. Then the series*

$$F_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z)_{\text{van}} = \sum_{n \in \mathbb{Z}^+} c_{\Gamma, \mathfrak{p}|\mathfrak{q}}^\kappa(m, n) e(nz) \quad (3.2.20)$$

converges absolutely for $z \in \mathbb{H}$. In particular, $F_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z)_{\text{van}}$ is a holomorphic function on $\mathbb{H}$ whenever $\kappa, 1-m \in \mathbb{Z}^+$ or $1-\kappa, m \in \mathbb{Z}^+$.

The proof of Proposition 3.2.7 uses simple estimates, such as those appearing in Lemmas 2.5

and 2.6 of [Kno62a]. By directly similar methods we see that the singular part of the formal series $\tilde{F}_S^{\kappa(m)}(q)$ defines an anti-holomorphic function on $\mathbb{H}$ when S is of the form $[\Sigma_p^{-1}\Sigma_q]^\times$.

Proposition 3.2.8. *Let Γ be a group commensurable with $G(\mathbb{Z})$, and let $\mathfrak{p}, \mathfrak{q} \in P_\Gamma$ be cusps of Γ . Let $\kappa, m \in \mathbb{Z}$ such that either $\kappa, 1 - m \in \mathbb{Z}^+$ or $1 - \kappa, m \in \mathbb{Z}^+$. Then the series*

$$F_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(\bar{z})_{\text{sing}} = \sum_{n \in \mathbb{Z}^+} c_{\Gamma, \mathfrak{p}|\mathfrak{q}}^\kappa(m, -n) e(-n\bar{z}) \quad (3.2.21)$$

converges absolutely for $z \in \mathbb{H}$. In particular, $F_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(\bar{z})_{\text{sing}}$ is an anti-holomorphic function on $\mathbb{H}$ whenever $\kappa, 1 - m \in \mathbb{Z}^+$ or $1 - \kappa, m \in \mathbb{Z}^+$.

The notation $F_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}$ suppresses the dependence upon the choice of scaling cosets Σ_p and Σ_q . The next result encodes this dependence precisely.

Proposition 3.2.9. *Let Γ be a group commensurable with $G(\mathbb{Z})$, and let $\mathfrak{p}, \mathfrak{q} \in P_\Gamma$. Let Σ_p and Σ'_p be scaling cosets for Γ at $\mathfrak{p}$ and let Σ_q and Σ'_q be scaling cosets for Γ at $\mathfrak{q}$. Set $S = [\Sigma_p^{-1}\Sigma_q]^\times$ and $S' = [(\Sigma'_p)^{-1}\Sigma'_q]^\times$. Then we have $\Sigma'_p = \Sigma_p T^\alpha$ and $\Sigma'_q = \Sigma_q T^\beta$ and $F_{S'}^{\kappa(m)}(z) = e(m\alpha)F_S^{\kappa(m)}(z + \beta)$ for some $\alpha, \beta \in \mathbb{Q}$.*

3.3 Convergence

Our main goal in this section is to establish the convergence, and Fourier series expansion, of the Rademacher sum $R_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z)$ (cf. §3.1), attached to a group Γ commensurable with $G(\mathbb{Z})$, and cusps $\mathfrak{p}$ and $\mathfrak{q}$ for Γ . We will require only slight modifications of the arguments furnished by Niebur in [Nie74]. An important technical tool in these arguments is the following result.

Lemma 3.3.1. *Let $K \in \mathbb{R}^+ - \mathbb{Z}^+$ and let $z \in \mathbb{H}$. Then for $\Re(s) \geq 1$ we have*

$$\sum_{n \in \mathbb{Z}^+} n^{(s-1)} e(nz) = \sum_{\substack{n \in \mathbb{Z} \\ -K < n < K}} (-2\pi i)^{-s} (z + n)^{-s} + \lambda_K(z, s) \quad (3.3.1)$$

where the error term $\lambda_K(z, s)$ is given by

$$\lambda_K(z, s) = i \int_{-\infty}^{\infty} \left(\frac{(2\pi t + 2\pi i K)^{-s}}{1 - e(-K + it - z)} - \frac{(2\pi t - 2\pi i K)^{-s}}{1 - e(K + it - z)} \right) dt. \quad (3.3.2)$$

Proof. The proof follows that of Lemma 4.1 in [Nie74]. Define a function $f(\xi)$ by setting $f(\xi) = (-2\pi i \xi)^{-s} (1 - e(\xi - z))^{-1}$. Then $f(\xi)$ has a pole at $z + n$ for each $n \in \mathbb{Z}$, and no other poles, and the residue at $\xi = z + n$ is $(-2\pi i)^{1-s} (z + n)^{-s}$. We obtain -1 times the right hand side of (3.3.1) by integrating $f(\xi)$ over the rectangular contour integral with corners $\pm K \pm iL$ and letting $L \rightarrow \infty$. The integrals over the vertical sides of this contour contribute the term $\lambda_K(z, s)$, and the contribution of the horizontal sides vanishes as L tends to infinity. To

obtain (-1) times) the left hand side of (3.3.1) we evaluate this integral another way, replacing $(1 - e(\xi - z))^{-1}$ with the series $\sum_{n \in \mathbb{Z}^+} (-1) e(n(z - \xi))$ and using Hankel's integral expression for the reciprocal of the Gamma function to recognize the integral of $(-2\pi i \xi)^{-s} e(-n\xi)$ over our infinite rectangle as $n^{(s-1)}$. $\square$

Taking the limit as K tends to infinity in (3.3.1) for $\Re(s) > 1$ we obtain the identity

$$\sum_{n \in \mathbb{Z}^+} n^{(s-1)} e(nz) = \sum_{n \in \mathbb{Z}} (-2\pi i)^{-s} (z + n)^{-s} \quad (3.3.3)$$

in which both sums are absolutely convergent. In the case that $s = 1$ the error term $\lambda_K(z, 1)$ tends to $-1/2$ as $K \rightarrow \infty$, and we thus obtain

$$\sum_{n \in \mathbb{Z}^+} e(nz) = -\frac{1}{2} + \lim_{K \rightarrow \infty} \sum_{\substack{n \in \mathbb{Z} \\ -K < n < K}} (-2\pi i)^{-1} (z + n)^{-1}. \quad (3.3.4)$$

The formulas (3.3.3) and (3.3.4) are collectively known as the *Lipschitz summation formula*. Their derivation as a consequence of the Poisson summation formula is explained nicely in [KR01] (although the approach there does not supply the error term $\lambda_K(z, s)$).

Theorem 3.3.2. *Let Γ be a group commensurable with $G(\mathbb{Z})$, and let $\mathfrak{p}, \mathfrak{q} \in \mathcal{P}_\Gamma$ be cusps of Γ . Let $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$, and let $m \in \mathbb{Z}^+$. Then the limit defining the classical Rademacher sum $R_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z)$ converges to a holomorphic function on $\mathbb{H}$, and we have*

$$R_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z) = \delta_{\Gamma, \mathfrak{p}|\mathfrak{q}} e(-mz) + \frac{1}{2} c_{\Gamma, \mathfrak{p}|\mathfrak{q}}^\kappa(m, 0) + F_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z)_{\text{van}}. \quad (3.3.5)$$

Proof. Set $U = \llbracket \Sigma_{\mathfrak{p}}^{-1} \Sigma_{\mathfrak{q}} \rrbracket^\times$ and $S = \llbracket \Sigma_{\mathfrak{p}}^{-1} \Sigma_{\mathfrak{q}} \rrbracket^\times$. The argument of Lemma 4.3 in [Nie74] demonstrates that the limit $\lim_{K \rightarrow \infty} \sum_{\llbracket \chi \in U \leq K} R_{\llbracket \chi}^{\kappa(m)}(z)$ defining $R_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z)$ converges to a holomorphic function on $\mathbb{H}$, and may be rewritten in the following way (as a sum of limits of sums, rather than a limit of double sums).

$$R_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z) = R_U^{\kappa(m)}(z) = \sum_{c \in \mathbb{Z}^+} \lim_{K \rightarrow \infty} \sum_{\substack{\llbracket \chi \in U \leq K \\ c(\chi) = c}} R_{\llbracket \chi}^{\kappa(m)}(z) \quad (3.3.6)$$

Indeed, using Lemma 3.1.1 we may identify the Rademacher component function $R_{\llbracket \chi}^{\kappa(m)}(z)$ with the function $s_{\llbracket \chi}(z) + t_{\llbracket \chi}(z)$ defined on page 376 of [Nie74]. Then the right hand side of (3.3.6) is, up to a certain constant function, the definition of Rademacher sum used by Niebur in [Nie74]. The argument of Lemma 4.2 of [Nie74] uses the Lipschitz summation formula(s) (3.3.3) and (3.3.4) to equate the right hand side of (3.3.6) with the right hand side of (3.3.5). $\square$

The next result identifies the Fourier series expansion of the conjugate Rademacher sum $\bar{R}_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z)$. As such it is a natural counterpart to Theorem 3.3.2. The method of proof is identical to that of Theorem 3.3.2.

Theorem 3.3.3. *Let Γ be a group commensurable with $G(\mathbb{Z})$, and let $\mathfrak{p}, \mathfrak{q} \in \mathcal{P}_\Gamma$ be cusps of Γ . Let $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$, and let $m \in \mathbb{Z}^+$. Then the limit defining the conjugate Rademacher sum $\bar{R}_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z)$ converges to an anti-holomorphic function on $\mathbb{H}$, and we have*

$$\bar{R}_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z) = \delta_{\Gamma, \mathfrak{p}|\mathfrak{q}} e(-m\bar{z}) - \frac{1}{2} c_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa}(m, 0) - F_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(\bar{z})_{\text{sing}}. \quad (3.3.7)$$

Theorems 3.3.2 and 3.3.3 furnish the Fourier expansions of the classical and conjugate Rademacher sums of non-positive weight. Directly analogous methods can be used to determine expressions for the Fourier coefficients of the Rademacher sums of positive weight, which are, after all, just the holomorphic Poincaré series (cf. §3.1). Indeed, there are fewer technical difficulties in the case of positive weight, since the sum appearing in the definition (3.1.2) of the Poincaré series $P_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}$ is absolutely convergent (at least when $\kappa > 1$). For completeness we state an analogue of Theorem 3.3.2 for $\kappa \in \mathbb{Z}^+$. The result is quite standard (cf. [Iwa02]).

Theorem 3.3.4. *Let Γ be a group commensurable with $G(\mathbb{Z})$, and let $\mathfrak{p}, \mathfrak{q} \in \mathcal{P}_\Gamma$ be cusps of Γ . Let $\kappa \in \mathbb{Z}^+$, and let $m \in \mathbb{Z}$ such that $1 - m \in \mathbb{Z}^+$. Then for $z \in \mathbb{H}$ we have*

$$R_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z) = \delta_{\Gamma, \mathfrak{p}|\mathfrak{q}} e(-mz) + F_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z)_{\text{van}}. \quad (3.3.8)$$

Recall from §3.1 that we write $R_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(m)}(z)$ and for $R_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{0(m)}(z)$ and similarly for $\bar{R}_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(m)}(z)$. To conclude this section we consider the difference $R_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(m)}(z) - \bar{R}_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(m)}(z)$. Combining Theorems 3.3.2 and 3.3.3 we obtain the following identification of the Fourier expansion of this function in terms of the exponential function $e(z)$ and its conjugate $e(-\bar{z})$.

Theorem 3.3.5. *Let Γ be a group commensurable with $G(\mathbb{Z})$, let $\mathfrak{p}, \mathfrak{q} \in \mathcal{P}_\Gamma$ be cusps of Γ , and let $m \in \mathbb{Z}^+$. Then we have*

$$\begin{aligned} R_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(m)}(z) - \bar{R}_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(m)}(z) &= \delta_{\Gamma, \mathfrak{p}|\mathfrak{q}} (e(-mz) - e(-m\bar{z})) \\ &\quad + F_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(m)}(\bar{z})_{\text{sing}} + c_{\Gamma, \mathfrak{p}|\mathfrak{q}}(m, 0) + F_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(m)}(z)_{\text{van}} \end{aligned} \quad (3.3.9)$$

for $z \in \mathbb{H}$.

3.4 Variance

Recall from §3.1 that we write $R_{\Gamma, \mathfrak{p}}^{\kappa(m)}(z)$ for $R_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z)$ in case $\mathfrak{q}$ is the infinite cusp $\Gamma \cdot \infty$. Suppose that Γ is commensurable with the modular group and has width one at infinity (cf. §2.6). We will establish in this section that the Rademacher sum $R_{\Gamma, \mathfrak{p}}^{\kappa(m)}(z)$, once corrected by a certain

constant function, is an automorphic integral of weight 2κ for Γ whenever $\kappa, 1 - m \in \mathbb{Z}^+$ or $1 - \kappa, m \in \mathbb{Z}^+$. This is quite general since, according to the discussion of §3.1, any Rademacher sum $R_{\Gamma, \mathfrak{p}|q}^{\kappa(m)}(z)$ can be expressed in the form $R_{\Gamma', \mathfrak{p}'}^{\kappa(m)}(z)$ for some group Γ' having width one at infinity (cf. (3.1.12)).

Let us begin by considering the weight 2κ action of $G(\mathbb{Q})$ on the Rademacher component function $R_{\mathbb{L}\chi}^{\kappa(m)}(z)$ of (3.1.4).

Lemma 3.4.1. *Let $\mathbb{L}\chi \in \mathbb{L}(G(\mathbb{Q}))^\times$ and $\sigma \in G(\mathbb{Q})$, and let $\kappa, m \in \mathbb{Z}$. In case $\kappa, 1 - m \in \mathbb{Z}^+$ we have*

$$\left(R_{\mathbb{L}\chi}^{\kappa(m)} \Big|^\kappa \sigma \right) (z) = R_{\mathbb{L}\chi\sigma}^{\kappa(m)}(z), \quad (3.4.1)$$

and in case $1 - \kappa, m \in \mathbb{Z}^+$ we have

$$\left(R_{\mathbb{L}\chi}^{\kappa(m)} \Big|^\kappa \sigma \right) (z) = R_{\mathbb{L}\chi\sigma}^{\kappa(m)}(z) - m^{1-2\kappa} \left(J_{\infty \cdot \mathbb{L}\sigma}^\kappa P_{\mathbb{L}\chi\sigma}^{1-\kappa(-m)} \right) (z). \quad (3.4.2)$$

Proof. Note that we have

$$\left(P_{\mathbb{L}\chi}^{\kappa(m)} \Big|^\kappa \sigma \right) (z) = P_{\mathbb{L}\chi\sigma}^{\kappa(m)}(z) \quad (3.4.3)$$

for all $\kappa, m \in \mathbb{Z}$, where $P_{\mathbb{L}\chi}^{\kappa(m)}(z) = e(-m \mathbb{L}\chi \cdot z) \text{jac}(\mathbb{L}\chi, z)^\kappa$ (cf. (3.1.1)), so the identity (3.4.1) follows from (3.4.3) and the fact that $R_{\mathbb{L}\chi}^{\kappa(m)}(z) = P_{\mathbb{L}\chi}^{\kappa(m)}(z)$ in case $\kappa \in \mathbb{Z}^+$, by the definition of $R_{\mathbb{L}\chi}^{\kappa(m)}(z)$ (cf. §3.1). For (3.4.2) we suppose $1 - \kappa, m \in \mathbb{Z}^+$ and use Lemma 3.1.1 to write the Rademacher component function $R_{\mathbb{L}\chi}^{\kappa(m)}(z)$ in terms of the functions $P_{\mathbb{L}\chi}^{\kappa(m)}(z)$ and the integral operator J_w^κ . Then, employing (3.4.3) and the identity (2.7.6), we compute

$$\begin{aligned} \left(R_{\mathbb{L}\chi}^{\kappa(m)} \Big|^\kappa \sigma \right) (z) &= \left(P_{\mathbb{L}\chi}^{\kappa(m)} \Big|^\kappa \sigma \right) (z) + m^{1-2\kappa} \left(\left(J_{\infty \cdot \mathbb{L}\chi}^\kappa P_{\mathbb{L}\chi}^{1-\kappa(-m)} \right) \Big|^\kappa \sigma \right) (z) \\ &= P_{\mathbb{L}\chi\sigma}^{\kappa(m)}(z) + m^{1-2\kappa} \left(J_{\infty \cdot \mathbb{L}\chi\sigma}^\kappa P_{\mathbb{L}\chi\sigma}^{1-\kappa(-m)} \right) (z) - m^{1-2\kappa} \left(J_{\infty \cdot \mathbb{L}\sigma}^\kappa P_{\mathbb{L}\chi\sigma}^{1-\kappa(-m)} \right) (z). \end{aligned} \quad (3.4.4)$$

Employing Lemma 3.1.1 once again we recognize the first two terms on the right hand side (of the second line) of (3.4.4) to be $R_{\mathbb{L}\chi\sigma}^{\kappa(m)}(z)$. We thus obtain the required identity (3.4.2). $\square$

As for technical tools, in addition to Lemma 3.3.1, we also use the following result which shows that there is some flexibility in the convergence of the limit (3.1.6) defining the classical Rademacher sum.

Lemma 3.4.2. *Let Γ be a group commensurable with $G(\mathbb{Z})$, let $\mathfrak{p} \in \mathcal{P}_\Gamma$ be a cusp of Γ and let*

$\sigma \in G(\mathbb{Q})$. Let $\kappa, m \in \mathbb{Z}$ such that either $\kappa, 1 - m \in \mathbb{Z}^+$ or $1 - \kappa, m \in \mathbb{Z}^+$. Then we have

$$\lim_{K \rightarrow \infty} \sum_{[\chi] \in (U_{\leq K})^\sigma} R_{[\chi]}^{\kappa(m)}(z) = \lim_{K \rightarrow \infty} \sum_{[\chi] \in (U^\sigma)_{\leq K}} R_{[\chi]}^{\kappa(m)}(z) \quad (3.4.5)$$

for $U = [\Sigma_{\mathfrak{p}}^{-1}]$ and $z \in \mathbb{H}$.

Proof. For $\kappa > 1$ and $1 - m \in \mathbb{Z}^+$, both sums in (3.4.5) are absolutely convergent, and the identity certainly holds. In case $\kappa = 1$ and $1 - m \in \mathbb{Z}^+$, or $1 - \kappa \in \mathbb{Z}^+$ and $m \in \mathbb{Z}^+$, the identity (3.4.5) is established via the argument of Lemma 4.3 in [Nie74]. $\square$

Theorem 3.4.3. *Let Γ be a group commensurable with $G(\mathbb{Z})$ that has width one at infinity and let $\mathfrak{p} \in P_\Gamma$ be a cusp of Γ . Let $\kappa \in \mathbb{Z}^+$, and let $m \in \mathbb{Z}$ such that $1 - m \in \mathbb{Z}^+$. Then the function $R_{\Gamma, \mathfrak{p}}^{\kappa(m)}(z)$ is a modular form of weight 2κ for Γ , and is a cusp form in case $m < 0$.*

Proof. Let κ and m be as in the statement of the proposition. Since Γ has width one at infinity we may take Γ to be the scaling coset for Γ at the infinite cusp. Let $\Sigma_{\mathfrak{p}}$ be a scaling coset for Γ at $\mathfrak{p}$ and set $U = [\Sigma_{\mathfrak{p}}^{-1}]$. Then in the case that $\kappa > 1$ we have

$$R_{\Gamma, \mathfrak{p}}^{\kappa(m)}(z) = \sum_{[\chi] \in U} R_{[\chi]}^{\kappa(m)}(z) = \sum_{[\chi] \in U} e(-m [\chi \cdot z]) \text{jac}([\chi], z)^\kappa, \quad (3.4.6)$$

with the sum(s) in (3.4.6) converging absolutely for $z \in \mathbb{H}$. Let $\sigma \in G(\mathbb{Q})$. Then we have

$$\left(R_{\Gamma, \mathfrak{p}}^{\kappa(m)} \Big|^\kappa \sigma \right) (z) = \sum_{[\chi] \in U} \left(R_{[\chi]}^{\kappa(m)} \Big|^\kappa \sigma \right) (z) = \sum_{[\chi] \in U} R_{[\chi\sigma]}^{\kappa(m)}(z) = \sum_{[\chi] \in U\sigma} R_{[\chi]}^{\kappa(m)}(z) \quad (3.4.7)$$

by Lemma 3.4.1, with all sums absolutely convergent. Taking $\sigma \in \Gamma$ we see that $R_{\Gamma, \mathfrak{p}}^{\kappa(m)}(z)$ is an unrestricted modular form of weight 2κ for Γ since $\Sigma_{\mathfrak{p}}^{-1}\sigma = \Sigma_{\mathfrak{p}}^{-1}$ for $\sigma \in \Gamma$. Taking $\mathfrak{q} \in P_\Gamma$ and $\sigma \in \Sigma_{\mathfrak{q}}$ we see that Theorem 3.3.4 gives the Fourier expansion of $R_{\Gamma, \mathfrak{p}}^{\kappa(m)}(z)$ at $\mathfrak{q}$, in the sense of §2.7, since $\Sigma_{\mathfrak{p}}^{-1}\sigma = \Sigma_{\mathfrak{p}}^{-1}\Sigma_{\mathfrak{q}}$ for any $\sigma \in \Sigma_{\mathfrak{q}}$. We conclude that the (a priori unrestricted) modular form $R_{\Gamma, \mathfrak{p}}^{\kappa(m)}(z)$ is vanishing at every cusp $\mathfrak{q} \in P_\Gamma$, except possibly for $\mathfrak{q} = \mathfrak{p}$, and vanishes even there just when $m < 0$, and thus the claim is verified for $\kappa > 1$.

Consider now the case that $\kappa = 1$. We have

$$R_{\Gamma, \mathfrak{p}}^{1(m)}(z) = \lim_{K \rightarrow \infty} \sum_{[\chi] \in U_{\leq K}} R_{[\chi]}^{1(m)}(z) = \lim_{K \rightarrow \infty} \sum_{[\chi] \in U_{\leq K}} e(-m [\chi \cdot z]) \text{jac}([\chi], z), \quad (3.4.8)$$

and the limit is necessary for convergence. Let $\sigma \in G(\mathbb{Q})$ and consider the weight 2 action of σ

on $R_{\Gamma, \mathfrak{p}}^{1(m)}(z)$.

$$\left(R_{\Gamma, \mathfrak{p}}^{1(m)} \Big|_1 \sigma\right)(z) = \lim_{K \rightarrow \infty} \sum_{\chi \in U_{\leq K}} \left(R_{\chi}^{1(m)} \Big|_1 \sigma\right)(z) = \lim_{K \rightarrow \infty} \sum_{\chi \in U_{\leq K}} R_{\chi^\sigma}^{1(m)}(z). \quad (3.4.9)$$

Taking $\sigma \in \Gamma$ and applying Lemma 3.4.2 we see that $R_{\Gamma, \mathfrak{p}}^{1(m)}(z)$ is a(n unrestricted) modular form for Γ . Applying Lemma 3.4.2 in the case that $\sigma \in \Sigma_{\mathfrak{q}}$ for some cusp $\mathfrak{q} \in \mathcal{P}_{\Gamma}$, we see that the Fourier expansion of the function $R_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{1(m)}(z)$ is the expansion of $R_{\Gamma, \mathfrak{p}}^{1(m)}(z)$ at $\mathfrak{q}$ in the sense of §2.7, so Theorem 3.3.4 applies also when $\kappa = 1$, and thus $R_{\Gamma, \mathfrak{p}}^{1(m)}(z)$ is a modular form of weight 2 for Γ when $m = 0$, and a cusp form in case $m < 0$. This completes the proof. $\square$

We now present an analogue of Theorem 3.4.3 for the case that $1 - \kappa \in \mathbb{Z}^+$.

Theorem 3.4.4. *Let Γ be a group commensurable with $G(\mathbb{Z})$ that has width one at infinity and let $\mathfrak{p} \in \mathcal{P}_{\Gamma}$ be a cusp of Γ . Let $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$, and let $m \in \mathbb{Z}^+$. Then the function $R_{\Gamma, \mathfrak{p}}^{\kappa(m)}(z) + c_{\Gamma, \mathfrak{p}}^{\kappa}(m, 0)/2$ is an automorphic integral of weight 2κ for Γ .*

Proof. Let κ and m be as in the statement of the theorem. Since Γ has width one at infinity we take Γ to be the scaling coset for Γ at the infinite cusp. Let $\Sigma_{\mathfrak{p}}$ be a scaling coset for Γ at $\mathfrak{p}$ and set $U = \llbracket \Sigma_{\mathfrak{p}}^{-1}$. Let $\sigma \in G(\mathbb{Q})$. Using Lemma 3.4.1 we compute

$$\begin{aligned} \left(R_U^{\kappa(m)} \Big|^\kappa \sigma\right)(z) &= \lim_{K \rightarrow \infty} \sum_{\chi \in U_{\leq K}} R_{\chi^\sigma}^{\kappa(m)}(z) - m^{1-2\kappa} \left(J_{\infty, \llbracket \sigma}^\kappa P_{\llbracket \chi^\sigma}^{1-\kappa(-m)}\right)(z) \\ &= \lim_{K \rightarrow \infty} \sum_{\chi \in (U_{\leq K})\sigma} R_{\chi}^{\kappa(m)}(z) - m^{1-2\kappa} \left(J_{\infty, \llbracket \sigma}^\kappa P_{\llbracket \chi}^{1-\kappa(-m)}\right)(z) \end{aligned} \quad (3.4.10)$$

for the weight 2κ action of σ on $R_U^{\kappa(m)}(z) = R_{\Gamma, \mathfrak{p}}^{\kappa(m)}(z)$. Now Lemma 3.4.2 shows that

$$\left(R_U^{\kappa(m)} \Big|^\kappa \sigma\right)(z) = R_{U\sigma}^{\kappa(m)}(z) - m^{1-2\kappa} \lim_{K \rightarrow \infty} \sum_{\chi \in (U_{\leq K})\sigma} \left(J_{\infty, \llbracket \sigma}^\kappa P_{\llbracket \chi}^{1-\kappa(-m)}\right)(z), \quad (3.4.11)$$

suggesting that $R_U^{\kappa(m)}(z)$ might be an automorphic integral of weight 2κ for Γ with associated cusp form $m^{1-2\kappa} P_U^{1-\kappa(-m)}(z)$ (cf. §2.7). This is not accurate, however, since the limit in (3.4.11) does not in general commute with the integral operator $J_{\infty, \llbracket \sigma}^\kappa$. Rather, by the argument of

Lemma 4.4. in [Nie74], we have the identity

$$\begin{aligned} & \lim_{K \rightarrow \infty} \left(J_{\infty, \llbracket \sigma}^{\kappa} \left(\sum_{\llbracket \chi \in (U_{\leq K})^{\sigma}} P_{\llbracket \chi}^{1-\kappa(-m)} \right) \right) (z) - \left(J_{\infty, \llbracket \sigma}^{\kappa} \left(\lim_{K \rightarrow \infty} \sum_{\llbracket \chi \in (U_{\leq K})^{\sigma}} P_{\llbracket \chi}^{1-\kappa(-m)} \right) \right) (z) \\ &= \frac{m^{2\kappa-1}}{2} (c_S^{\kappa}(m, 0) \text{jac}(\sigma, z)^{\kappa} - c_{S'}^{\kappa}(m, 0)) \end{aligned} \quad (3.4.12)$$

where $S = \{\llbracket \chi \rrbracket \mid \llbracket \chi \rrbracket \in U^{\times}\}$ and $S' = \{\llbracket \chi \rrbracket \mid \llbracket \chi \rrbracket \in (U\sigma)^{\times}\}$. Lemma 3.3.1 plays a crucial rôle in the verification of (3.4.12). Applying (3.4.12) to (3.4.11) we find that

$$\left(R_U^{\kappa(m)} + \frac{1}{2} c_S^{\kappa}(m, 0) \right) \Big|_{\sigma}^{\kappa} = R_{U'}^{\kappa(m)} + \frac{1}{2} c_{S'}^{\kappa}(m, 0) - m^{1-2\kappa} J_{\infty, \llbracket \sigma}^{\kappa} P_{U'}^{1-\kappa(-m)} \quad (3.4.13)$$

for $U' = U\sigma$. Taking $\sigma \in \Gamma$, and applying the identity (4.4.29), we obtain

$$\left(R_{\Gamma, \mathbf{p}}^{\kappa(m)} + \frac{1}{2} c_{\Gamma, \mathbf{p}}^{\kappa}(m, 0) \right) \Big|_{\sigma}^{\kappa} = R_{\Gamma, \mathbf{p}}^{\kappa(m)} + \frac{1}{2} c_{\Gamma, \mathbf{p}}^{\kappa}(m, 0) - m^{1-2\kappa} J_{\infty, \llbracket \sigma}^{\kappa} P_{\Gamma, \mathbf{p}}^{1-\kappa(-m)} \quad (3.4.14)$$

from (3.4.13), demonstrating that the function $R_{\Gamma, \mathbf{p}}^{\kappa(m)}(z) + c_{\Gamma, \mathbf{p}}^{\kappa}(m, 0)/2$ is an unrestricted automorphic integral of weight 2κ for Γ with associated cusp form $m^{1-2\kappa} P_{\Gamma, \mathbf{p}}^{1-\kappa(-m)}(z)$. Letting $\mathbf{q} \in \mathbf{P}_{\Gamma}$ and applying (3.4.13) with $\sigma \in \Sigma_{\mathbf{q}}$ we obtain

$$\begin{aligned} R_{\Gamma, \mathbf{p}|\mathbf{q}}^{\kappa(m)} + \frac{1}{2} c_{\Gamma, \mathbf{p}|\mathbf{q}}^{\kappa}(m, 0) &= \left(R_{\Gamma, \mathbf{p}}^{\kappa(m)} + \frac{1}{2} c_{\Gamma, \mathbf{p}}^{\kappa}(m, 0) \right) \Big|_{\sigma}^{\kappa} + m^{1-2\kappa} J_{\infty, \llbracket \sigma}^{\kappa} P_{\Gamma, \mathbf{p}|\mathbf{q}}^{1-\kappa(-m)} \\ &= \left(R_{\Gamma, \mathbf{p}}^{\kappa(m)} + \frac{1}{2} c_{\Gamma, \mathbf{p}}^{\kappa}(m, 0) - m^{1-2\kappa} J_{\sigma, \infty}^{\kappa} P_{\Gamma, \mathbf{p}}^{1-\kappa(-m)} \right) \Big|_{\sigma}^{\kappa}, \end{aligned} \quad (3.4.15)$$

demonstrating that the Fourier expansion of the function $R_{\Gamma, \mathbf{p}|\mathbf{q}}^{\kappa(m)}(z) + c_{\Gamma, \mathbf{p}|\mathbf{q}}^{\kappa}(m, 0)/2$ is the expansion at $\mathbf{q}$ of the automorphic integral $R_{\Gamma, \mathbf{p}}^{\kappa(m)}(z) + c_{\Gamma, \mathbf{p}}^{\kappa}(m, 0)/2$, in the sense of §2.7. Now Theorem 3.3.2 confirms that $R_{\Gamma, \mathbf{p}|\mathbf{q}}^{\kappa(m)}(z) + c_{\Gamma, \mathbf{p}|\mathbf{q}}^{\kappa}(m, 0)/2$ is meromorphic at the cusps of Γ , and is thus an automorphic integral of weight 2κ for Γ . This completes the proof of the proposition. $\square$

We now specialize to the case that $\kappa = 0$. Our final goal in this section is to establish the Γ -invariance of the harmonic function $R_{\Gamma, \mathbf{p}}^{(m)}(z) - \bar{R}_{\Gamma, \mathbf{p}}^{(m)}(z)$ for $m \in \mathbb{Z}^+$ in case Γ has width one at infinity. For this we utilize the spectral theory of the hyperbolic Laplacian $\Delta = (z - \bar{z})^2 \partial_z \partial_{\bar{z}}$.

Consider the *Whittaker functions* $V(z, s)$ and $W(z, s)$, defined by setting

$$V(z, s) = e(z) \Gamma(s) \Phi(s, 2s, z - \bar{z}) (2\pi \mathbf{i}(\bar{z} - z))^s, \quad (3.4.16)$$

$$W(z, s) = \frac{\sin(\pi s)}{\sin(2\pi s)} (V(z, s) - V(z, 1 - s)), \quad (3.4.17)$$

where the function Φ is defined in (2.1.3). Then $V(z, s)$ and $W(z, s)$ are eigenfunctions for Δ with eigenvalue $s(1-s)$, and the identity $V(z, 1) = e(\bar{z}) - e(z)$ hints at a connection with the right hand side of (3.3.9). Define the *free space Green's function* $G(w, z, s)$ by setting

$$G(w, z, s) = \frac{\Gamma(s)^2}{\Gamma(2s)} {}_2F_1(s, s; 2s; h) h^s = \sum_{k \in \mathbb{N}} \frac{\Gamma(k+s)^2}{\Gamma(k+2s)\Gamma(k+1)} h^{k+s}, \quad (3.4.18)$$

where ${}_2F_1(a, b; c; x)$ denotes the Gauss hypergeometric function, and $h = h(w, z)$ is given as follows, and depends only on the hyperbolic distance between w and z .

$$h(w, z) = \frac{(w - \bar{w})(z - \bar{z})}{(w - z)(\bar{w} - \bar{z})} = \frac{4\Im(w)\Im(z)}{|w - z|^2} \quad (3.4.19)$$

Then the function $z \mapsto G(w, z, s)$ satisfies $\Delta G(w, z, s) = s(1-s)G(w, z, s)$ for $z \neq w$, and the integral operator with kernel $G(w, z, s)$ furnishes a right-inverse to the differential operator $f \mapsto (\Delta - \lambda)f$ when $\lambda = s(1-s)$.

Let Γ be a group commensurable with $G(\mathbb{Z})$, let $\mathfrak{p}$ and $\mathfrak{q}$ be cusps of Γ , and let $\Sigma_{\mathfrak{p}}$ and $\Sigma_{\mathfrak{q}}$ be scaling cosets for Γ at $\mathfrak{p}$ and $\mathfrak{q}$, respectively. Define $G_{\Gamma, \mathfrak{p}|\mathfrak{q}}(w, z, s)$ by setting

$$G_{\Gamma, \mathfrak{p}|\mathfrak{q}}(w, z, s) = \sum_{\chi \in \Sigma_{\mathfrak{p}}^{-1}\Sigma_{\mathfrak{q}}} G(w, \chi \cdot z, s). \quad (3.4.20)$$

The series (3.4.20) converges absolutely when $\Re(s) > 1$ and $\Gamma \cdot w \neq \Gamma \cdot z$, and the functions $w \mapsto G_{\Gamma, \mathfrak{p}|\mathfrak{q}}(w, z, s)$ and $z \mapsto G_{\Gamma, \mathfrak{p}|\mathfrak{q}}(w, z, s)$ are invariant for the actions of $\Gamma^{\mathfrak{p}}$ and $\Gamma^{\mathfrak{q}}$, respectively. According to [Hej83] (see also [Iwa02]) we have the Fourier–Whittaker series expansion

$$G_{\Gamma, \mathfrak{p}|\mathfrak{q}}(w, z, s) = \sum_{m \in \mathbb{Z}^+} G_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(-m)}(z, s) W(-m\bar{w}, s) + G_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(0)}(z, s) \Im(w)^{1-s} + \sum_{m \in \mathbb{Z}^+} G_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(m)}(z, s) W(mw, s), \quad (3.4.21)$$

converging absolutely when $\Re(s) > 1$ so long as $\Im(w) > \Im(z)$ and $\Im(z)\Im(w) > \varrho_{\Gamma, \mathfrak{p}|\mathfrak{q}}$, for a certain constant $\varrho_{\Gamma, \mathfrak{p}|\mathfrak{q}}$ depending only on Γ and $\mathfrak{p}$ and $\mathfrak{q}$, where the coefficient functions $G_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(m)}(z, s)$ are $\Gamma^{\mathfrak{q}}$ -invariant, and themselves admit the following series expansions of Poincaré

type, converging absolutely for $\Re(s) > 1$ and $\Im(z) > 0$.

$$G_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(-m)}(z, s) = \frac{1}{m} \sum_{\llbracket \chi \in \llbracket \Sigma_{\mathfrak{p}}^{-1} \Sigma_{\mathfrak{q}} \rrbracket} V(m \llbracket \chi \cdot z, s) \quad (3.4.22)$$

$$G_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(0)}(z, s) = \frac{4\pi}{2s-1} \sum_{\llbracket \chi \in \llbracket \Sigma_{\mathfrak{p}}^{-1} \Sigma_{\mathfrak{q}} \rrbracket} \Im(\llbracket \chi \cdot z \rrbracket^s) \quad (3.4.23)$$

$$G_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(m)}(z, s) = \frac{1}{m} \sum_{\llbracket \chi \in \llbracket \Sigma_{\mathfrak{p}}^{-1} \Sigma_{\mathfrak{q}} \rrbracket} V(-m \llbracket \chi \cdot \bar{z}, s) \quad (3.4.24)$$

Let $m \in \mathbb{Z}^+$. Then we have the following analogue of (3.4.21) for the Green's function coefficient $G_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(m)}(z, s)$.

$$\begin{aligned} G_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(m)}(z, s) &= \delta_{\Gamma, \mathfrak{p}|\mathfrak{q}} \frac{1}{m} V(-m \bar{z}, s) + \\ &\sum_{n \in \mathbb{Z}^+} g_{\Gamma, \mathfrak{p}|\mathfrak{q}}(m, -n, s) W(-n \bar{z}, s) + g_{\Gamma, \mathfrak{p}|\mathfrak{q}}(m, 0, s) \Im(z)^{1-s} + \sum_{n \in \mathbb{Z}^+} g_{\Gamma, \mathfrak{p}|\mathfrak{q}}(m, n, s) W(nz, s) \end{aligned} \quad (3.4.25)$$

This identity (3.4.25) is again reminiscent of the right hand side of (3.3.9). The coefficients in (3.4.25) may be defined via the following formulas, in which we assume $m, n \in \mathbb{Z}^+$.

$$\begin{aligned} g_{\Gamma, \mathfrak{p}|\mathfrak{q}}(m, -n, s) &= \\ \frac{1}{m^{1-s} n^s} \sum_{\llbracket \chi \rrbracket \in S} \lim_{k \in \mathbb{N}} \sum e(-m \llbracket \chi \rrbracket \cdot \infty) e(-n \llbracket \chi^{-1} \rrbracket \cdot \infty) (4\pi^2 \varrho \llbracket \chi \rrbracket)^{k+s} m^{(k)} n^{(k+2s-1)} \end{aligned} \quad (3.4.26)$$

$$g_{\Gamma, \mathfrak{p}|\mathfrak{q}}(m, 0, s) = \frac{1}{m^{1-s}} \frac{4\pi}{2s-1} \sum_{\llbracket \chi \rrbracket \in S} \lim_{k \in \mathbb{N}} \sum e(-m \llbracket \chi \rrbracket \cdot \infty) \varrho \llbracket \chi \rrbracket^s \frac{\pi^s}{\Gamma(s)} \quad (3.4.27)$$

$$\begin{aligned} g_{\Gamma, \mathfrak{p}|\mathfrak{q}}(m, n, s) &= \\ \frac{1}{m^{1-s} n^s} \sum_{\llbracket \chi \rrbracket \in S} \lim_{k \in \mathbb{N}} \sum e(-m \llbracket \chi \rrbracket \cdot \infty) e(n \llbracket \chi^{-1} \rrbracket \cdot \infty) (-1)^k (4\pi^2 \varrho \llbracket \chi \rrbracket)^{k+s} m^{(k)} n^{(k+2s-1)} \end{aligned} \quad (3.4.28)$$

Comparing with Lemma (3.2.4) we obtain a direct relationship between the coefficient functions $g_{\Gamma, \mathfrak{p}|\mathfrak{q}}(m, n, s)$ arising from the Green's function, and the coefficient functions $c_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa}(m, n)$ of §3.2 which arise from the Rademacher sums.

Proposition 3.4.5. *Let Γ be a group commensurable with $G(\mathbb{Z})$, let $\mathfrak{p}, \mathfrak{q} \in \mathcal{P}_{\Gamma}$, and let $m, \kappa \in$*

$\mathbb{Z}^+$. Then we have the following identities for $n \in \mathbb{Z}^+$.

$$c_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{1-\kappa}(m, -n) = (-1)^{1-\kappa} m^\kappa n^{1-\kappa} g_{\Gamma, \mathfrak{p}|\mathfrak{q}}(m, -n, \kappa) \quad (3.4.29)$$

$$c_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{1-\kappa}(m, 0) = (-1)^{1-\kappa} m^\kappa g_{\Gamma, \mathfrak{p}|\mathfrak{q}}(m, 0, \kappa) \quad (3.4.30)$$

$$c_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{1-\kappa}(m, n) = (-1)^{1-\kappa} m^\kappa n^{1-\kappa} g_{\Gamma, \mathfrak{p}|\mathfrak{q}}(m, n, \kappa) \quad (3.4.31)$$

Taking $\kappa = 1$ in Proposition 3.4.5 we obtain the identity $c_{\Gamma, \mathfrak{p}|\mathfrak{q}}(m, n) = m g_{\Gamma, \mathfrak{p}|\mathfrak{q}}(m, n, 1)$, which, by Proposition 3.3.9, the series expansion (3.4.25), and the identity $W(z, 1) = e(z)$, may be reformulated as

$$R_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(m)}(z) - \bar{R}_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(m)}(z) = m G_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(m)}(z, 1). \quad (3.4.32)$$

The Γ^q -invariance of the difference $R_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(m)}(z) - \bar{R}_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(m)}(z)$ of classical and conjugate Rademacher sums of weight 0 now follows from the Γ^q -invariance of the Green's function coefficient $G_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(m)}(z, s)$ at $s = 1$ (cf. [Nie73]). In particular, we have the following result.

Theorem 3.4.6. *Let Γ be a group commensurable with $G(\mathbb{Z})$ that has width one at infinity, let $\mathfrak{p} \in P_\Gamma$ be a cusp of Γ , and let $m \in \mathbb{Z}^+$. Then the harmonic function $R_{\Gamma, \mathfrak{p}}^{(m)}(z) - \bar{R}_{\Gamma, \mathfrak{p}}^{(m)}(z)$ is Γ -invariant.*

4 Modified Rademacher sums

We have seen in §3.4 that the classical Rademacher sum $R_{\Gamma, \mathfrak{p}}^{\kappa(m)}(z)$ of §3 is an automorphic integral for Γ only after the addition of a particular constant function, which does not generally vanish. In this section we introduce a modification of Rademacher's construction via which the correct constant term appears naturally. Our approach employs an analytic continuation of the component functions defining the classical Rademacher sums, and entails the assignment of a Dirichlet series to each triple $(\Gamma, \mathfrak{p}, \mathfrak{q})$ where Γ is a group commensurable with the modular group and $\mathfrak{p}, \mathfrak{q} \in P_\Gamma$ are cusps for Γ .

4.1 Construction

For $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$, for $m \in \mathbb{Z}^+$ and for $[\chi] \in [G(\mathbb{Q})]^\times$ we define the *continued Rademacher component function*, denoted $(z, s) \mapsto T_{[\chi]}^{\kappa(m)}(z, s)$, by setting

$$T_{[\chi]}^{\kappa(m)}(z, s) = e(-m [\chi \cdot z]) t^\kappa(m, [\chi], z, s) \text{jac}([\chi], z)^\kappa, \quad (4.1.1)$$

where $t^\kappa(m, \llbracket \chi, z, s)$ is the *continued Rademacher regularization factor of weight 2κ* , which is in turn given by

$$t^\kappa(m, \llbracket \chi, z, s) = \Phi(s - 2\kappa, 1 + s - 2\kappa, m \llbracket \chi \cdot z - m \llbracket \chi \cdot \infty)(2\pi i(m \llbracket \chi \cdot \infty - m \llbracket \chi \cdot z))^{s-2\kappa} \quad (4.1.2)$$

in case $\llbracket \chi \in \llbracket G(\mathbb{Q})^\times$, and $t^\kappa(m, \llbracket \chi, z, s) = 1$ otherwise. We also define a function $s \mapsto T_{0, \llbracket \chi}^{\kappa(m)}(s)$ by setting

$$\begin{aligned} T_{0, \llbracket \chi}^{\kappa(m)}(s) &= e(-m \llbracket \chi \cdot \infty) (2\pi i(m \llbracket \chi \cdot \infty - m \llbracket \chi \cdot 0))^{(s-2\kappa)} \text{jac}(\llbracket \chi, 0)^\kappa \\ &\quad - e(s/2) e(-m \llbracket \chi \cdot \infty) (2\pi i(m \llbracket \chi \cdot 0 - m \llbracket \chi \cdot \infty))^{(s-2\kappa)} \text{jac}(\llbracket \chi, 0)^\kappa \end{aligned} \quad (4.1.3)$$

in case $\llbracket \chi \in \llbracket G(\mathbb{Q})^{\times \times}$, and by setting $T_{0, \llbracket \chi}^{\kappa(m)}(s) = 0$ otherwise, and we define the *modified continued Rademacher component function*, denoted $(z, s) \mapsto Q_{\llbracket \chi}^{\kappa(m)}(z, s)$, by subtracting $T_{0, \llbracket \chi}^{\kappa(m)}(z, s)$ from $T_{\llbracket \chi}^{\kappa(m)}(z, s)$.

$$Q_{\llbracket \chi}^{\kappa(m)}(z, s) = T_{\llbracket \chi}^{\kappa(m)}(z, s) - T_{0, \llbracket \chi}^{\kappa(m)}(z, s) \quad (4.1.4)$$

Now for $U \subset \llbracket G(\mathbb{Q})$, for $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$, and for $m \in \mathbb{Z}^+$, we define the *continued Rademacher sum* $T_U^{\kappa(m)}(z, s)$, and the *modified continued Rademacher sum* $Q_U^{\kappa(m)}(z, s)$, by setting

$$T_U^{\kappa(m)}(z, s) = \sum_{\llbracket \chi \in U} T_{\llbracket \chi}^{\kappa(m)}(z, s), \quad (4.1.5)$$

$$Q_U^{\kappa(m)}(z, s) = \sum_{\llbracket \chi \in U} Q_{\llbracket \chi}^{\kappa(m)}(z, s), \quad (4.1.6)$$

and we define $T_U^{\kappa(m)}(z)$ and $Q_U^{\kappa(m)}(z)$ by taking the limit as s tends to 1 through the region $\Re(s) > 1$ in $T_U^{\kappa(m)}(z, s)$ and $Q_U^{\kappa(m)}(z, s)$, respectively.

$$T_U^{\kappa(m)}(z) = \lim_{s \rightarrow 1^+} T_U^{\kappa(m)}(z, s) \quad (4.1.7)$$

$$Q_U^{\kappa(m)}(z) = \lim_{s \rightarrow 1^+} Q_U^{\kappa(m)}(z, s) \quad (4.1.8)$$

The function $Q_U^{\kappa(m)}(z)$ is the most important from the point of view of automorphy for subgroups of $G(\mathbb{Q})$, while the function $T_U^{\kappa(m)}(z)$ plays a special rôle in the critical case that $\kappa = 0$. We call $Q_U^{\kappa(m)}(z)$ the *modified Rademacher sum of weight 2κ and order m associated to U* , and we call $T_U^{\kappa(m)}(z)$ the *normalized Rademacher sum of weight 2κ and order m associated to U* .

Observe that we recover the Rademacher component function $R_{\llbracket \chi}^{\kappa(m)}(z)$ by taking $s = 1$

in either the continued Rademacher component function $T_{\mathbb{X}}^{\kappa(m)}(z, s)$ or the modified continued Rademacher component function $Q_{\mathbb{X}}^{\kappa(m)}(z, s)$.

Lemma 4.1.1. *Let $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$, let $m \in \mathbb{Z}^+$, and let $\mathbb{X} \in \mathbb{G}(\mathbb{Q})$. Then we have $T_{\mathbb{X}}^{\kappa(m)}(z, 1) = R_{\mathbb{X}}^{\kappa(m)}(z)$ and $T_{0, \mathbb{X}}^{\kappa(m)}(1) = 0$, and hence $Q_{\mathbb{X}}^{\kappa(m)}(z, 1) = R_{\mathbb{X}}^{\kappa(m)}(z)$.*

Despite the result of Lemma 4.1.1, it is generally not the case that the modified Rademacher sum $Q_U^{\kappa(m)}(z)$ and the classical Rademacher sum $R_U^{\kappa(m)}(z)$ coincide (cf. Proposition 4.4.10).

We typically take U to be of the form $U = \llbracket \Sigma_{\mathbf{p}}^{-1} \Sigma_{\mathbf{q}} \rrbracket$ where $\{\Sigma_{\mathbf{p}} \mid \mathbf{p} \in \mathbf{P}_{\Gamma}\}$ is a system of scaling cosets (cf. §2.6) for some group Γ commensurable with the modular group, and $\mathbf{p}, \mathbf{q} \in \mathbf{P}_{\Gamma}$ are cusps of Γ . In this case we write $Q_{\Gamma, \mathbf{p}|\mathbf{q}}^{\kappa(m)}(z)$ for $Q_U^{\kappa(m)}(z)$ and $T_{\Gamma, \mathbf{p}|\mathbf{q}}^{\kappa(m)}(z)$ for $T_U^{\kappa(m)}(z)$, suppressing the choice of scaling cosets from notation. A change in the choice of scaling cosets $\Sigma_{\mathbf{p}}$ and $\Sigma_{\mathbf{q}}$ replaces $Q_{\Gamma, \mathbf{p}|\mathbf{q}}^{\kappa(m)}(z)$ with a function of the form $e(\alpha)Q_{\Gamma, \mathbf{p}|\mathbf{q}}^{\kappa(m)}(z + \beta)$ for some $\alpha, \beta \in \mathbb{Q}$, and similarly for $T_{\Gamma, \mathbf{p}|\mathbf{q}}^{\kappa(m)}(z)$.

In the case that $\mathbf{p}$ or $\mathbf{q}$ is the *infinite cusp* $\Gamma \cdot \infty$ we omit it from notation, writing $Q_{\Gamma, \mathbf{p}}^{\kappa(m)}(z)$ for $Q_{\Gamma, \mathbf{p}|\Gamma \cdot \infty}^{\kappa(m)}(z)$, and $Q_{\Gamma, \mathbf{q}}^{\kappa(m)}(z)$ for $Q_{\Gamma, \Gamma \cdot \infty|\mathbf{q}}^{\kappa(m)}(z)$, and similarly for the functions $T_{\Gamma, \mathbf{p}|\mathbf{q}}^{\kappa(m)}(z)$. The functions $Q_{\Gamma, \mathbf{p}}^{\kappa(m)}(z)$ and $T_{\Gamma, \mathbf{p}}^{\kappa(m)}(z)$ are the most important, for we shall see in §4.5 that $Q_{\Gamma, \mathbf{p}}^{\kappa(m)}(z)$ is an automorphic integral of weight 2κ for Γ with a single pole at the cusp $\mathbf{p}$ in case Γ has width one at infinity, and we will see in §4.4 that $Q_{\Gamma, \mathbf{p}}^{\kappa(m)}(z)$ and $T_{\Gamma, \mathbf{p}}^{\kappa(m)}(z)$ differ only by a constant function. We call $Q_{\Gamma, \mathbf{p}}^{\kappa(m)}(z)$ the *modified Rademacher sum of weight 2κ and order m associated to Γ at the cusp $\mathbf{p}$* , and we call $T_{\Gamma, \mathbf{p}}^{\kappa(m)}(z)$ the *normalized Rademacher sum of weight 2κ and order m associated to Γ at the cusp $\mathbf{p}$* . We shall see also in §4.5 that the function $Q_{\Gamma, \mathbf{p}|\mathbf{q}}^{\kappa(m)}(z)$ may be regarded as encoding the behavior of $Q_{\Gamma, \mathbf{p}}^{\kappa(m)}(z)$ at the cusp $\mathbf{q}$.

Just as for the classical Rademacher sums (cf. §3.1) we have the result that every modified Rademacher sum $Q_{\Gamma, \mathbf{p}|\mathbf{q}}^{\kappa(m)}(z)$ may be expressed in the form $Q_{\Gamma', \mathbf{p}'|\mathbf{q}'}^{\kappa(m)}(z)$ for some group Γ' with width one at infinity, and some cusp $\mathbf{p}'$ of Γ' , and similarly for the normalized Rademacher sums $T_{\Gamma, \mathbf{p}|\mathbf{q}}^{\kappa(m)}(z)$. Precisely, we have

$$Q_{\Gamma, \mathbf{p}|\mathbf{q}}^{\kappa(m)}(z) = Q_{\Gamma^{\mathbf{q}}, \mathbf{p}^{\mathbf{q}}}^{\kappa(m)}(z) \quad (4.1.9)$$

$$T_{\Gamma, \mathbf{p}|\mathbf{q}}^{\kappa(m)}(z) = T_{\Gamma^{\mathbf{q}}, \mathbf{p}^{\mathbf{q}}}^{\kappa(m)}(z) \quad (4.1.10)$$

subject to the understanding that the data defining the right hand sides of (4.1.9) and (4.1.10) is related to the data defining the left hand sides of (4.1.9) and (4.1.10) by $\Gamma^{\mathbf{q}} = \Sigma_{\mathbf{q}}^{-1} \Sigma_{\mathbf{q}}$ and $\mathbf{p}^{\mathbf{q}} = \Sigma_{\mathbf{q}}^{-1} \cdot \mathbf{p}$ and $\Sigma_{\mathbf{p}^{\mathbf{q}}} = \Sigma_{\mathbf{q}}^{-1} \Sigma_{\mathbf{p}}$.

Our primary interest in this article is in the distinguished case that $\kappa = 0$. In order to simplify notation, and maintain consistency with the notation of §1, we write $Q_{\Gamma, \mathbf{p}|\mathbf{q}}^{(m)}(z)$ as a shorthand for $Q_{\Gamma, \mathbf{p}|\mathbf{q}}^{0(m)}(z)$, and we write $T_{\Gamma, \mathbf{p}|\mathbf{q}}^{(m)}(z)$ as a shorthand for $T_{\Gamma, \mathbf{p}|\mathbf{q}}^{0(m)}(z)$.

4.2 Coefficients

In order to recover explicit expressions of the Fourier coefficients of the modified Rademacher sums we employ the following generalizations of $\text{Bl}_{[\chi]}^\kappa(m, n)$ and $c_S^\kappa(m, n)$ for $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$, and for $m, n \in \mathbb{Z}^+$. We define functions $s \mapsto \text{Bl}_{[\chi]}^\kappa(m, n, s)$, for $1 - \kappa, m, n \in \mathbb{Z}^+$, by setting

$$\text{Bl}_{[\chi]}^\kappa(m, n) = (-1)^\kappa \sum_{k \in \mathbb{N}} (4\pi^2)^{k-\kappa+s} \varrho([\chi])^{k-2\kappa+s} m^{(k-2\kappa+s)} n^{(k+s-1)}, \quad (4.2.1)$$

so that $\text{Bl}_{[\chi]}^\kappa(m, n, 1) = \text{Bl}_{[\chi]}^\kappa(m, n)$. We then define the *continued coefficient function* $s \mapsto c_S^\kappa(m, n, s)$, for $1 - \kappa, m, n \in \mathbb{Z}^+$, by setting

$$c_S^\kappa(m, n, s) = \sum_{[\chi] \in S} \text{Kl}_{[\chi]}(m, n, \kappa) \text{Bl}_{[\chi]}^\kappa(m, n, s). \quad (4.2.2)$$

Lemma 4.2.1. *The functions $s \mapsto c_S^\kappa(m, n, s)$ admit the following series representation.*

$$\begin{aligned} c_S^\kappa(m, n, s) &= (-1)^\kappa \lim_{[\chi] \in S} \sum_{k \in \mathbb{N}} \sum_{k \in \mathbb{N}} e(-m[\chi] \cdot \infty) e(-n[\chi^{-1}] \cdot \infty) (4\pi^2 \varrho([\chi]))^{k-\kappa+s} m^{(k-2\kappa+s)} n^{(k+s-1)} \end{aligned} \quad (4.2.3)$$

We have $c_S^\kappa(m, n, 1) = c_S^\kappa(m, n)$.

We usually take S to be of the form $S = [\Sigma_p^{-1} \Sigma_q]^\times$ where Σ_p and Σ_q are scaling cosets (cf. §2.6) for a group Γ commensurable with $G(\mathbb{Z})$ at cusps $\mathfrak{p}, \mathfrak{q} \in P_\Gamma$. We write $c_{\Gamma, \mathfrak{p}|\mathfrak{q}}^\kappa(m, n, s)$ for $c_S^\kappa(m, n, s)$ in the case that $S = [\Sigma_p^{-1} \Sigma_q]^\times$.

Utilizing the continued coefficient functions $s \mapsto c_S^\kappa(m, n, s)$ we generalize the function $F_S^{\kappa(m)}(z)_{\text{van}}$ in case $1 - \kappa, m \in \mathbb{Z}^+$ by setting

$$F_S^{\kappa(m)}(z, s)_{\text{van}} = \sum_{n \in \mathbb{Z}^+} c_S^\kappa(m, n, s) e(nz). \quad (4.2.4)$$

We write $F_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z, s)_{\text{van}}$ for $F_S^{\kappa(m)}(z, s)_{\text{van}}$ in the case that $S = [\Sigma_p^{-1} \Sigma_q]^\times$ for Γ a group commensurable with $G(\mathbb{Z})$ and for $\Sigma_p, \Sigma_q \in \Gamma \backslash G(\mathbb{Q})$ scaling cosets for Γ at cusps $\mathfrak{p}, \mathfrak{q} \in P_\Gamma$. We write $F_S^{(m)}(z, s)_{\text{van}}$ as a shorthand for $F_S^{0(m)}(z, s)_{\text{van}}$, and interpret the notation $F_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(m)}(z, s)_{\text{van}}$ similarly.

We have the following analogue of Proposition 3.2.7 for the functions $F_S^{\kappa(m)}(z, s)_{\text{van}}$ in case S is of the form $[\Sigma_p^{-1} \Sigma_q]^\times$.

Proposition 4.2.2. *Let Γ be a group commensurable with $G(\mathbb{Z})$, and let $\mathfrak{p}, \mathfrak{q} \in P_\Gamma$ be cusps of*

Γ . Let $\kappa, m \in \mathbb{Z}$ such that $1 - \kappa, m \in \mathbb{Z}^+$. Then the series

$$F_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z, s)_{\text{van}} = \sum_{n \in \mathbb{Z}^+} c_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa}(m, n, s) e(nz) \quad (4.2.5)$$

converges absolutely for $z \in \mathbb{H}$ and $\Re(s) \geq 1$. In particular, the assignment $z \mapsto F_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z, s)_{\text{van}}$ is a holomorphic function on $\mathbb{H}$ whenever $1 - \kappa, m \in \mathbb{Z}^+$ and $\Re(s) \geq 1$.

4.3 Dirichlet series

Given $[\chi] \in [G(\mathbb{Q})]^\times$, and $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$, and $m \in \mathbb{Z}^+$, we define a Dirichlet series $Z_{[\chi]}^{\kappa(m)}(s)$ by setting

$$Z_{[\chi]}^{\kappa(m)}(s) = (-1)^{1-\kappa} \sum_{n \in \mathbb{Z}^\times} e(-m [\chi] \cdot \infty) e(-n [\chi^{-1}] \cdot \infty) (4\pi^2 \varrho([\chi]))^{1-s-\kappa} m^{(1-s-2\kappa)} n^{(-s)}. \quad (4.3.1)$$

This series (4.3.1) converges absolutely in the half plane $\Re(s) > 1$, and admits an analytic continuation to all of $\mathbb{C}$. Given $S \subset [G(\mathbb{Q})]^\times$ we define the *zeta function of weight 2κ and order m associated to S* , to be denoted $Z_S^{\kappa(m)}(s)$, by setting

$$Z_S^{\kappa(m)}(s) = \sum_{[\chi] \in S} Z_{[\chi]}^{\kappa(m)}(s). \quad (4.3.2)$$

In order to describe the analytic continuation of the zeta function $s \mapsto Z_S^{\kappa(m)}(s)$ explicitly, we formulate an identity which expresses it in terms of the Hurwitz zeta function. In preparation for this we define a function $s \mapsto \tilde{Z}_{[\chi]}^{\kappa(m)}(s)$, for each $[\chi] \in [G(\mathbb{Q})]^\times$, by setting

$$\tilde{Z}_{[\chi]}^{\kappa(m)}(s) = e(-m [\chi] \cdot \infty) \varrho([\chi])^{s-\kappa} (2\pi i m)^{(s-2\kappa)} (e(s/2) - e(-s/2)) \zeta(1 - \alpha_{[\chi]}, s) \quad (4.3.3)$$

where $\zeta(\alpha, s)$ is defined by $\zeta(\alpha, s) = \sum_{n \in \mathbb{N}} (n + \alpha)^{-s}$ for $\Re(\alpha) > 0$ and $\Re(s) > 1$, and $\alpha_{[\chi]} \in \mathbb{Q}$ is chosen so that

$$0 \leq \alpha_{[\chi]} < 1, \quad \alpha_{[\chi]} + \mathbb{Z} = -\chi^{-1} \cdot \infty + \mathbb{Z}. \quad (4.3.4)$$

We then define $\tilde{Z}_S^{\kappa(m)}(s)$ by setting $\tilde{Z}_S^{\kappa(m)}(s) = \sum_{[\chi] \in S} \tilde{Z}_{[\chi]}^{\kappa(m)}(s)$.

Lemma 4.3.1. *We have $Z_S^{\kappa(m)}(1-s) = \tilde{Z}_S^{\kappa(m)}(s)$.*

Proof. Let $[\chi] \in [G(\mathbb{Q})]^\times$ and choose $\alpha_{[\chi]} \in \mathbb{Q}$ such that $0 \leq \alpha_{[\chi]} < 1$ and $\alpha_{[\chi]} + \mathbb{Z} = -\chi^{-1} \cdot \infty + \mathbb{Z}$ for any $\chi \in [\chi]$. We will show that $Z_{[\chi]}^{\kappa(m)}(s) = \tilde{Z}_{[\chi]}^{\kappa(m)}(1-s)$. Observe that we

have

$$\sum_{n \in \mathbb{Z}^\times} e(-n \llbracket \chi^{-1} \rrbracket \cdot \infty) n^{(-s)} = \frac{1}{\Gamma(1-s)} \left(F(-\llbracket \chi^{-1} \rrbracket \cdot \infty, s) + e(-s/2) F(\llbracket \chi^{-1} \rrbracket \cdot \infty, s) \right), \quad (4.3.5)$$

where $F(\alpha, s)$ denotes the periodic zeta function, defined by $F(\alpha, s) = \sum_{n \in \mathbb{Z}^+} e(n\alpha) n^{-s}$ for $\Re(s) > 1$. The *Hurwitz relation*, which may be stated in the form

$$F(\alpha, s) + e(-s/2) F(-\alpha, s) = \frac{(-2\pi i)^s}{\Gamma(s)} \zeta(1-\alpha, 1-s) \quad (4.3.6)$$

(cf. [KR01]) implies the identity

$$\sum_{n \in \mathbb{Z}^\times} e(-n \llbracket \chi^{-1} \rrbracket \cdot \infty) n^{(-s)} = \frac{(-2\pi i)^s}{\Gamma(1-s)\Gamma(s)} \zeta(1-\alpha_{\llbracket \chi \rrbracket}, 1-s), \quad (4.3.7)$$

so the expression (4.3.1) may be reformulated as

$$Z_{\llbracket \chi \rrbracket}^{\kappa(m)}(s) = (-1)^{1-\kappa} e(-m \llbracket \chi \rrbracket \cdot \infty) (4\pi^2 \varrho \llbracket \chi \rrbracket)^{1-s-\kappa} m^{(1-s-2\kappa)} \frac{(-2\pi i)^s}{\Gamma(1-s)\Gamma(s)} \zeta(1-\alpha_{\llbracket \chi \rrbracket}, 1-s). \quad (4.3.8)$$

We obtain the coincidence of the right hand side of (4.3.8) with $\tilde{Z}_{\llbracket \chi \rrbracket}^{\kappa(m)}(1-s)$ by applying the functional equation for the Gamma function, which may be expressed as

$$(e(s/2) - e(-s/2)) \Gamma(s) \Gamma(1-s) = 2\pi i \quad (4.3.9)$$

This completes the proof of the claim. $\square$

We write $Z_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(s)$ for $Z_S^{\kappa(m)}(s)$ in the case that $S = \llbracket \Sigma_{\mathfrak{p}}^{-1} \Sigma_{\mathfrak{q}} \rrbracket^\times$ for Γ a group commensurable with $G(\mathbb{Z})$ and for $\Sigma_{\mathfrak{p}}, \Sigma_{\mathfrak{q}} \in \Gamma \backslash G(\mathbb{Q})$ scaling cosets for Γ at cusps $\mathfrak{p}, \mathfrak{q} \in \mathcal{P}_\Gamma$.

4.4 Convergence

Our main objective in this section is to furnish explicit expressions for the Fourier expansions of the functions $z \mapsto T_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z, s)$ and $z \mapsto Q_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z, s)$, defined by the continued Rademacher sums and modified continued Rademacher sums, respectively. This will yield the Fourier expansions of the modified Rademacher sums $Q_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z)$ and the normalized Rademacher sums $T_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z)$. In preparation for the derivation of these expansions we verify the absolute convergence of the series defining $T_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z, s)$ and $Q_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z, s)$ in case $\Re(s) > 1$.

Proposition 4.4.1. *Let Γ be a group commensurable with $G(\mathbb{Z})$ and let $\mathfrak{p}, \mathfrak{q} \in P_\Gamma$ be cusps of Γ . Let $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$, and let $m \in \mathbb{Z}^+$. Then the continued Rademacher sum*

$$T_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z, s) = \sum_{\llbracket \chi \in \llbracket \Sigma_{\mathfrak{p}}^{-1} \Sigma_{\mathfrak{q}}} T_{\llbracket \chi}^{\kappa(m)}(z, s) \quad (4.4.1)$$

is absolutely convergent for $z \in \mathbb{H}$ and $\Re(s) > 1$.

Proof. We begin by observing that we have

$$T_{\llbracket \chi}^{\kappa(m)}(z, s) = e(-m \llbracket \chi \cdot \infty) e(m \llbracket \chi \cdot \infty - m \llbracket \chi \cdot z, s - 2\kappa) \text{jac}(\llbracket \chi, z)^\kappa \quad (4.4.2)$$

in case $\llbracket \chi \in \llbracket G(\mathbb{Q})^\times$, and $T_{\llbracket \chi}^{\kappa(m)}(z, s) = e(-m \llbracket \chi \cdot z) \text{jac}(\llbracket \chi, z)^\kappa$ otherwise, where $e(z, s)$ denotes the generalized exponential function of (2.1.6). This identity (4.4.2) follows from an application of the *Kummer transformation* $\Phi(a, b, z) = e(z)\Phi(b - a, b, -z)$ to the expression (4.1.1) defining $T_{\llbracket \chi}^{\kappa(m)}(z, s)$. The absolute convergence of (4.4.2) now follows from standard estimates. $\square$

Proposition 4.4.2. *Let Γ be a group commensurable with $G(\mathbb{Z})$ and let $\mathfrak{p}, \mathfrak{q} \in P_\Gamma$ be cusps of Γ . Let $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$, and let $m \in \mathbb{Z}^+$. Then the sum*

$$\sum_{\llbracket \chi \in \llbracket \Sigma_{\mathfrak{p}}^{-1} \Sigma_{\mathfrak{q}}} T_{0, \llbracket \chi}^{\kappa(m)}(s) \quad (4.4.3)$$

is absolutely convergent for $z \in \mathbb{H}$ and $\Re(s) > 1$.

Propositions 4.4.1 and 4.4.2 imply the absolute convergence of the expression defining the modified continued Rademacher sum $Q_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z, s)$ for $z \in \mathbb{H}$ and $\Re(s) > 1$.

Proposition 4.4.3. *Let Γ be a group commensurable with $G(\mathbb{Z})$, let $\mathfrak{p}, \mathfrak{q} \in P_\Gamma$ be cusps of Γ , and let $\Sigma_{\mathfrak{p}}$ and $\Sigma_{\mathfrak{q}}$ be scaling cosets for Γ at $\mathfrak{p}$ and $\mathfrak{q}$, respectively. Let $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$, and let $m \in \mathbb{Z}^+$. Then the modified Rademacher sum*

$$Q_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z, s) = \sum_{\llbracket \chi \in \llbracket \Sigma_{\mathfrak{p}}^{-1} \Sigma_{\mathfrak{q}}} T_{\llbracket \chi}^{\kappa(m)}(z, s) - T_{0, \llbracket \chi}^{\kappa(m)}(s) \quad (4.4.4)$$

converges absolutely for $z \in \mathbb{H}$ and $\Re(s) > 1$.

We next seek to relate the modified Rademacher sums $Q_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z, s)$ to the functions $F_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z, s)$ and $Z_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(s)$ of §4.2 and §4.3, respectively.

Proposition 4.4.4. *Let Γ be a group commensurable with $G(\mathbb{Z})$, let $\mathfrak{p}, \mathfrak{q} \in P_\Gamma$ be cusps of Γ , with scaling matrices $\Sigma_{\mathfrak{p}}$ and $\Sigma_{\mathfrak{q}}$, respectively. Let $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$ and let $m \in \mathbb{Z}^+$.*

Then we have

$$\sum_{\llbracket \chi \in \llbracket \Sigma_p^{-1} \Sigma_q \rrbracket} T_{0, \llbracket \chi \rrbracket}^{\kappa(m)}(s) = -Z_{\Gamma, p|q}^{\kappa(m)}(1-s) \quad (4.4.5)$$

for $z \in \mathbb{H}$ and $\Re(s) > 1$.

Proof. Set $U = \llbracket \Sigma_p^{-1} \Sigma_q \rrbracket$ and $S = \llbracket \Sigma_p^{-1} \Sigma_q \rrbracket^\times$. Recall that $T_{0, \llbracket \chi \rrbracket}^{\kappa(m)}(s) = 0$ when $\llbracket \chi \rrbracket \notin \llbracket G(\mathbb{Q}) \rrbracket^{\times \times}$, and otherwise

$$\begin{aligned} T_{0, \llbracket \chi \rrbracket}^{\kappa(m)}(s) = & e(-m \llbracket \chi \rrbracket \cdot \infty) (2\pi i(m \llbracket \chi \rrbracket \cdot \infty - m \llbracket \chi \rrbracket \cdot 0))^{(s-2\kappa)} \text{jac}(\llbracket \chi \rrbracket, 0)^\kappa \\ & - e(s/2) e(-m \llbracket \chi \rrbracket \cdot \infty) (2\pi i(m \llbracket \chi \rrbracket \cdot 0 - m \llbracket \chi \rrbracket \cdot \infty))^{(s-2\kappa)} \text{jac}(\llbracket \chi \rrbracket, 0)^\kappa, \end{aligned} \quad (4.4.6)$$

by the definition (cf. §3.1) of $T_{0, \llbracket \chi \rrbracket}^{\kappa(m)}(s)$. We apply the identities $\chi \cdot \infty - \chi \cdot 0 = \varrho(\chi)/(-\chi^{-1} \cdot \infty)$ and $\text{jac}(\llbracket \chi \rrbracket, 0) = \varrho(\chi)/(-\chi^{-1} \cdot \infty)^2$ (cf. §2.4) so as to write

$$T_{0, \llbracket \chi \rrbracket}^{\kappa(m)}(s) = e(-m \llbracket \chi \rrbracket \cdot \infty) \varrho \llbracket \chi \rrbracket^{s-\kappa} (2\pi i m)^{(s-2\kappa)} ((-\chi^{-1} \llbracket \chi \rrbracket \cdot \infty)^{-s} - e(s/2)(\chi^{-1} \llbracket \chi \rrbracket \cdot \infty)^{-s}). \quad (4.4.7)$$

Let $\llbracket \chi \rrbracket \in \llbracket G(\mathbb{Q}) \rrbracket^\times$. Then we have

$$\sum_{\llbracket \chi \in \llbracket \chi \rrbracket} T_{0, \llbracket \chi \rrbracket}^{\kappa(m)}(s) = e(-m \llbracket \chi \rrbracket \cdot \infty) \varrho \llbracket \chi \rrbracket^{s-\kappa} (2\pi i m)^{(s-2\kappa)} \sum_{n \in \mathbb{Z}}^* ((n + \alpha_{\llbracket \chi \rrbracket})^{-s} - e(s/2)(n - \alpha_{\llbracket \chi \rrbracket})^{-s}) \quad (4.4.8)$$

where $\alpha_{\llbracket \chi \rrbracket} \in \mathbb{Q}$ is chosen so that $0 \leq \alpha_{\llbracket \chi \rrbracket} < 1$ and $\alpha_{\llbracket \chi \rrbracket} + \mathbb{Z} = -\chi^{-1} \cdot \infty + \mathbb{Z}$, and the superscript in the summation $\sum_{n \in \mathbb{Z}}^*$ indicates to omit the term corresponding to $n = 0$ in case $\alpha_{\llbracket \chi \rrbracket} = 0$. We compute

$$\sum_{n \in \mathbb{Z}}^* ((n + \alpha_{\llbracket \chi \rrbracket})^{-s} - e(s/2)(n - \alpha_{\llbracket \chi \rrbracket})^{-s}) = (e(-s/2) - e(s/2))\zeta(1 - \alpha_{\llbracket \chi \rrbracket}, s) \quad (4.4.9)$$

where $\zeta(\alpha, s)$ denotes the Hurwitz zeta function (cf. §4.1), so we have

$$\sum_{\llbracket \chi \in U^{\times \times} \rrbracket} T_{0, \llbracket \chi \rrbracket}^{\kappa(m)}(s) = \sum_{\llbracket \chi \rrbracket \in S^\times} e(-m \llbracket \chi \rrbracket \cdot \infty) \varrho \llbracket \chi \rrbracket^{s-\kappa} (2\pi i m)^{(s-2\kappa)} (e(-s/2) - e(s/2))\zeta(1 - \alpha_{\llbracket \chi \rrbracket}, s). \quad (4.4.10)$$

Comparing with the expression (4.3.3) defining $Z_{\llbracket \chi \rrbracket}^{\kappa(m)}(s)'$ we see that the right hand side of (4.4.10) coincides with $-Z_{S^\times}^{\kappa(m)}(s)'$. Now Lemma 4.3.1 yields the required identity (4.4.5). This completes the proof. $\square$

Theorem 4.4.5. *Let Γ be a group commensurable with $G(\mathbb{Z})$ and let $\mathfrak{p}, \mathfrak{q} \in \mathcal{P}_\Gamma$ be cusps of Γ . Let $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$, and let $m \in \mathbb{Z}^+$. Then we have*

$$Q_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z, s) = \delta_{\Gamma, \mathfrak{p}|\mathfrak{q}} e(-mz) + Z_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(1 - s) + F_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z, s)_{\text{van}} \quad (4.4.11)$$

for $z \in \mathbb{H}$ and $\Re(s) > 1$.

Proof. Let $\kappa, m \in \mathbb{Z}^+$, let $s \in \mathbb{C}$ with $\Re(s) > 1$, set $U = \llbracket \Sigma_{\mathfrak{p}}^{-1} \Sigma_{\mathfrak{q}} \rrbracket$ and set $S = \llbracket \Sigma_{\mathfrak{p}}^{-1} \Sigma_{\mathfrak{q}} \rrbracket^\times$. By the defining properties of scaling cosets (cf. §2.6) and by the definition (4.1.1) of the continued Rademacher component function $T_{\llbracket \chi \rrbracket}^{1-\kappa(m)}(z, s)$ we have

$$\sum_{\llbracket \chi \rrbracket \in U} T_{\llbracket \chi \rrbracket}^{1-\kappa(m)}(z, s) = \delta_{\Gamma, \mathfrak{p}|\mathfrak{q}} e(-mz) + \sum_{\llbracket \chi \rrbracket \in U^\times} T_{\llbracket \chi \rrbracket}^{1-\kappa(m)}(z, s) \quad (4.4.12)$$

so by Proposition 4.4.4, and the definition of the modified continued Rademacher component function $Q_{\llbracket \chi \rrbracket}^{1-\kappa(m)}(z, s)$, it suffices for us to show that

$$\sum_{\llbracket \chi \rrbracket \in U^\times} T_{\llbracket \chi \rrbracket}^{1-\kappa(m)}(z, s) = F_S^{1-\kappa(m)}(z, s)_{\text{van}}. \quad (4.4.13)$$

We will verify the equality of (4.4.13) by using the Lipschitz summation formula (3.3.3) to transform the expression on the right hand side of (4.4.13) into that on the left. This is essentially the approach employed originally by Rademacher in [Rad39], and the reverse of that employed by Niebur in Lemma 4.2 of [Nie74], except that we can avoid the technical difficulties of [Rad39] and [Nie74], such as the need to employ the identity (3.3.4), by working only with absolutely convergent series.

We begin by inspecting Lemma 4.2.1 to find that

$$\begin{aligned} & (-1)^{1-\kappa} c_S^{1-\kappa}(m, n, s) \\ &= \sum_{\llbracket \chi \rrbracket \in S} \sum_{k \in \mathbb{N}} e(-m \llbracket \chi \rrbracket \cdot \infty) e(-n \llbracket \chi^{-1} \rrbracket \cdot \infty) (4\pi^2 \varrho \llbracket \chi \rrbracket)^{k+\kappa+s-1} m^{(k+2\kappa+s-2)} n^{(k+s-1)}, \end{aligned} \quad (4.4.14)$$

As a shorthand let us set $F_{\text{van}} = F_S^{1-\kappa(m)}(z, s)_{\text{van}}$. Then we have the following expression for F_{van} as a triple sum.

$$\begin{aligned} & (-1)^{1-\kappa} F_{\text{van}} = \\ & \lim_{n \in \mathbb{Z}^+} \sum_{\llbracket \chi \rrbracket \in S} \sum_{k \in \mathbb{N}} e(-m \llbracket \chi \rrbracket \cdot \infty) e(-n \llbracket \chi^{-1} \rrbracket \cdot \infty) (4\pi^2 \varrho \llbracket \chi \rrbracket)^{k+\kappa+s-1} m^{(k+2\kappa+s-2)} n^{(k+s-1)} e(nz) \end{aligned} \quad (4.4.15)$$

We now move the summation over n past the other two summations, simultaneously pulling the terms $e(-m \llbracket \chi \rrbracket \cdot \infty)$ outside the summation over k , and combining the exponents involving n . We thus obtain

$$(-1)^{1-\kappa} F_{\text{van}} = \lim_{\llbracket \chi \rrbracket \in S} e(-m \llbracket \chi \rrbracket \cdot \infty) \sum_{k \in \mathbb{N}} (4\pi^2 \varrho \llbracket \chi \rrbracket)^{k+\kappa+s-1} m^{(k+2\kappa+s-2)} \sum_{n \in \mathbb{Z}^+} n^{(k+s-1)} e(n(z - \llbracket \chi^{-1} \rrbracket \cdot \infty)). \quad (4.4.16)$$

Applying the Lipschitz summation formula (3.3.3) to each summation over n in (4.4.16) we find that

$$(-1)^{1-\kappa} F_{\text{van}} = \lim_{\llbracket \chi \rrbracket \in S} e(-m \llbracket \chi \rrbracket \cdot \infty) \sum_{k \in \mathbb{N}} (4\pi^2 \varrho \llbracket \chi \rrbracket)^{\kappa-1} m^{(k+2\kappa+s-2)} \sum_{n \in \mathbb{Z}} (2\pi i \varrho \llbracket \chi \rrbracket)^{k+s} (z + n - \llbracket \chi^{-1} \rrbracket \cdot \infty)^{-k-s} \quad (4.4.17)$$

where $\llbracket \chi^{-1} \rrbracket$ is any (left) coset of $B(\mathbb{Z})$ in the double coset $\llbracket \chi^{-1} \rrbracket$, for each $\llbracket \chi \rrbracket \in S$. Using the identity (2.4.5) we write F_{van} as $F_{\text{van}} = \sum_{\llbracket \chi \rrbracket \in S} e(-m \llbracket \chi \rrbracket \cdot \infty) F_{\text{van}, \llbracket \chi \rrbracket}$ where

$$F_{\text{van}, \llbracket \chi \rrbracket} = \sum_{k \in \mathbb{N}} \sum_{n \in \mathbb{Z}} (2\pi i \varrho \llbracket \chi \rrbracket m)^{k+2\kappa+s-2} (z + n - \llbracket \chi^{-1} \rrbracket \cdot \infty)^{-k-2\kappa-s+2} \text{jac}(\llbracket \chi \rrbracket, z + n)^{1-\kappa}. \quad (4.4.18)$$

Recalling the definition of the generalized exponential function $e(z, s)$ from (2.1.6) we see that

$$F_{\text{van}} = \sum_{\llbracket \chi \rrbracket \in U^\times} e(-m \llbracket \chi \rrbracket \cdot \infty) e\left(\frac{\varrho \llbracket \chi \rrbracket m}{z - \llbracket \chi^{-1} \rrbracket \cdot \infty}, s - 2(1 - \kappa)\right) \text{jac}(\llbracket \chi \rrbracket, z)^{1-\kappa}. \quad (4.4.19)$$

According to (2.4.4) we have $\varrho \llbracket \chi \rrbracket / (z - \llbracket \chi^{-1} \rrbracket \cdot \infty) = \chi \cdot \infty - \chi \cdot z$. Using this identity together with the Kummer transformation $\Phi(a, b, z) = e(z)\Phi(b - a, b, -z)$ (cf. (2.1.3)) we readily find that

$$e(-m \llbracket \chi \rrbracket \cdot \infty) e\left(\frac{\varrho \llbracket \chi \rrbracket m}{z - \llbracket \chi^{-1} \rrbracket \cdot \infty}, s - 2(1 - \kappa)\right) = e(-m \llbracket \chi \rrbracket \cdot z) t^{1-\kappa}(m, \llbracket \chi \rrbracket, z, s), \quad (4.4.20)$$

so that we have

$$F_{\text{van}} = \sum_{\llbracket \chi \rrbracket \in U^\times} T_{\llbracket \chi \rrbracket}^{1-\kappa(m)}(z, s). \quad (4.4.21)$$

This establishes the required identity (4.4.13). $\square$

In the course of proving Theorem 4.4.5 we have also established the Fourier expansion of the continued Rademacher sums $T_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z, s)$. We record the result as follows.

Theorem 4.4.6. *Let Γ be a group commensurable with $G(\mathbb{Z})$ and let $\mathfrak{p}, \mathfrak{q} \in \mathcal{P}_\Gamma$ be cusps of Γ . Let $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$ and let $m \in \mathbb{Z}^+$. Then we have*

$$T_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z, s) = \delta_{\Gamma, \mathfrak{p}|\mathfrak{q}} e(-mz) + F_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z, s)_{\text{van}} \quad (4.4.22)$$

for $z \in \mathbb{H}$ and $\Re(s) > 1$.

Theorem 4.4.5 will facilitate the identification of the Fourier expansion of the modified Rademacher sum $Q_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z)$ obtained by sending s to 1 in $Q_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z, s)$. Indeed, we will see presently (cf. Proposition 4.4.8) that the series $\tilde{F}_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(q)_{\text{reg}}$ encodes the regular part of the fourier series expansion of $Q_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z)$ precisely, in the case that $1 - \kappa \in \mathbb{Z}^+$. In preparation for this we state the following result which identifies the value of the zeta function $Z_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(s)$ at $s = 0$.

Proposition 4.4.7. *Let $S \subset [G(\mathbb{Q})]^\times$, let $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$, let $m \in \mathbb{Z}^+$. Then we have*

$$Z_S^{\kappa(m)}(0) = c_S^\kappa(m, 0). \quad (4.4.23)$$

Proof. According to the proof of Lemma 4.3.1 the function $\Gamma(s)Z_{[\chi]}^{\kappa(m)}(s)$ coincides with

$$(-1)^{1-\kappa} e(-m [\chi] \cdot \infty) (4\pi^2 \varrho [\chi])^{1-s-\kappa} m^{(1-s-2\kappa)} (F(\alpha_{[\chi]}, s) + e(-s/2) F(-\alpha_{[\chi]}, s)) \quad (4.4.24)$$

where $0 \leq \alpha_{[\chi]} < 1$ and $\alpha_{[\chi]} + \mathbb{Z} = -\chi^{-1} \cdot \infty + \mathbb{Z}$. We have $F(\alpha, 0) + F(-\alpha, 0) = -1$ for $\alpha \in \mathbb{R}$ (cf. [KR01]), so taking $s = 0$ in (4.4.24) we obtain

$$Z_{[\chi]}^{\kappa(m)}(0) = (-1)^\kappa e(-m [\chi] \cdot \infty) (4\pi^2 \varrho [\chi])^{1-\kappa} m^{(1-2\kappa)}. \quad (4.4.25)$$

Upon inspection of (3.2.13) we see that

$$c_S^\kappa(m, 0) = (-1)^\kappa \sum_{[\chi] \in S} e(-m [\chi] \cdot \infty) (4\pi^2 \varrho [\chi])^{1-\kappa} m^{(1-2\kappa)} \quad (4.4.26)$$

for $1 - \kappa \in \mathbb{Z}^+$, and the claim now follows since $Z_S^{\kappa(m)}(s) = \sum_{[\chi] \in S} Z_{[\chi]}^{\kappa(m)}(s)$ by definition. $\square$

Proposition 4.4.8. *Let Γ be a group commensurable with $G(\mathbb{Z})$, and let $\mathfrak{p}, \mathfrak{q} \in P_\Gamma$ be cusps of Γ . Let $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$ and let $m \in \mathbb{Z}^+$. Then we have*

$$Q_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z) = \delta_{\Gamma, \mathfrak{p}|\mathfrak{q}} e(-mz) + F_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z)_{\text{reg}} \quad (4.4.27)$$

for $z \in \mathbb{H}$.

Proof. We take the limit as s tends to 1 in the right hand side of the identity (4.4.11). We have $F_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z, 1)_{\text{van}} = F_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z)_{\text{van}}$ by definition (cf. §4.2), and we have $Z_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(0) = c_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa}(m, 0)$ by Proposition 4.4.7. The required identity (4.4.27) now follows from the fact that $F_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z)_{\text{reg}} = c_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa}(m, 0) + F_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z)_{\text{van}}$, also by definition (cf. §3.2). $\square$

The proof of Proposition 4.4.8 implies a result analogous to (4.4.27) for the normalized Rademacher sum $T_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z)$.

Proposition 4.4.9. *Let Γ be a group commensurable with $G(\mathbb{Z})$ and let $\mathfrak{p}, \mathfrak{q} \in P_\Gamma$ be cusps of Γ . Let $m \in \mathbb{Z}^+$. Then we have*

$$T_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z) = \delta_{\Gamma, \mathfrak{p}|\mathfrak{q}} e(-mz) + F_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z)_{\text{van}} \quad (4.4.28)$$

for $z \in \mathbb{H}$.

We may compare the functions $Q_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z)$ and $R_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z)$ defined by the modified and classical Rademacher sums, respectively. Inspecting Theorem 3.3.2 and Proposition 4.4.8 we obtain the precise relationship, which we record in the following proposition.

Proposition 4.4.10. *Let Γ be a group commensurable with $G(\mathbb{Z})$ and let $\mathfrak{p}, \mathfrak{q} \in P_\Gamma$ be cusps of Γ . Let $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$, and let $m \in \mathbb{Z}^+$. Then we have*

$$Q_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z) = R_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z) + \frac{1}{2} c_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa}(m, 0). \quad (4.4.29)$$

According to Propositions 4.4.8 and 4.4.9 the modified Rademacher sum $Q_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z)$ and the normalized Rademacher sum $T_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z)$ also differ only by a constant function.

Proposition 4.4.11. *Let Γ be a group commensurable with $G(\mathbb{Z})$ and let $\mathfrak{p}, \mathfrak{q} \in P_\Gamma$ be cusps of Γ . Let $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$, and let $m \in \mathbb{Z}^+$. Then we have*

$$Q_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z) = T_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z) + c_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa}(m, 0). \quad (4.4.30)$$

4.5 Variance

In this section we examine how the modified Rademacher sum $Q_{\Gamma, \mathfrak{p}}^{\kappa(m)}(z)$ transforms under the weight 2κ action of Γ . Combining Theorem 3.4.4 and Proposition 4.4.10 we obtain the result

that the modified Rademacher sum $Q_{\Gamma, \mathfrak{p}}^{\kappa(m)}(z)$ is an automorphic integral of weight 2κ for Γ in case Γ has width one at infinity.

Theorem 4.5.1. *Let Γ be a group commensurable with $G(\mathbb{Z})$ that has width one at infinity and let $\mathfrak{p} \in P_\Gamma$ be a cusp of Γ . Let $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$, and let $m \in \mathbb{Z}^+$. Then the modified Rademacher sum $Q_{\Gamma, \mathfrak{p}}^{\kappa(m)}(z)$ is an automorphic integral of weight 2κ for Γ .*

The proof of Theorem 3.4.4 yields the following explicit description of the associated cusp form map $I_\kappa(\Gamma) \rightarrow S_{1-\kappa}(\Gamma)$ on the subspace of $I_\kappa(\Gamma)$ spanned by the modified Rademacher sums $Q_{\Gamma, \mathfrak{p}}^{\kappa(m)}(z)$.

Proposition 4.5.2. *Let Γ be a group commensurable with $G(\mathbb{Z})$ that has width one at infinity and let $\mathfrak{p} \in P_\Gamma$ be a cusp of Γ . Let $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$, and let $m \in \mathbb{Z}^+$. Then*

$$m^\kappa Q_{\Gamma, \mathfrak{p}}^{\kappa(m)} \mapsto m^{1-\kappa} P_{\Gamma, \mathfrak{p}}^{1-\kappa(-m)} \mapsto 0, \quad (4.5.1)$$

under the maps $I_\kappa(\Gamma) \mapsto S_{1-\kappa}(\Gamma)$ of §2.7.

We next seek to determine the utility of Proposition 4.5.2 by identifying the subspace of $I_\kappa(\Gamma)$ that is spanned by the modified Rademacher sums $Q_{\Gamma, \mathfrak{p}}^{\kappa(m)}(z)$.

Theorem 4.5.3. *Let Γ be a group commensurable with $G(\mathbb{Z})$ that has width one at infinity, and let $\kappa \in \mathbb{Z}$ with $1 - \kappa \in \mathbb{Z}^+$. Then the set $\{Q_{\Gamma, \mathfrak{p}}^{\kappa(m)}(z) \mid \mathfrak{p} \in P_\Gamma, m \in \mathbb{Z}^+\}$ is a basis for the space of automorphic integrals of weight 2κ for Γ in case $\kappa < 0$. When $\kappa = 0$ the set $\{Q_{\Gamma, \mathfrak{p}}^{(m)}(z) \mid \mathfrak{p} \in P_\Gamma, m \in \mathbb{Z}^+\}$ spans a subspace of $I_0(\Gamma)$ of codimension 1.*

Proof. The function $Q_{\Gamma, \mathfrak{p}}^{\kappa(m)}(z)$ has principal part $e(-mz)$ at the cusp $\mathfrak{p}$, and vanishes at all the other cusps, so the collection $\{Q_{\Gamma, \mathfrak{p}}^{\kappa(m)}(z) \mid \mathfrak{p} \in P_\Gamma, m \in \mathbb{Z}^+\}$ is linearly independent. To show that it furnishes a basis we follow Niebur's proof of Theorem 3.3 in [Nie74], employing Petersson's generalized Riemann–Roch Theorem to show that the dimension of the subspace of $I_\kappa(\Gamma)$ containing automorphic integrals with poles degree not more than m say at $\mathfrak{p}$ is bounded above by m less the dimension of the space $S_{1-\kappa}(\Gamma)$, for m sufficiently large. $\square$

We write $I'_0(\Gamma)$ for the subspace of $I_0(\Gamma)$ spanned by the modified Rademacher sums $Q_{\Gamma, \mathfrak{p}}^{(m)}(z)$ of weight 0 for varying $\mathfrak{p} \in P_\Gamma$ and $m \in \mathbb{Z}^+$.

Theorem 4.5.4. *Let Γ be a group commensurable with $G(\mathbb{Z})$ that has width one at infinity. Then for any $\kappa \in \mathbb{Z}$ the sequence*

$$0 \rightarrow M_\kappa(\Gamma) \rightarrow I_\kappa(\Gamma) \rightarrow S_{1-\kappa}(\Gamma) \rightarrow 0 \quad (4.5.2)$$

is exact.

Proof. The second map of (4.5.2) is an inclusion by definition, and we observed already in §2.7 that $M_\kappa(\Gamma)$ is the kernel of the associated cusp form map $I_\kappa(\Gamma) \rightarrow S_{1-\kappa}(\Gamma)$ (cf. Lemma 2.7.1), so we require to show only that the second to last map of (4.5.2) is surjective. For this we recall the fact (cf. [Iwa97]) that the holomorphic Poincaré series $P_{\Gamma, \mathfrak{p}}^{1-\kappa(-m)}(z)$, with varying $m \in \mathbb{Z}^+$, span the space $S_{1-\kappa}(\Gamma)$ of cusp forms of weight $2 - 2\kappa$ for Γ whenever $1 - \kappa \in \mathbb{Z}^+$. Indeed, one need not vary the cusp $\mathfrak{p}$ in order to obtain a spanning set. Given an arbitrary cusp form $g \in S_{1-\kappa}(\Gamma)$ we may then write

$$g = \sum_{n \in \mathbb{Z}^+} a^{-n} P_\Gamma^{1-\kappa(-n)} \quad (4.5.3)$$

for some $a^{-n} \in \mathbb{C}$, with only finitely many a^{-n} non-zero. Employing Proposition 4.5.2 we see that the automorphic integral

$$f = \sum_{n \in \mathbb{Z}^+} n^{2\kappa-1} a^{-n} Q_\Gamma^{\kappa(n)} \quad (4.5.4)$$

is mapped to g by the associated cusp form map. We conclude that the associated cusp form map is surjective. This completes the proof. $\square$

Theorem 4.5.3 and Proposition 4.5.2 together describe the associated cusp form map $I_\kappa(\Gamma) \rightarrow S_{1-\kappa}(\Gamma)$ explicitly, and thus we obtain a powerful criterion for determining when an automorphic integral $f \in I_\kappa(\Gamma)$ lies in the subspace $M_\kappa(\Gamma)$ of modular forms. For example, we have the following corollary, indicating exactly when the modified Rademacher sum $Q_{\Gamma, \mathfrak{p}}^{\kappa(m)}(z)$ is a modular form for Γ in the sense of §2.7.

Corollary 4.5.5. *Let Γ be a group commensurable with $G(\mathbb{Z})$ that has width one at infinity and let $\mathfrak{p} \in P_\Gamma$ be a cusp of Γ . Let $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$, and let $m \in \mathbb{Z}^+$. Then the modified Rademacher sum $Q_{\Gamma, \mathfrak{p}}^{\kappa(m)}(z)$ is a modular form of weight 2κ for Γ if and only if the cusp form $P_{\Gamma, \mathfrak{p}}^{1-\kappa(-m)}(z)$ vanishes identically.*

5 Structural applications

We have seen already in §4 various applications of the modified Rademacher sums, and the continuation procedure introduced in §4.1, such as the basis theorem (Theorem 4.5.3) for automorphic integrals, and the explicit description (cf. Proposition 4.5.2) of the associated cusp form map $I_\kappa(\Gamma) \rightarrow S_{1-\kappa}(\Gamma)$. In this section we explore further consequences of the modified Rademacher sum construction for the structure of the spaces of automorphic integrals associated to groups commensurable with the modular group. Applications of the modified Rademacher sums to monstrous moonshine and quantum gravity will be developed in §§6,7.

5.1 Constants

Perhaps the most striking application of the continuation procedure of §4.1 is the correction of the constant terms appearing in the classical Rademacher sums. For negative weights, the constant term in the Fourier expansion of an automorphic integral $f \in I_\kappa(\Gamma)$ is determined by the automorphy condition (2.7.13). At weight 0 constant functions are themselves automorphic integrals, so the classical Rademacher sums $R_{\Gamma, \mathfrak{p}}^{(m)}(z)$ of weight 0 do not fail to be automorphic, and it is natural then to wonder about the significance of the constant term $c_{\Gamma, \mathfrak{p}}(m, 0)$ appearing in the Fourier expansion of the modified Rademacher sum $Q_{\Gamma, \mathfrak{p}}^{(m)}(z)$ of weight 0 associated to a group Γ at a cusp $\mathfrak{p} \in \mathcal{P}_\Gamma$.

Let $\mathfrak{G}$ denote the set of subgroups of $G(\mathbb{Q})$ that are commensurable with $G(\mathbb{Z})$. Recall from §2.7 that for $\Gamma \in \mathfrak{G}$ we write $M_0(\Gamma)$ for the space of holomorphic functions on $\mathbb{H}$ that are invariant for the natural action of Γ and are meromorphic at the cusps of Γ . Let $\mathfrak{M}_0$ be the union of the spaces $M_0(\Gamma)$ for $\Gamma \in \mathfrak{G}$.

$$\mathfrak{M}_0 = \bigcup_{\Gamma \in \mathfrak{G}} M_0(\Gamma) \quad (5.1.1)$$

Since $\mathfrak{G}$ is closed under intersections, the set $\mathfrak{M}_0$ is in fact a subalgebra of the space $\mathcal{O}(\mathbb{H})$ of holomorphic functions on $\mathbb{H}$. Since $G(\mathbb{Q})$ is the commensurator of $G(\mathbb{Z})$, the algebra $\mathfrak{M}_0$ is $G(\mathbb{Q})$ -invariant. The constant functions evidently constitute a $G(\mathbb{Q})$ -invariant subspace $\mathbb{C}1 \subset \mathfrak{M}_0$. We may enquire as to the $G(\mathbb{Q})$ -module structure of the quotient $\mathfrak{M}_0/\mathbb{C}1$. According to [Nor84] this quotient is irreducible, and the natural map $\mathfrak{M}_0 \rightarrow \mathfrak{M}_0/\mathbb{C}1$ admits a section.

Theorem 5.1.1 ([Nor84]). *There exists a unique $G(\mathbb{Q})$ -submodule $\mathfrak{M}'_0 \subset \mathfrak{M}_0$ with the property that the natural inclusions $\mathbb{C}1, \mathfrak{M}'_0 \subset \mathfrak{M}_0$ induce an isomorphism*

$$\mathfrak{M}_0 \cong \mathbb{C}1 \oplus \mathfrak{M}'_0 \quad (5.1.2)$$

of $G(\mathbb{Q})$ -modules.

According to Theorem 5.1.1, for each $f \in \mathfrak{M}_0$ there is a unique $c(f) \in \mathbb{C}$ with the property that $f + c(f) \in \mathfrak{M}'_0$. The value $c(f)$ is called the *Rademacher constant of f* . The assignment $f \mapsto c(f)$ defines a linear function on $\mathfrak{M}_0$ with kernel $\mathfrak{M}'_0$ which we call the *Rademacher constant function*. We suggest that the Rademacher constant function may be generalized to automorphic integrals of weight 0 in the following way. Let $\mathfrak{I}_0$ be the union of the spaces $I_0(\Gamma)$ for $\Gamma \in \mathfrak{G}$.

$$\mathfrak{I}_0 = \bigcup_{\Gamma \in \mathfrak{G}} I_0(\Gamma) \quad (5.1.3)$$

Since $\mathfrak{G}$ is closed under intersections, the set $\mathfrak{I}_0$ is also a subalgebra of $\mathcal{O}(\mathbb{H})$, and since $G(\mathbb{Q})$

is the commensurator of $G(\mathbb{Z})$, the algebra $\mathfrak{I}_0$ is also stable under the action of $G(\mathbb{Q})$. The constant functions again constitute a $G(\mathbb{Q})$ -invariant subspace $\mathbb{C}1 \subset \mathfrak{I}_0$. We conjecture that the natural analogue of Theorem 5.1.1 holds for $\mathfrak{I}_0$. Precisely, we conjecture that there is a unique $G(\mathbb{Q})$ -submodule $\mathfrak{I}'_0 \subset \mathfrak{I}_0$ with the property that the natural inclusions induce an isomorphism

$$\mathfrak{I}_0 \cong \mathbb{C}1 \oplus \mathfrak{I}'_0 \quad (5.1.4)$$

of $G(\mathbb{Q})$ -modules. Further, we conjecture that $\mathfrak{I}'_0$ is just the space spanned by the modified Rademacher sums $Q_{\Gamma, \mathfrak{p}}^{\kappa(m)}(z)$ for $\Gamma \in \mathfrak{G}$ and $\mathfrak{p} \in P_\Gamma$ and $m \in \mathbb{Z}^+$.

5.2 Inner products

Let Γ be a group commensurable with $G(\mathbb{Z})$ and let $\kappa \in \mathbb{Z}^+$. Then the space $S_\kappa(\Gamma)$ of cusp forms of weight 2κ for Γ becomes a Hilbert space when equipped with the *Petersson inner product*, defined by setting

$$\langle f, g \rangle = \int_{\mathfrak{F}_\Gamma} f(z) \overline{g(z)} \mathfrak{I}(z)^{2\kappa} d\mu(z) \quad (5.2.1)$$

for $f, g \in S_\kappa(\Gamma)$, where $\mathfrak{F}_\Gamma$ is a fundamental domain for Γ . Let us define a *normalized Petersson inner product of weight 2κ for Γ* , to be denoted $\langle \cdot, \cdot \rangle_\Gamma^\kappa$, by setting

$$\langle f, g \rangle_\Gamma^\kappa = (4\pi)^{(2\kappa-2)} \int_{\mathfrak{F}_\Gamma} f(z) \overline{g(z)} \mathfrak{I}(z)^{2\kappa} d\mu(z) \quad (5.2.2)$$

for $f, g \in S_\kappa(\Gamma)$. It is well known (and follows from Theorem 3.4.3), that for $\kappa, -m \in \mathbb{Z}^+$ and $\mathfrak{p}$ a cusp for Γ , the holomorphic Poincaré series $P_{\Gamma, \mathfrak{p}}^{\kappa(m)}(z)$ is a cusp form of weight 2κ for Γ in case Γ has width one at infinity. A standard calculation (cf. [Iwa97]) shows that we have

$$\left\langle P_{\Gamma, \mathfrak{p}}^{\kappa(m)}, P_{\Gamma, \mathfrak{q}}^{\kappa(n)} \right\rangle_\Gamma^\kappa = (-n)^{2-2\kappa} (\delta_{\Gamma, \mathfrak{p}|\mathfrak{q}} \delta_{m,n} + c_{\Gamma, \mathfrak{p}|\mathfrak{q}}^\kappa(m, -n)) \quad (5.2.3)$$

for $\mathfrak{p}, \mathfrak{q} \in P_\Gamma$ and $-m, -n \in \mathbb{Z}^+$. According to Theorem 3.4.3 the right hand side of (5.2.3) is, up to the scalar factor $n^{2-2\kappa}$, the coefficient of $e(-nz)$ in the expansion $P_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z)$ of the Poincaré series $P_{\Gamma, \mathfrak{p}}^{\kappa(m)}(z)$ at $\mathfrak{q}$. Now the modified Rademacher sums $Q_{\Gamma, \mathfrak{p}}^{\kappa(m)}(z)$, for $-\kappa, m \in \mathbb{Z}^+$, span the space $I_\kappa(\Gamma)$ of automorphic integrals for weight 2κ for Γ , according to Theorem 4.5.3, and thus serve an analogous rôle for the spaces $I_\kappa(\Gamma)$ as the Poincaré series $P_{\Gamma, \mathfrak{p}}^{\kappa(m)}(z)$ do for the spaces $S_\kappa(\Gamma)$. It is natural then to use the right hand side of (5.2.3) to extend the normalized Petersson inner product to negative weights. In light of the fact, established in Proposition 4.4.8, that the coefficient of $e(-nz)$ in the expansion $Q_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m)}(z)$ of the modified Rademacher sum $Q_{\Gamma, \mathfrak{p}}^{\kappa(m)}(z)$ at $\mathfrak{q}$

is $\delta_{\Gamma, \mathbf{p}|\mathbf{q}} \delta_{m,n}$ for $m, n \in \mathbb{Z}^+$, we define an inner product $\langle \cdot, \cdot \rangle_{\Gamma}^{\kappa}$ on $I_{\kappa}(\Gamma)$ by setting

$$\left\langle Q_{\Gamma, \mathbf{p}}^{\kappa(m)}, Q_{\Gamma, \mathbf{q}}^{\kappa(n)} \right\rangle_{\Gamma}^{\kappa} = n^{2-2\kappa} \delta_{\Gamma, \mathbf{p}|\mathbf{q}} \delta_{m,n} \quad (5.2.4)$$

for $\mathbf{p}, \mathbf{q} \in \mathbf{P}_{\Gamma}$ and $m, n \in \mathbb{Z}^+$, when $-\kappa \in \mathbb{Z}^+$. In case $\kappa = 0$ the modified Rademacher sums $Q_{\Gamma, \mathbf{p}}^{\kappa(m)}(z)$ span a subspace $I'_0(\Gamma)$ of $I_0(\Gamma)$ of codimension 1 (cf. Theorem 4.5.3 and §5.1). We may use the formula (5.2.4) to define an inner product on this codimension 1 subspace $I'_0(\Gamma)$.

$$\left\langle Q_{\Gamma, \mathbf{p}}^{(m)}, Q_{\Gamma, \mathbf{q}}^{(n)} \right\rangle_{\Gamma}^0 = n^2 \delta_{\Gamma, \mathbf{p}|\mathbf{q}} \delta_{m,n} \quad (5.2.5)$$

A complement to the subspace $I'_0(\Gamma)$ of $I_0(\Gamma)$ is spanned by the constant functions. We extend the inner product $\langle \cdot, \cdot \rangle_{\Gamma}^0$ to all of $I_0(\Gamma)$ by adopting the convention that

$$\left\langle Q_{\Gamma, \mathbf{p}}^{(m)}, f \right\rangle_{\Gamma}^0 = 0 \quad (5.2.6)$$

for all $\mathbf{p} \in \mathbf{P}_{\Gamma}$ and $m \in \mathbb{Z}^+$ whenever f is identically constant.

5.3 Branching

Let Δ and Γ be groups commensurable with $G(\mathbb{Z})$ and suppose that Δ is a subgroup of Γ . Then an automorphic integral for Γ is also an automorphic integral for Δ . By Theorem 4.5.3 the modified Rademacher sums span the spaces of automorphic integrals, so it is natural to consider the problem of expressing the modified Rademacher sums $Q_{\Gamma, \mathbf{p}}^{\kappa(m)}$ associated to Γ in terms of the modified Rademacher sums $Q_{\Delta, \mathbf{r}}^{\kappa(n)}$ associated to Δ .

Suppose then that Δ and Γ are groups commensurable with $G(\mathbb{Z})$ and Δ is a subgroup of Γ . Let $\mathbf{p}$ be a cusp for Γ , let $\Sigma_{\mathbf{p}}$ be a scaling coset for Γ at $\mathbf{p}$, and choose a system $\{\Sigma_r \mid r \in \mathbf{P}_{\Delta}\}$ of scaling cosets for Δ . Suppose for now that κ is a negative integer. Then by Theorem 4.5.3 we have

$$Q_{\Gamma, \mathbf{p}}^{\kappa(m)}(z) = \sum_{r \in \mathbf{P}_{\Delta}} b_r^n Q_{\Delta, r}^{\kappa(n)}(z) \quad (5.3.1)$$

for some $b_r^n \in \mathbb{C}$, and these b_r^n are non-zero for only finitely many $n \in \mathbb{Z}^+$. Furthermore, since the modified Rademacher sum $Q_{\Gamma, \mathbf{p}}^{\kappa(m)}(z)$ has no poles away from $\mathbf{p}$, the coefficient b_r^n can be non-zero only when r lies in the preimage of $\mathbf{p}$ under the natural map $\mathbf{X}_{\Delta} \rightarrow \mathbf{X}_{\Gamma}$, which is to say, b_r^n is non-zero only when $r \in \Delta \backslash \mathbf{p}$. It is natural then to consider the map

$$\begin{aligned} \Delta \backslash \Sigma_{\mathbf{p}} &\rightarrow \Delta \backslash \mathbf{p} \\ \Delta \sigma &\mapsto \Delta \sigma \cdot \infty \end{aligned} \quad (5.3.2)$$

sending right cosets of Δ in Σ_p to cusps of Δ contained in $\mathfrak{p} \subset \hat{\mathbb{Q}}$. The map (5.3.2) is always surjective. We may ask under what circumstances it is also injective.

Lemma 5.3.1. *Let Δ and Γ be groups commensurable with $G(\mathbb{Z})$ and suppose that Δ is a subgroup of Γ . Let $\mathfrak{p}$ be a cusp for Γ and let Σ_p be a scaling coset for Γ at $\mathfrak{p}$. Then the map $\Delta \backslash \Sigma_p \rightarrow \Delta \backslash \mathfrak{p}$ of (5.3.2) is injective if and only if $\Delta\sigma$ is a scaling coset for Δ at $\Delta\sigma \cdot \infty$ for every $\Delta\sigma \in \Delta \backslash \Sigma_p$.*

Proof. Suppose the map $\Delta \backslash \Sigma_p \rightarrow \Delta \backslash \mathfrak{p}$ of (5.3.2) is injective, and let $\sigma \in \Sigma_p$. The forward implication of the lemma follows if we can show that $\Delta\sigma$ is a scaling coset for Δ at $\Delta\sigma \cdot \infty$. For this it suffices to show that $(\sigma^{-1}\Delta\sigma)_\infty = B(\mathbb{Z})$. Since $\sigma^{-1}\Delta\sigma$ is a subgroup of $\sigma^{-1}\Gamma\sigma$, and $(\sigma^{-1}\Gamma\sigma)_\infty = (\Sigma_p^{-1}\Sigma_p)_\infty = B(\mathbb{Z})$ by the defining properties of Σ_p , we have the inclusion $(\sigma^{-1}\Delta\sigma)_\infty \subset B(\mathbb{Z})$, so it suffices to show that the translation T belongs to $\sigma^{-1}\Delta\sigma$. Now $\Delta\sigma T$ is also a coset of Δ in Σ_p , since $\Gamma\sigma = \Sigma_p$ is a union of left cosets of $B(\mathbb{Z})$. Since T fixes ∞ we have $\Delta\sigma T \cdot \infty = \Delta\sigma \cdot \infty$, and thus the cosets $\Delta\sigma T$ and $\Delta\sigma$ coincide by the assumed injectivity of the map (5.3.2). It follows that $T \in \sigma^{-1}\Delta\sigma$, so that $\Delta\sigma$ is indeed a scaling coset for Δ .

For the reverse implication suppose that $\Delta\sigma$ is a scaling coset for Δ for every $\sigma \in \Sigma_p$. Let $\sigma, \sigma' \in \Sigma_p$ and suppose that $\Delta\sigma \cdot \infty = \Delta\sigma' \cdot \infty$. Then $\sigma^{-1}\delta\sigma' \in (\Sigma_p^{-1}\Sigma_p)_\infty = B(\mathbb{Z})$ for some $\delta \in \Delta$, so $\sigma^{-1}\delta\sigma' = T^n$ for some $n \in \mathbb{Z}$, and this implies $\Delta\sigma' = \Delta\sigma T^n$. Now the assumption that $\Delta\sigma$ is a scaling coset for Δ implies that $\Delta\sigma$ is a union of left cosets of $B(\mathbb{Z})$, so $\Delta\sigma T^n = \Delta\sigma$. We conclude that $\Delta\sigma' = \Delta\sigma$, so that the map $\Delta \backslash \Sigma_p \rightarrow \Delta \backslash \mathfrak{p}$ is indeed injective. This completes the proof of the lemma. $\square$

According to the discussion of §2.6 a coset $\Delta\sigma \in \Delta \backslash \Sigma_p$ becomes a scaling coset for Δ at $\Delta\sigma \cdot \infty$ only once we multiply it on the right by $[\beta]$ (cf. (2.3.3)) for some $\beta \in \mathbb{Q}^+$. In fact, this β is a positive integer, for we have $(\sigma^{-1}\Delta\sigma)_\infty \subset (\Sigma_p^{-1}\Sigma_p)_\infty = B(\mathbb{Z})$, so that $(\sigma^{-1}\Delta\sigma)_\infty$ is generated by T^n for some $n \in \mathbb{Z}^+$. In order that $\Delta\sigma[\beta]$ be a scaling coset for Δ at $\Delta\sigma \cdot \infty$ we should have $([1/\beta]\sigma^{-1}\Delta\sigma[\beta])_\infty = B(\mathbb{Z})$, so the computation $[1/\beta]T^n[\beta] = T^{n/\beta}$ shows that $\beta = n$. Evidently we may define a function $\Delta \backslash \Sigma_p \rightarrow \mathbb{Z}^+$ by mapping the coset $\Delta\sigma$ to the positive integer n such that T^n generates $(\sigma^{-1}\Delta\sigma)_\infty$. We next show that this function $\Delta \backslash \Sigma_p \rightarrow \mathbb{Z}^+$ factors through the map $\Delta \backslash \Sigma_p \rightarrow \Delta \backslash \mathfrak{p}$ of (5.3.2).

Lemma 5.3.2. *Let Δ and Γ be groups commensurable with $G(\mathbb{Z})$, and suppose that Δ is a subgroup of Γ . Let $\mathfrak{p} \in \mathcal{P}_\Gamma$ be a cusp of Γ and let Σ_p be a scaling coset for Γ at $\mathfrak{p}$. Then for any $\sigma_1, \sigma_2 \in \Sigma_p$ the identity $\Delta\sigma_1 \cdot \infty = \Delta\sigma_2 \cdot \infty$ implies $(\sigma_1^{-1}\Delta\sigma_1)_\infty = (\sigma_2^{-1}\Delta\sigma_2)_\infty$.*

Proof. Set $\mathfrak{r} = \Delta\sigma_1 \cdot \infty = \Delta\sigma_2 \cdot \infty$. Since $(\sigma^{-1}\Gamma\sigma)_\infty = (\Sigma_p^{-1}\Sigma_p)_\infty = B(\mathbb{Z})$ for any $\sigma \in \Sigma_p$ the groups $(\sigma_1^{-1}\Delta\sigma_1)_\infty$ and $(\sigma_2^{-1}\Delta\sigma_2)_\infty$ are both contained in $B(\mathbb{Z})$. Let n_1 and n_2 be the positive integers such that $(\sigma_1^{-1}\Delta\sigma_1)_\infty$ is generated by T^{n_1} and $(\sigma_2^{-1}\Delta\sigma_2)_\infty$ is generated by T^{n_2} . Then $\Delta\sigma_1[n_1]$ and $\Delta\sigma_2[n_2]$ are both scaling cosets for Δ at $\mathfrak{r}$, so $\Delta\sigma_1[n_1] = \Delta\sigma_2[n_2]T^\alpha$

for some $\alpha \in \mathbb{Q}$. This implies that $[n_2]T^\alpha[1/n_1]$ is an element of $(\Sigma_{\mathbf{p}}^{-1}\Sigma_{\mathbf{p}})_{\infty} = B(\mathbb{Z})$, so that $[n_2]T^\alpha[1/n_1] = T^k$ for some $k \in \mathbb{Z}$. Now we compute

$$[n_2]T^\alpha[1/n_1] = \begin{bmatrix} n_2 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & \alpha \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & n_1 \end{bmatrix} = \begin{bmatrix} n_2 & n_2\alpha n_1 \\ 0 & n_1 \end{bmatrix} = \begin{bmatrix} 1 & k \\ 0 & 1 \end{bmatrix} \quad (5.3.3)$$

and conclude that $n_1 = n_2$, as we required to show. $\square$

On the basis of Lemma 5.3.2 we may define a function $\Delta \backslash \mathbf{p} \rightarrow \mathbb{Z}^+$ for each cusp $\mathbf{p}$ of Γ by first choosing a scaling coset $\Sigma_{\mathbf{p}}$ for Γ at $\mathbf{p}$, and then sending the cusp $\mathbf{r} \in \Delta \backslash \mathbf{p}$ of Δ to the unique positive integer n satisfying $(\sigma^{-1}\Delta\sigma)_{\infty} = \langle T^n \rangle$ when $\mathbf{r} = \Delta\sigma \cdot \infty$. Observe now that this map does not depend upon the choice of scaling coset $\Sigma_{\mathbf{p}}$, for if we replace $\sigma \in \Sigma_{\mathbf{p}}$ with $\sigma' = \sigma T^\alpha \in \Sigma_{\mathbf{p}} T^\alpha$ for some $\alpha \in \mathbb{Q}$, then we have

$$((\sigma')^{-1}\Delta\sigma')_{\infty} = (T^{-\alpha}\sigma^{-1}\Delta\sigma T^\alpha)_{\infty} = T^{-\alpha}(\sigma^{-1}\Delta\sigma)_{\infty} T^\alpha = (\sigma^{-1}\Delta\sigma)_{\infty}. \quad (5.3.4)$$

In light of this we may define a map $w_{\Gamma} : \mathbf{P}_{\Delta} \rightarrow \mathbb{Z}^+$, which we call the *width function associated to Γ* , by setting $w_{\Gamma}(\mathbf{r}) = n$ in case $\mathbf{r} = \Delta\sigma \cdot \infty$ and T^n generates $(\sigma^{-1}\Delta\sigma)_{\infty}$ and $\Gamma\sigma$ is a scaling coset for Γ at the unique cusp of Γ containing $\mathbf{r}$.

$$\begin{aligned} w_{\Gamma} : \mathbf{P}_{\Delta} &\rightarrow \mathbb{Z}^+ \\ \mathbf{r} = \Delta\sigma \cdot \infty, \Gamma\sigma \in \mathfrak{S}_{\Gamma\mathbf{r}} &\implies (\sigma^{-1}\Delta\sigma)_{\infty} = \langle T^{w_{\Gamma}(\mathbf{r})} \rangle \end{aligned} \quad (5.3.5)$$

We call $w_{\Gamma}(\mathbf{r})$ the *width of $\mathbf{r}$ with respect to Γ* . The next result gives upper bounds for the width functions.

Lemma 5.3.3. *Let Δ and Γ be groups commensurable with $G(\mathbb{Z})$, and suppose that Δ is a subgroup of Γ . Then we have*

$$\sum_{\mathbf{r} \in \Delta \backslash \mathbf{p}} w_{\Gamma}(\mathbf{r}) = \# \Delta \backslash \Gamma \quad (5.3.6)$$

for every cusp $\mathbf{p}$ of Γ .

Proof. Let $\mathbf{p}$ be a cusp of Γ and choose a scaling coset $\Sigma_{\mathbf{p}}$ for Γ at $\mathbf{p}$, and let us temporarily write $c_{\mathbf{p}}$ for the map $\Delta\sigma \mapsto \Delta\sigma \cdot \infty$ of (5.3.2). We claim that $w_{\Gamma}(\mathbf{r})$ is the cardinality of the preimage of $\mathbf{r}$ under $c_{\mathbf{p}}$, for each $\mathbf{r} \in \Delta \backslash \mathbf{p}$. The identity (5.3.6) follows from this because the cardinality $\# \Delta \backslash \Sigma_{\mathbf{p}}$ of the source of $c_{\mathbf{p}}$ is just the number of cosets of Δ in Γ . To compute $\# c_{\mathbf{p}}^{-1}(\mathbf{r})$ suppose $\Delta\sigma \cdot \infty = \Delta\sigma' \cdot \infty$ for some $\sigma, \sigma' \in \Sigma_{\mathbf{p}}$. Then $\sigma^{-1}\delta\sigma' \in (\Sigma_{\mathbf{p}}^{-1}\Sigma_{\mathbf{p}})_{\infty} = B(\mathbb{Z})$ for some $\delta \in \Delta$, so that $\Delta\sigma' = \Delta\sigma T^k$ for some $k \in \mathbb{Z}$. Now $\Delta\sigma T$ is an element of $\Delta \backslash \Sigma_{\mathbf{p}}$ whenever $\Delta\sigma$ is, for $\Sigma_{\mathbf{p}}$ is a union of left cosets of $B(\mathbb{Z})$ by construction. So we have $c_{\mathbf{p}}^{-1}(\mathbf{r}) = \{\Delta\sigma T^k \mid k \in \mathbb{Z}\}$ for any

$\sigma \in \Sigma_{\mathfrak{p}}$ satisfying $\mathfrak{r} = \Delta\sigma \cdot \infty$. If $w_{\Gamma}(\mathfrak{r}) = n$, so that $(\sigma^{-1}\Delta\sigma)_{\infty} = \langle T^n \rangle$ then $\Delta\sigma T^k = \Delta\sigma T^l$ if and only if $n|(k-l)$. This establishes the claim that $\#c_{\mathfrak{p}}^{-1}(\mathfrak{r}) = w_{\Gamma}(\mathfrak{r})$, and completes the proof. $\square$

We can now compute the coefficients b_r^n in (5.3.1) explicitly, and thus obtain a branching theorem for the modified Rademacher sums. Even though we formulated the expression (5.3.1) under the assumption that κ be negative, our methods will apply for all $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$.

Theorem 5.3.4. *Let Δ and Γ be groups commensurable with $G(\mathbb{Z})$. Suppose that Δ is a subgroup of Γ , and suppose that Δ and Γ both have width one at infinity. Let $\mathfrak{p}$ be a cusp of Γ and let $\Sigma_{\mathfrak{p}}$ be a scaling coset for Γ at $\mathfrak{p}$. Let $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$ and let $m \in \mathbb{Z}^+$. Then there exists a system of scaling cosets $\{\Sigma_r \mid \mathfrak{r} \in \mathcal{P}_{\Delta}\}$ for Δ for which we have*

$$Q_{\Gamma, \mathfrak{p}}^{\kappa(m)}(z) = \sum_{\mathfrak{r} \in \Delta \setminus \mathfrak{p}} w_{\Gamma}(\mathfrak{r}) Q_{\Delta, \mathfrak{r}}^{\kappa(mw_{\Gamma}(\mathfrak{r}))}(z). \quad (5.3.7)$$

Proof. Let Δ and Γ be as in the statement of the lemma. Under the assumption that Γ has width one at infinity, and the convention that $\Sigma_{\Gamma \cdot \infty} = \Gamma$ in this case (cf. §2.6), we have

$$Q_{\Gamma, \mathfrak{p}}^{\kappa(m)}(z) = Q_{\Sigma_{\mathfrak{p}}^{-1}}^{\kappa(m)}(z) = \sum_{\Delta\sigma \in \Delta \setminus \Sigma_{\mathfrak{p}}} Q_{\|\sigma^{-1}\Delta}^{\kappa(m)}(z), \quad (5.3.8)$$

and we may attempt to write each summand $Q_{\|\sigma^{-1}\Delta}^{\kappa(m)}(z)$ in terms of the modified Rademacher sums $Q_{\Delta, \mathfrak{r}}^{\kappa(n)}(z)$ associated to Δ . Let $\Delta\sigma \in \Delta \setminus \Sigma_{\mathfrak{p}}$ and set $\mathfrak{r} = \Delta\sigma \cdot \infty$. Supposing that $w_{\Gamma}(\mathfrak{r}) = n$, so that $(\sigma^{-1}\Delta\sigma)_{\infty} = \langle T^n \rangle$, we set $\Sigma_r = \Delta\sigma[n]$. Then we have

$$Q_{\|\sigma^{-1}\Delta}^{\kappa(m)}(z) = Q_{\|[n][1/n]\sigma^{-1}\Delta}^{\kappa(m)}(z) = Q_{\|\Sigma_r^{-1}}^{\kappa(mn)}(z), \quad (5.3.9)$$

since $Q_{\|[n]\chi}^{\kappa(m)}(z, s) = Q_{\|\chi}^{\kappa(mn)}(z, s)$ (cf. §4.1). Under the assumption that Δ has width one at infinity, so that $\Sigma_{\Delta \cdot \infty} = \Delta$, we may rewrite the right most term in (5.3.9) as $Q_{\Delta, \mathfrak{r}}^{\kappa(mn)}(z)$. We see then that a coset $\Delta\sigma$ of Δ in $\Sigma_{\mathfrak{p}}$ contributes a term $Q_{\Delta, \mathfrak{r}}^{\kappa(mw_{\Gamma}(\mathfrak{r}))}(z)$ to the right hand side of (5.3.7), where $\mathfrak{r} = \Delta\sigma \cdot \infty$. According to the proof of Lemma 5.3.3 there are exactly $w_{\Gamma}(\mathfrak{r})$ cosets $\Delta\sigma$ in $\Delta \setminus \Sigma_{\mathfrak{p}}$ satisfying $\mathfrak{r} = \Delta\sigma \cdot \infty$. The required identity (5.3.7) now follows. $\square$

From the proof of Theorem 5.3.4 we see that the existence of the scaling cosets of the conclusion is verified constructively: if $\mathfrak{r} = \Delta\sigma \cdot \infty$ for some coset $\Delta\sigma \in \Delta \setminus \Sigma_{\mathfrak{p}}$, then we may take $\Sigma_r = \Delta\sigma[w_{\Gamma}(\mathfrak{r})]$.

By utilizing the inner products $\langle \cdot, \cdot \rangle_{\Delta}^{\kappa}$ of §5.2 we can reformulate Theorem 5.3.4 in a way that is independent of scaling coset choices.

Theorem 5.3.5. *Let Δ and Γ be groups commensurable with $G(\mathbb{Z})$. Suppose that Δ is a subgroup of Γ , and suppose that Δ and Γ both have width one at infinity. Let $\mathfrak{p}$ be a cusp of Γ , let $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$ and let $m \in \mathbb{Z}^+$. Then we have*

$$\left| \left\langle Q_{\Gamma, \mathfrak{p}}^{\kappa(m)}, Q_{\Delta, r}^{\kappa(n)} \right\rangle_{\Delta}^{\kappa} \right|^2 = w_{\Gamma}(r) \delta_{\Gamma, \mathfrak{p} | \Gamma r} \delta_{mw_{\Gamma}(r), n} \quad (5.3.10)$$

for all $r \in P_{\Delta}$ and $n \in \mathbb{Z}^+$, for any choice of scaling coset systems for Δ and Γ .

To conclude this section we comment on the branching of Rademacher sums in the case that Γ both contains and normalizes Δ . Suppose then that Δ and Γ have width one at infinity, and consider the case that $\mathfrak{p}$ is the infinite cusp $\Gamma \cdot \infty$ in (5.3.2). Then we are speaking of the map

$$\begin{aligned} \Delta \backslash \Gamma &\rightarrow \Delta \backslash \Gamma \cdot \infty \\ \Delta \gamma &\mapsto \Delta \gamma \cdot \infty, \end{aligned} \quad (5.3.11)$$

and this map is injective (and thus bijective) when Γ normalizes Δ . Indeed, since Δ is supposed to have width one at infinity we have $(\gamma^{-1} \Delta \gamma)_{\infty} = \Delta_{\infty} = B(\mathbb{Z})$, so that every coset $\Delta \gamma \in \Delta \backslash \Gamma$ is indeed a scaling coset for Δ at $r = \Delta \gamma \cdot \infty$, and the injectivity of (5.3.11) follows from Lemma 5.3.1. Applying Theorem 5.3.4 now with $\mathfrak{p} = \Gamma \cdot \infty$ we obtain the following result.

Proposition 5.3.6. *Suppose that Δ and Γ are groups commensurable with $G(\mathbb{Z})$ that both have width one at infinity, and suppose that Γ contains and normalizes Δ . Let $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$ and let $m \in \mathbb{Z}^+$. Then we have*

$$Q_{\Gamma}^{\kappa(m)}(z) = \sum_{r \in \Delta \backslash \mathfrak{p}} Q_{\Delta, r}^{\kappa(m)}(z) \quad (5.3.12)$$

when the scaling cosets for Δ at the cusps $r \in \Delta \backslash \Gamma \cdot \infty$ are taken to lie in $\Delta \backslash \Gamma$.

Theorem 5.3.5 now implies the following reformulation of Proposition 5.3.6 which is independent of scaling coset choices.

Proposition 5.3.7. *Suppose that Δ and Γ are groups commensurable with $G(\mathbb{Z})$ that both have width one at infinity, and suppose that Γ contains and normalizes Δ . Then we have*

$$\left| \left\langle Q_{\Gamma}^{\kappa(1)}, Q_{\Delta, r}^{\kappa(1)} \right\rangle_{\Delta}^{\kappa} \right|^2 = \delta_{\Gamma, \mathfrak{p} | \Gamma r} \quad (5.3.13)$$

for all $r \in P_{\Delta}$, for any choice of scaling coset system for Δ .

5.4 Fractional orders

Let Γ be a group commensurable with $G(\mathbb{Z})$, let $\phi, \psi \in G(\mathbb{Q})$ and set $U = \llbracket \phi^{-1} \Gamma \psi \rrbracket$. Let us consider the problem of writing the modified Rademacher sum $Q_U^{\kappa(m)}(z)$ in terms of Rademacher

sums of the form $Q_{\Gamma', \mathbf{p}'}^{\kappa(m')}(z)$ for some group Γ' and some cusp $\mathbf{p}' \in \mathbf{P}_{\Gamma'}$. According to §4.1 we have $Q_{\Gamma, \mathbf{p}|\mathbf{q}}^{\kappa(m)}(z) = Q_{\Gamma', \mathbf{p}'}^{\kappa(m)}(z)$ for $\Gamma' = \Gamma^{\mathfrak{q}}$ and $\mathbf{p}' = \mathbf{p}^{\mathfrak{q}}$ (cf. (4.1.9)), so we are done if we can write $Q_U^{\kappa(m)}(z)$ in terms of $Q_{\Gamma, \mathbf{p}|\mathbf{q}}^{\kappa(n)}(z)$ for some cusps $\mathbf{p}, \mathbf{q} \in \mathbf{P}_{\Gamma}$ for Γ . Set $\mathbf{p} = \Gamma\phi \cdot \infty$ and $\mathbf{q} = \Gamma\psi \cdot \infty$. By the discussion of §2.6 there exist unique $\mu, \nu \in \mathbb{Q}^+$ such that the cosets $\Sigma_{\mathbf{p}} = \Gamma\phi[\mu]$ and $\Sigma_{\mathbf{q}} = \Gamma\psi[\nu]$ (cf. (2.3.3)) are scaling cosets for Γ at $\mathbf{p}$ and $\mathbf{q}$, respectively. We then have $U = \llbracket [\mu]\Sigma_{\mathbf{p}}^{-1}\Sigma_{\mathbf{q}}[1/\nu] \rrbracket$. Let $\chi \in \Sigma_{\mathbf{q}}^{-1}\Sigma_{\mathbf{p}}$ and consider the contribution $Q_{\llbracket [\mu]\chi[1/\nu] \rrbracket}^{\kappa(m)}(z, s)$ of the coset $\llbracket [\mu]\chi[1/\nu] \rrbracket \in U$ to the modified continued Rademacher sum $Q_U^{\kappa(m)}(z, s)$. As the following lemma demonstrates, the factor $[1/\nu]$ induces a re-scaling of the input variable z .

Lemma 5.4.1. *Let $\chi \in G(\mathbb{Q})$ and let $\nu \in \mathbb{Q}^+$. Let $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$ and let $m \in \mathbb{Z}^+$. Then we have*

$$Q_{\llbracket \chi[1/\nu] \rrbracket}^{\kappa(m)}(z, s) = \frac{1}{\nu^{\kappa}} Q_{\llbracket \chi \rrbracket}^{\kappa(m)}\left(\frac{z}{\nu}, s\right). \quad (5.4.1)$$

Proof. We recall from (4.1.4) that the modified continued Rademacher component function $Q_{\llbracket \chi[1/\nu] \rrbracket}^{\kappa(m)}(z, s)$ is, by definition, the difference $T_{\llbracket \chi[1/\nu] \rrbracket}^{\kappa(m)}(z, s) - T_{0, \llbracket \chi[1/\nu] \rrbracket}^{\kappa(m)}(s)$, where the functions $T_{\llbracket \chi[1/\nu] \rrbracket}^{\kappa(m)}(z, s)$ and $T_{0, \llbracket \chi[1/\nu] \rrbracket}^{\kappa(m)}(s)$ are defined by (4.1.1) and (4.1.3), respectively. We compute $\text{jac}(\llbracket \chi[1/\nu] \rrbracket, z) = \text{jac}(\llbracket \chi \rrbracket, z/\nu)/\nu$, and use the fact that $[1/\nu]$ fixes both ∞ and 0 to verify that

$$T_{\llbracket \chi[1/\nu] \rrbracket}^{\kappa(m)}(z, s) = \frac{1}{\nu^{\kappa}} T_{\llbracket \chi \rrbracket}^{\kappa(m)}\left(\frac{z}{\nu}, s\right), \quad T_{0, \llbracket \chi[1/\nu] \rrbracket}^{\kappa(m)}(s) = \frac{1}{\nu^{\kappa}} T_{0, \llbracket \chi \rrbracket}^{\kappa(m)}(s). \quad (5.4.2)$$

The claim follows from these identities. $\square$

We now seek to describe the effect of the factor $[\mu]$ in $Q_{\llbracket [\mu]\chi \rrbracket}^{\kappa(m)}(z, s)$, for arbitrary $\chi \in G(\mathbb{Q})$. Inspecting (4.1.1) we find that

$$T_{\llbracket [\mu]\chi \rrbracket}^{\kappa(m)}(z, s) = \mu^{\kappa} e(-m\mu\chi \cdot z) t^{\kappa}(m\mu, \chi, z, s) \text{jac}(\chi, z)^{\kappa}, \quad (5.4.3)$$

which suggests that we generalize the notion of Rademacher sum so as to allow for fractional orders. Recall from (2.3.5) that $B(\alpha\mathbb{Z})$ denotes the subgroup of $B_u(\mathbb{Q})$ generated by T^{α} (cf. (2.3.4)). For general $\mu \in \mathbb{Q}^+$ the right hand side of (5.4.3) will not be invariant under the replacement of χ by $T\chi$, so a Rademacher sum of fractional order is not naturally defined by collections of cosets of $B(\mathbb{Z})$, but rather, by cosets of $B(h\mathbb{Z})$ for a suitably chosen positive integer $h \in \mathbb{Z}^+$.

Given $h \in \mathbb{Z}^+$ and $\chi \in G(\mathbb{Q})$ let us write $\llbracket^h \chi$ as a shorthand for $B(h\mathbb{Z})\chi$, and for $X \subset G(\mathbb{Q})$ let us write $\llbracket^h X$ as a shorthand for the set of right cosets of $B(h\mathbb{Z})$ determined by elements of X , so that

$$\llbracket^h X = \{B(h\mathbb{Z})\chi \mid \chi \in X\}. \quad (5.4.4)$$

Let $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$. We define modified and normalized fractional Rademacher sums as follows, in analogy with the constructions of §4.1. For $\mu = g/h$ with $(g, h) = 1$ and $g, h \in \mathbb{Z}^+$, and for $[\![^h\chi] \in [\![^hG(\mathbb{Q})$, we define the *continued fractional Rademacher component function of weight 2κ and order μ associated to $[\![^h\chi$* , denoted $(z, s) \mapsto T_{[\![^h\chi}^{\kappa(\mu)}(z, s)$, by setting

$$T_{[\![^h\chi}^{\kappa(\mu)}(z, s) = e(-\mu [\![^h\chi \cdot z) t^\kappa(\mu, [\![^h\chi, z, s) \text{jac}([\![^h\chi, z)^\kappa, \quad (5.4.5)$$

where $\text{jac}([\![^h\chi, z) = \text{jac}(\chi, z)$, and $t^\kappa(\mu, [\![^h\chi, z, s)$ is the *continued fractional Rademacher regularization factor of weight 2κ* , which is in turn given by

$$t^\kappa(\mu, [\![^h\chi, z, s) = \Phi(s - 2\kappa, 1 + s - 2\kappa, \mu [\![^h\chi \cdot z - \mu [\![^h\chi \cdot \infty)(2\pi i(\mu [\![^h\chi \cdot \infty - \mu [\![^h\chi \cdot z))^{s-2\kappa} \quad (5.4.6)$$

in case $[\![^h\chi \in [\![^hG(\mathbb{Q})^\times$, and $t^\kappa(\mu, [\![^h\chi, z, s) = 1$ otherwise. We define a function $s \mapsto T_{0, [\![^h\chi}^{\kappa(\mu)}(s)$ by setting

$$\begin{aligned} T_{0, [\![^h\chi}^{\kappa(\mu)}(s) &= e(-\mu [\![^h\chi \cdot \infty) (2\pi i(\mu [\![^h\chi \cdot \infty - \mu [\![^h\chi \cdot 0))^{(s-2\kappa)} \text{jac}([\![^h\chi, 0)^\kappa \\ &\quad - e(s/2) e(-\mu [\![^h\chi \cdot \infty) (2\pi i(\mu [\![^h\chi \cdot 0 - \mu [\![^h\chi \cdot \infty))^{(s-2\kappa)} \text{jac}([\![^h\chi, 0)^\kappa \end{aligned} \quad (5.4.7)$$

in case $[\![^h\chi \in [\![G(\mathbb{Q})^{\times \times}$, and by setting $T_{0, [\![^h\chi}^{\kappa(\mu)}(s) = 0$ otherwise, and we define the *modified continued fractional Rademacher component function of weight 2κ and order μ associated to $[\![^h\chi$* , denoted $(z, s) \mapsto Q_{[\![^h\chi}^{\kappa(\mu)}(z, s)$, by subtracting $T_{0, [\![^h\chi}^{\kappa(\mu)}(z, s)$ from $T_{[\![^h\chi}^{\kappa(\mu)}(z, s)$.

$$Q_{[\![^h\chi}^{\kappa(\mu)}(z, s) = T_{[\![^h\chi}^{\kappa(\mu)}(z, s) - T_{0, [\![^h\chi}^{\kappa(\mu)}(z, s) \quad (5.4.8)$$

For $U \subset [\![^hG(\mathbb{Q})$ we now define the *continued fractional Rademacher sum of weight 2κ and order μ associated to U* , denoted $T_U^{\kappa(\mu)}(z, s)$, and the *modified continued fractional Rademacher sum of weight 2κ and order μ associated to U* , denoted $Q_U^{\kappa(\mu)}(z, s)$, by setting

$$T_U^{\kappa(\mu)}(z, s) = \sum_{[\![^h\chi \in U} T_{[\![^h\chi}^{\kappa(\mu)}(z, s), \quad (5.4.9)$$

$$Q_U^{\kappa(\mu)}(z, s) = \sum_{[\![^h\chi \in U} Q_{[\![^h\chi}^{\kappa(\mu)}(z, s), \quad (5.4.10)$$

and we define $T_U^{\kappa(\mu)}(z)$ and $Q_U^{\kappa(\mu)}(z)$ by taking the limit as s tends to 1 in $T_U^{\kappa(\mu)}(z, s)$ and

$Q_U^{\kappa(\mu)}(z, s)$, respectively.

$$T_U^{\kappa(\mu)}(z) = \lim_{s \rightarrow 1^+} T_U^{\kappa(\mu)}(z, s) \quad (5.4.11)$$

$$Q_U^{\kappa(\mu)}(z) = \lim_{s \rightarrow 1^+} Q_U^{\kappa(\mu)}(z, s) \quad (5.4.12)$$

We call $T_U^{\kappa(\mu)}(z)$ the *normalized fractional Rademacher sum of weight 2κ and order μ associated to U* , and we call $Q_U^{\kappa(\mu)}(z)$ the *modified fractional Rademacher sum of weight 2κ and order μ associated to U* .

We have the following vanishing result for fractional Rademacher sums.

Proposition 5.4.2. *Let X be a union of cosets of $B(\mathbb{Z})$ in $G(\mathbb{Q})$, let $\mu = g/h$ for some $g, h \in \mathbb{Z}^+$ and $(g, h) = 1$, and set $U = \llbracket^h X$. Suppose that the normalized fractional Rademacher sum $T_U^{\kappa(\mu)}(z)$ and the modified fractional Rademacher sum $Q_U^{\kappa(\mu)}(z)$ converge. Then they vanish identically unless $h = 1$.*

Proof. Suppose that $h \neq 1$. By hypothesis we have a disjoint decomposition $X = \bigcup_i \llbracket \chi_i$ for some $\{\chi_i\} \subset X$. This implies a disjoint decomposition $X = \bigcup_i \bigcup_{k=0}^{h-1} \llbracket^h T^k \chi_i$, so that we have

$$\begin{aligned} T_U^{\kappa(\mu)}(z, s) &= \sum_i \sum_{k=0}^{h-1} e\left(-\frac{g}{h} T^k \chi_i \cdot z\right) t^\kappa\left(\frac{g}{h}, T^k \chi_i, z, s\right) \text{jac}(T^k \chi_i, z)^\kappa \\ &= \sum_i \left(\sum_{k=0}^{h-1} e\left(-\frac{gk}{h}\right) \right) e\left(-\frac{g}{h} \chi_i \cdot z\right) t^\kappa\left(\frac{g}{h}, \chi_i, z, s\right) \text{jac}(\chi_i, z)^\kappa \end{aligned} \quad (5.4.13)$$

since the continued fractional regularization factor is unaffected when the second argument is multiplied by an element of $B(\mathbb{Z})$ on the left. The sum $\sum_{k=0}^{h-1} e(-gk/h)$ vanishes for g coprime to h unless $h = 1$. This shows that the continued fractional Rademacher sum $T_U^{\kappa(\mu)}(z, s)$ vanishes for all z and s when $h \neq 1$. A directly analogous computation shows that the function $T_{0,U}^{\kappa(\mu)}(s)$ vanishes in case $h \neq 1$. We conclude that both the continued fractional Rademacher sum $T_U^{\kappa(\mu)}(z, s)$ and the modified continued fractional Rademacher sum $Q_U^{\kappa(\mu)}(z, s)$ vanish identically in case $h \neq 1$. The claim of the proposition follows. $\square$

We return now to the identification of the Rademacher sum $Q_U^{\kappa(m)}(z)$, in the case that $U = \llbracket [\mu] \Sigma_{\mathfrak{p}}^{-1} \Sigma_{\mathfrak{q}} [1/\nu]$.

Theorem 5.4.3. *Let Γ be a group commensurable with $G(\mathbb{Z})$, let $\mathfrak{p}, \mathfrak{q} \in \mathcal{P}_\Gamma$ be cusps for Γ , and let $\Sigma_{\mathfrak{p}}$ and $\Sigma_{\mathfrak{q}}$ be scaling cosets for Γ at $\mathfrak{p}$ and $\mathfrak{q}$, respectively. Let $\mu, \nu \in \mathbb{Q}^+$ and set $U = \llbracket [\mu] \Sigma_{\mathfrak{p}}^{-1} \Sigma_{\mathfrak{q}} [1/\nu]$. Let $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$, and let $m \in \mathbb{Z}^+$. Let h be the smallest*

positive integer such that $h\mu \in \mathbb{Z}$. If h divides m then we have

$$Q_U^{\kappa(m)}(z) = h \frac{\mu^\kappa}{\nu^\kappa} Q_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m\mu)}\left(\frac{z}{\nu}\right), \quad (5.4.14)$$

and if h does not divide m then $Q_U^{\kappa(m)}(z)$ vanishes identically.

Proof. Set $U' = \llbracket [\mu] \Sigma_{\mathfrak{p}}^{-1} \Sigma_{\mathfrak{q}} \rrbracket$ and $U'' = \llbracket \Sigma_{\mathfrak{p}}^{-1} \Sigma_{\mathfrak{q}} \rrbracket$, so that $U = U'[1/\nu] = [\mu]U''[1/\nu]$. By Lemma 5.4.1 we have $Q_U^{\kappa(m)}(z) = Q_{U'}^{\kappa(m)}(z/\nu)/\nu^\kappa$, so it suffices for us to show that $Q_{U'}^{\kappa(m)}(z)$ is $h\mu^\kappa Q_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m\mu)}(z)$ or vanishing, according as h divides m or not.

Set $X = \Sigma_{\mathfrak{p}}^{-1} \Sigma_{\mathfrak{q}} \subset G(\mathbb{Q})$. Then X is a union of right cosets of $B(\mathbb{Z})$, so we have a disjoint decomposition $X = \bigcup_i \llbracket \chi_i \rrbracket$ for some $\chi_i \in G(\mathbb{Q})$. By the choice of h we have $\mu = g/h$ for some $g \in \mathbb{Z}^+$ with $(g, h) = 1$. Then for the set $[\mu]X$ we have

$$[\mu]X = \bigcup_i [\mu] \llbracket \chi_i \rrbracket = \bigcup_i \bigcup_{k=0}^{h-1} [\mu] T^k \llbracket \chi_i \rrbracket = \bigcup_i \bigcup_{k=0}^{h-1} T^{k\mu} \llbracket [\mu] \chi_i \rrbracket \quad (5.4.15)$$

with all the unions disjoint, from which we conclude that $\llbracket [\mu] X \rrbracket$ admits the disjoint decomposition $\llbracket [\mu] X \rrbracket = \bigcup_{k=0}^{h-1} \bigcup_i T^{k\mu} \llbracket [\mu] \chi_i \rrbracket$. We have

$$Q_{T^{k\mu} \llbracket [\mu] \chi_i \rrbracket}^{\kappa(m)}(z, s) = e(-mk\mu) Q_{\llbracket [\mu] \chi_i \rrbracket}^{\kappa(m)}(z, s). \quad (5.4.16)$$

If h divides m then $m\mu \in \mathbb{Z}$, and $e(-mk\mu) = 1$ for all k . Further, we have $Q_{\llbracket [\mu] \chi_i \rrbracket}^{\kappa(m)}(z, s) = \mu^\kappa Q_{\llbracket \chi_i \rrbracket}^{\kappa(m\mu)}(z, s)$ in this case, and so we deduce the required identity $Q_{U'}^{\kappa(m)}(z) = h\mu^\kappa Q_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{\kappa(m\mu)}(z)$. On the other hand, if h does not divide m , so that $m\mu$ is not an integer, then the sum $\sum_{k=0}^{h-1} e(-mk\mu)$ vanishes, and this implies the vanishing of the modified Rademacher sum $Q_{U'}^{\kappa(m)}(z)$. The proof is complete. $\square$

Taking $\mathfrak{q} = \Gamma \cdot \infty$ in Theorem 5.4.3 we obtain the following result.

Theorem 5.4.4. *Let Γ be a group commensurable with $G(\mathbb{Z})$, let Z be a finite union of left cosets of Γ in $G(\mathbb{Q})$ and set $U = \llbracket Z \rrbracket$. Let $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$ and let $m \in \mathbb{Z}^+$. Then the modified Rademacher sum $Q_U^{\kappa(m)}(z)$ is an automorphic integral of weight 2κ for Γ .*

5.5 Hecke operators

For the modified Rademacher sums, in contrast to the classical Rademacher sums, it is natural to consider not only their variance with respect to group actions, but also how they vary with respect to Hecke operators. The discrepancy in constant terms between the classical and modified Rademacher sums illustrated by Proposition 4.4.10 is a stubborn barrier to an exposition of the interaction between Hecke operators and the classical Rademacher sums.

Recall that $I_\kappa(\Gamma)$ denotes the space of automorphic integrals of weight 2κ for Γ , in the sense of §2.7, and recall the operator $I_\kappa(\Gamma) \rightarrow \mathcal{O}(\mathbb{H})$ of (2.7.14), denoted $f \mapsto f|_\Gamma^\kappa X$, and defined for a right coset $X \in \Gamma \backslash G(\mathbb{Q})$ by setting $f|_\Gamma^\kappa X = (f - J_{\chi, \infty}^\kappa g)|^\kappa \chi$ for any representative $\chi \in X$ where $g \in S_{1-\kappa}(\Gamma)$ is the cusp form associated to f (cf. (2.7.13)). Let us generalize this operator by setting

$$f|_\Gamma^\kappa X = \sum_i (f - J_{\chi_i, \infty}^\kappa g)|^\kappa \chi_i \quad (5.5.1)$$

in case X is a finite union of right cosets of Γ in $G(\mathbb{Q})$ and the χ_i furnish a transversal $X = \bigcup_i \Gamma \chi_i$ for X over Γ . The next lemma verifies that the operator $f \mapsto f|_\Gamma^\kappa X$ makes sense in case X is a double coset $X = \Gamma \sigma \Gamma$ of Γ in $G(\mathbb{Q})$.

Lemma 5.5.1. *Let Γ be a group commensurable with $G(\mathbb{Z})$, let $\sigma \in G(\mathbb{Q})$, set $\Delta = \Gamma \cap \sigma^{-1} \Gamma \sigma$ and $\Delta' = \sigma \Gamma \sigma^{-1} \cap \Gamma$, and suppose that*

$$\Gamma = \bigcup_i \lambda'_i \Delta', \quad \Gamma = \bigcup_i \Delta \rho_i, \quad (5.5.2)$$

are left and right transversals for Γ over Δ' and Δ , respectively. Then the double coset $\Gamma \sigma \Gamma$ admits disjoint decompositions

$$\Gamma \sigma \Gamma = \bigcup_i \lambda'_i \sigma \Gamma, \quad \Gamma \sigma \Gamma = \bigcup_i \Gamma \sigma \rho_i, \quad (5.5.3)$$

into left and right cosets for Γ .

Proof. Since $G(\mathbb{Q})$ is the commensurator of $G(\mathbb{Z})$, the intersection $\Delta = \Gamma \cap \sigma^{-1} \Gamma \sigma$ has finite index in both Γ and $\sigma^{-1} \Gamma \sigma$, so there are only finitely many λ'_i and ρ_i . Observe next that $\sigma \Delta = \Delta' \sigma = \sigma \Gamma \cap \Gamma \sigma$. Consequently, we have $\Delta' \sigma \subset \sigma \Gamma$ and $\sigma \Delta \subset \Gamma \sigma$, so that

$$\Gamma \sigma \Gamma = \bigcup \lambda'_i \Delta' \sigma \Gamma \subset \bigcup \lambda'_i \sigma \Gamma \Gamma = \bigcup \lambda'_i \sigma \Gamma, \quad (5.5.4)$$

$$\Gamma \sigma \Gamma = \bigcup \Gamma \sigma \Delta \rho_i \subset \bigcup \Gamma \sigma \rho_i = \bigcup \Gamma \sigma \rho_i. \quad (5.5.5)$$

The reverse inclusions hold since all the λ'_i and ρ_i lie in Γ . The unions are disjoint, for if $\Gamma \sigma \rho_i = \Gamma \sigma \rho_j$ say, then $\rho_i \rho_j^{-1} \in \sigma^{-1} \Gamma \sigma$, but $\rho_i \rho_j^{-1} \in \Gamma$ by our choice of the ρ_i , so $\rho_i \rho_j^{-1} \in \Delta$, and this implies $i = j$. A similar argument applies to the cosets $\lambda'_i \sigma \Gamma$. $\square$

The right hand identity of (5.5.3) shows that the operator $f \mapsto f|_\Gamma^\kappa \Gamma \sigma \Gamma$ is well-defined for $f \in I_\kappa(\Gamma)$ and $\sigma \in G(\mathbb{Q})$. The left hand identity of (5.5.3) shows that the function $f|_\Gamma^\kappa \Gamma \sigma \Gamma$ again lies in $I_\kappa(\Gamma)$. Indeed, if the cusp form associated to $f \in I_\kappa(\Gamma)$ is $g \in S_{1-\kappa}(\Gamma)$ then the cusp form associated to $f|_\Gamma^\kappa \Gamma \sigma \Gamma$ is $g|_\Gamma^\kappa \Gamma \sigma \Gamma$. We call the operator $I_\kappa(\Gamma) \rightarrow I_\kappa(\Gamma)$ given by $f \mapsto f|_\Gamma^\kappa \Gamma \sigma \Gamma$

the *weight* 2κ *Hecke operator associated to* σ . In order to ease notation we set

$$f \dagger_{\Gamma}^{\kappa} \sigma = f \|_{\Gamma}^{\kappa} \Gamma \sigma \Gamma \quad (5.5.6)$$

for $\sigma \in G(\mathbb{Q})$. We have $f \dagger_{\Gamma}^{\kappa} \sigma = \sum_i f |^{\kappa} \chi_i$ when $\{\chi_i\} \subset G(\mathbb{Q})$ is a right transversal $\Gamma \sigma \Gamma = \bigcup_i \Gamma \chi_i$ for Γ in $\Gamma \sigma \Gamma$. If σ belongs to the normalizer of Γ then we have $\Gamma \sigma \Gamma = \Gamma \sigma$ and hence $f \dagger_{\Gamma}^{\kappa} \sigma = (f - J_{\sigma \cdot \infty}^{\kappa}) |^{\kappa} \sigma$ for any $f \in I_{\kappa}(\Gamma)$. More generally, Lemma 5.5.1 yields for us the formula

$$f \dagger_{\Gamma}^{\kappa} \sigma = \sum_i (f - J_{\sigma \rho_i \cdot \infty}^{\kappa}) |^{\kappa} (\sigma \rho_i) \quad (5.5.7)$$

in the case that $\{\rho_i\}$ is a right transversal for Γ over the intersection $\Delta = \Gamma \cap \Gamma^{\sigma}$.

The action of the operator $f \mapsto f \|_{\Gamma}^{\kappa} X$ on holomorphic Poincaré series can be described without reference to transversals. Indeed, if U is the set of right cosets of $B(\mathbb{Z})$ determined by a union of left cosets of some group Γ commensurable with $G(\mathbb{Z})$, and if X is a finite union $\bigcup_j \Gamma \chi_j$ say, of right cosets of Γ , then, taking $\kappa > 1$ to ensure absolute convergence, we have

$$P_U^{\kappa(m)} \|_{\Gamma}^{\kappa} X = \sum_j P_U^{\kappa(m)} |^{\kappa} \chi_j = \sum_j P_{U \chi_j}^{\kappa(m)} = P_{UX}^{\kappa(m)}, \quad (5.5.8)$$

so that the Poincaré series are stable under the action of the operators $f \mapsto f \|_{\Gamma}^{\kappa} X$. In particular, for the Hecke operator $f \mapsto f \dagger_{\Gamma}^{\kappa} \sigma$ we have $P_U^{\kappa(m)} \dagger_{\Gamma}^{\kappa} \sigma = P_{U \sigma \Gamma}^{\kappa(m)}$. This identity $P_U^{\kappa(m)} \|_{\Gamma}^{\kappa} X = P_{UX}^{\kappa(m)}$ extends naturally to the modified Rademacher sums.

Proposition 5.5.2. *Let Γ be a group commensurable with $G(\mathbb{Z})$, let Z be a finite union of left cosets of Γ in $G(\mathbb{Q})$ and set $U = \llbracket Z$. Let $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$ and let $m \in \mathbb{Z}^+$. Then for X a finite union of right cosets of Γ in $G(\mathbb{Q})$ we have*

$$Q_U^{\kappa(m)} \|_{\Gamma}^{\kappa} X = Q_{UX}^{\kappa(m)}. \quad (5.5.9)$$

Proof. The modified Rademacher sum $Q_U^{\kappa(m)}(z)$ is an automorphic integral of weight 2κ for Γ according to Theorem 5.4.4, so it has an associated cusp form $g \in S_{1-\kappa}(\Gamma)$ say. We may express X as a disjoint union $X = \bigcup_i \Gamma \chi_i$ for some finitely many $\chi_i \in G(\mathbb{Q})$. We then compute

$$Q_U^{\kappa(m)} \|_{\Gamma}^{\kappa} X = \sum_i (Q_U^{\kappa(m)} - J_{\chi_i \cdot \infty}^{\kappa}) |^{\kappa} \chi_i = \sum_i Q_{U \chi_i}^{\kappa(m)} = Q_{UX}^{\kappa(m)}. \quad (5.5.10)$$

This proves the claim. $\square$

In the case that Γ is the modular group $G(\mathbb{Z})$ we have the classical Hecke operators $\hat{T}(n)$,

defined for $n \in \mathbb{Z}^+$ by setting

$$n^{1-\kappa}(\hat{T}(n)f)(z) = \sum_{\substack{ad=n \\ 0 \leq b < d}} f\left(\frac{az+b}{d}\right) \frac{a^\kappa}{d^\kappa} \quad (5.5.11)$$

for $f \in M_\kappa(\Gamma)$. We deduce the relation between the operators $\hat{T}(n)$ and $f \mapsto f \dagger_\Gamma^\kappa \sigma$, for $\Gamma = G(\mathbb{Z})$, by observing the following coincidence of disjoint unions of cosets and double cosets of Γ .

$$\bigcup_{\substack{ad=n \\ 0 \leq b < d}} \Gamma \begin{bmatrix} a & b \\ 0 & d \end{bmatrix} = \bigcup_{\substack{ad=n \\ d|a}} \Gamma \begin{bmatrix} a & 0 \\ 0 & d \end{bmatrix} \Gamma \quad (5.5.12)$$

Both sides of (5.5.12) are decompositions of the image in $G(\mathbb{Q})$ of the set of 2×2 matrices with integral entries and determinant n . From these decompositions we deduce the following result.

Lemma 5.5.3. *For $n \in \mathbb{Z}^+$ and $\kappa \in \mathbb{Z}$ we have*

$$n^{1-\kappa}(\hat{T}(n)f) = \sum_{d \in \mathbb{Z}^+, d^2 | n} f \dagger_\Gamma^\kappa [n/d^2] \quad (5.5.13)$$

for any $f \in I_\kappa(\Gamma)$.

As a partial converse to Lemma 5.5.3, observe that for $\Gamma = G(\mathbb{Z})$ the action of any Hecke operator $f \mapsto f \dagger_\Gamma^\kappa \sigma$ can be expressed in terms of the Hecke operator associated to a diagonal element $[\mu] \in B(\mathbb{Q})$. Indeed, if $\sigma \in B(\mathbb{Q})$ then, according to (5.5.12), we have $\Gamma\sigma\Gamma = \Gamma[\mu]\Gamma$ for some $\mu \in \mathbb{Q}^+$. In case σ does not lie in $B(\mathbb{Q})$ we have $\sigma \cdot \infty = p \neq \infty$ for some $p \in \mathbb{Q}$. Since the modular group acts transitively on $\hat{\mathbb{Q}}$ there is some $\sigma_p \in \Gamma$ with $\sigma_p \cdot \infty = p$, and then $\Gamma\sigma\Gamma = \Gamma\tilde{\sigma}\Gamma$ where $\tilde{\sigma} = \sigma_p^{-1}\sigma$ evidently lies in $B(\mathbb{Q})$.

Lemma 5.5.4. *Let $\Gamma = G(\mathbb{Z})$ and let $\sigma \in G(\mathbb{Q})$. Then there exists $\mu \in \mathbb{Q}^+$ such that*

$$f \dagger_\Gamma^\kappa \sigma = f \dagger_\Gamma^\kappa [\mu] \quad (5.5.14)$$

for all $\kappa \in \mathbb{Z}$ and all $f \in I_\kappa(\Gamma)$.

We now consider the action of the Hecke operators $\hat{T}(n)$ on the modified Rademacher sum $Q_\Gamma^{(1)}(z) = Q_{\llbracket \Gamma \rrbracket}^{(1)}(z)$ of weight 0 and order 1 associated to the modular group $\Gamma = G(\mathbb{Z})$. From Theorem 4.5.3 we have that $Q_{\llbracket \Gamma \rrbracket}^{(m)}(z)$ is an automorphic integral of weight 0 (that is, an abelian integral) for Γ . Since there are no non-zero cusp forms of weight 2 for the modular group, we see from Corollary 4.5.5 that $Q_{\llbracket \Gamma \rrbracket}^{(m)}(z)$ is in fact a Γ -invariant function on $\mathbb{H}$ for all $m \in \mathbb{Z}^+$. Consequently we have $Q_{\llbracket \Gamma \rrbracket}^{(1)} \dagger_\Gamma^0 [n] = Q_{\llbracket \Gamma[n] \rrbracket}^{(1)}$ for $n \in \mathbb{Z}^+$, by Proposition 5.5.2 and the definition (5.5.6) of $f \dagger_\Gamma^\kappa \sigma$. Suppose that n is square-free. Then, according to Lemma 5.5.3, the action of the operator $f \mapsto f \dagger_\Gamma^0 [n]$ on $M_0(\Gamma)$ coincides with that of $n\hat{T}(n)$. We anticipate an application

of Lemma 5.5.1. For $\sigma = [n]$ we have $\Delta' = \sigma\Gamma\sigma^{-1} \cap \Gamma = \Gamma^0(n)$, and for a left transversal of $\Delta' = \Gamma^0(n)$ in Γ we may take

$$\Gamma = \bigcup_{e \parallel n} \bigcup_{k=0}^{n/e-1} T^k ST^e \Delta' \quad (5.5.15)$$

where the first union is over exact divisors of n (cf. §2.1). We deduce that

$$\Gamma[n]\Gamma = \bigcup_{e \parallel n} \bigcup_{k=0}^{n/e-1} T^k ST^e [n]\Gamma \quad (5.5.16)$$

by Lemma 5.5.1. Now $\llbracket T^k X = \llbracket X$ for any subset $X \subset G(\mathbb{Q})$ and any $k \in \mathbb{Z}$, so we find that

$$n(\hat{T}(n)Q_{\llbracket \Gamma}^{(1)}) = Q_{\llbracket \Gamma[n]\Gamma}^{(1)} = \sum_{e \parallel n} Q_{\llbracket ST^e [n]\Gamma}^{(1)}. \quad (5.5.17)$$

Lemma 5.5.5. *Let e be an exact divisor of n . Then we have $\llbracket ST^e [n]\Gamma = \llbracket [n/e^2]\Gamma$ when Γ is the modular group $G(\mathbb{Z})$.*

Proof. We compute

$$ST^e [n] = \begin{bmatrix} 0 & -1 \\ n & e \end{bmatrix}, \quad ST^e [n] ST^{n/e} S = \begin{bmatrix} n/e & -1 \\ 0 & e \end{bmatrix}, \quad (5.5.18)$$

and observe that $T^k AT^l = [n/e^2]$ for $k, l \in \mathbb{Z}$ such that $ke + ln/e = 1$, where A is given by $A = ST^e [n] ST^{n/e} S$. $\square$

We can now write

$$n(\hat{T}(n)Q_{\llbracket \Gamma}^{(1)}) = \sum_{e \parallel n} Q_{\llbracket [n/e^2]\Gamma}^{(1)} \quad (5.5.19)$$

in the case that n is square-free, and we can employ the methods of §5.4 to rewrite (5.5.19) in terms of the fractional Rademacher sums. Observe that $B(g\mathbb{Z})[g/h] = [g/h]B(h\mathbb{Z})$. We thus have

$$\llbracket [\mu]\chi = \bigcup_{k=0}^{g-1} T^k \llbracket [g[\mu]]\chi = \bigcup_{k=0}^{g-1} T^k [\mu] \llbracket^h \chi \quad (5.5.20)$$

for any $\chi \in G(\mathbb{Q})$ when $\mu = g/h$ with $g, h \in \mathbb{Z}^+$ and $(g, h) = 1$. We rewrite the right hand side

of (5.5.19), taking $g = n/h$ and $h = e$, as

$$\sum_{h \parallel n} Q_{\llbracket n/h^2 \rrbracket \Gamma}^{(1)} = \sum_{h \parallel n} Q_{\llbracket n/h^2 \rrbracket \llbracket^h \Gamma}^{(1)} = \sum_{h \parallel n} Q_{\llbracket^h \Gamma}^{(n/h^2)}, \quad (5.5.21)$$

and thus obtain the following result, relating the actions of Hecke operators to the fractional Rademacher sums.

Theorem 5.5.6. *Let $n \in \mathbb{Z}^+$ be square-free and let $\Gamma = G(\mathbb{Z})$. Then we have*

$$n(\hat{T}(n)Q_{\llbracket \Gamma}^{(1)})(z) = \sum_{h \parallel n} Q_{\llbracket^h \Gamma}^{(n/h^2)}(z) \quad (5.5.22)$$

where the sum is over exact divisors of n .

It is interesting to compare the result (5.5.22) of Theorem 5.5.6 with the identity

$$n(\hat{T}(n)Q_{\llbracket \Gamma}^{(1)})(z) = Q_{\llbracket \Gamma}^{(n)}(z), \quad (5.5.23)$$

which holds since both sides are holomorphic on $\mathbb{H}$, and have the same singular terms in their Fourier expansion at $\Gamma \cdot \infty$. This proves that the Rademacher sum $Q_U^{(n/h^2)}(z)$ vanishes for $U = \llbracket^h \Gamma$ whenever h is an exact divisor of n , and thus recovers a special case of the vanishing result of Theorem 5.4.3.

6 Moonshine

Monstrous moonshine associates a group Γ_g commensurable with $G(\mathbb{Z})$ and having width one at infinity to each (conjugacy class of) element(s) g in the Monster group $\mathbb{M}$. In the article [CMS04] the set of groups $\{\Gamma_g \mid g \in \mathbb{M}\}$ is characterized in purely group theoretic terms. In this section we describe a reformulation of this characterization in terms of normalized Rademacher sums, and a certain family of moduli problems for solid tori with conformal structure on the boundary.

6.1 Genera

The main theorem of [CMS04] gives four conditions which, taken together, characterize the groups Γ_g for $g \in \mathbb{M}$. The first of these conditions is the requirement that a group Γ have genus zero. Our first result in this section is a reformulation of the genus zero condition in terms of Rademacher sums.

Theorem 6.1.1. *Let Γ be a group commensurable with $G(\mathbb{Z})$ that has width one at infinity and let $\mathfrak{p}$ be a cusp for Γ . Then the normalized Rademacher sum $T_{\Gamma, \mathfrak{p}}^{(1)}(z)$ defines a Γ -invariant*

function on $\mathbb{H}$ if and only if Γ has genus zero.

Proof. Let Γ be as in the statement of the theorem and let $\mathbf{p}, \mathbf{q} \in \mathbf{P}_\Gamma$ be cusps of Γ . By Theorem 4.4.6 we have

$$T_{\Gamma, \mathbf{p}|\mathbf{q}}^{(1)}(z) = \delta_{\Gamma, \mathbf{p}|\mathbf{q}} e(-z) + F_{\Gamma, \mathbf{p}|\mathbf{q}}^{(1)}(z)_{\text{van}}. \quad (6.1.1)$$

Taking $\mathbf{q} = \Gamma \cdot \infty$ in (6.1.1) we see that $T_{\Gamma, \mathbf{p}}^{(1)}(z)$ is holomorphic on $\mathbb{H}$, and letting $\mathbf{q}$ range over $\mathbf{P}_\Gamma$ we see from (6.1.1) that the only pole of $T_{\Gamma, \mathbf{p}}^{(1)}(z)$ is a simple pole at the cusp $\mathbf{p}$. We see then that if the function $T_{\Gamma, \mathbf{p}}^{(1)}(z)$ is Γ -invariant then it defines a morphism ϕ say, of Riemann surfaces $\phi : \mathbf{X}_\Gamma \rightarrow \hat{\mathbb{C}}$ (cf. §2.5) which has degree one since the preimage of $\infty \in \hat{\mathbb{C}}$ under ϕ is the single point $\mathbf{p} \in \mathbf{X}_\Gamma$. We conclude that ϕ is an isomorphism, so that Γ indeed has genus zero.

Conversely, if Γ has genus zero then, since the space $S_1(\Gamma)$ of cusp forms of weight 2 for Γ is isomorphic to the space of holomorphic differentials on $\mathbf{X}_\Gamma$, we have $S_1(\Gamma) = \{0\}$, so that any automorphic integral of weight 0 for Γ is in fact a Γ -invariant function, by Corollary 4.5.5. The modified Rademacher sum $Q_{\Gamma, \mathbf{p}}^{(1)}(z)$ is an automorphic integral for Γ by Theorem 4.5.3. The normalized Rademacher sum $T_{\Gamma, \mathbf{p}}^{(1)}(z)$ differs from $Q_{\Gamma, \mathbf{p}}^{(1)}(z)$ by a constant function by Proposition 4.4.11, and thus $T_{\Gamma, \mathbf{p}}^{(1)}(z)$ is Γ -invariant whenever $Q_{\Gamma, \mathbf{p}}^{(1)}(z)$ is. We conclude that $T_{\Gamma, \mathbf{p}}^{(1)}(z)$ is Γ -invariant in case Γ has genus zero. This completes the proof. $\square$

From the first part of the proof of Theorem 6.1.1 we see that the normalized Rademacher sum $T_{\Gamma, \mathbf{p}}^{(1)}(z)$ associated to Γ at a cusp $\mathbf{p}$ defines an isomorphism $\mathbf{X}_\Gamma \rightarrow \hat{\mathbb{C}}$ mapping $\mathbf{p}$ to $\infty \in \hat{\mathbb{C}}$ in case Γ is a group of genus zero.

Theorem 6.1.2. *Let Γ be a group commensurable with $G(\mathbb{Z})$ that has width one at infinity and let $\mathbf{p} \in \mathbf{P}_\Gamma$ be a cusp of Γ . If Γ has genus zero then the normalized Rademacher sum $T_{\Gamma, \mathbf{p}}^{(1)}(z)$ associated to Γ at the cusp $\mathbf{p}$ induces an isomorphism $\mathbf{X}_\Gamma \rightarrow \hat{\mathbb{C}}$ mapping $\mathbf{p}$ to ∞ .*

Applying Theorem 6.1.2 with $\mathbf{p} = \Gamma \cdot \infty$ the infinite cusp we see that the normalized Rademacher sum $T_\Gamma^{(1)}(z)$ induces an isomorphism $\mathbf{X}_\Gamma \rightarrow \hat{\mathbb{C}}$ mapping the infinite cusp to $\infty \in \hat{\mathbb{C}}$. Applying Proposition 4.4.9 with $\mathbf{p} = \mathbf{q} = \Gamma \cdot \infty$ we see that the Fourier expansion of $T_\Gamma^{(1)}(z)$ has vanishing constant term. This shows that the normalized Rademacher sum $T_\Gamma^{(1)}(z)$ associated to Γ at the infinite cusp is the normalized hauptmodul for Γ (cf. §2.5) whenever this statement makes sense; i.e. whenever Γ has genus zero.

Theorem 6.1.3. *Let Γ be a group commensurable with $G(\mathbb{Z})$ that has width one at infinity. If Γ has genus zero then the normalized Rademacher sum $T_\Gamma^{(1)}(z)$ is the normalized hauptmodul for Γ .*

From the second part of the proof of Theorem 6.1.1 we see that if Γ has genus zero then not only the first order but in fact all the higher order normalized Rademacher sums $T_{\Gamma, \mathbf{p}}^{(m)}(z)$

associated to Γ at an arbitrary cusp $\mathfrak{p} \in \mathcal{P}_\Gamma$ are Γ -invariant.

Theorem 6.1.4. *Let Γ be a subgroup of $G(\mathbb{Q})$ that is commensurable with $G(\mathbb{Z})$ and has width one at infinity and let $\mathfrak{p} \in \mathcal{P}_\Gamma$ be a cusp of Γ . Let $m \in \mathbb{Z}^+$. If Γ has genus zero then the normalized Rademacher sum $T_{\Gamma, \mathfrak{p}}^{(m)}(z)$ is Γ -invariant.*

6.2 Moduli

The second condition of the main theorem of [CMS04] is that a group Γ be of $n||h$ -type (cf. §2.5). In this section we provide a geometric description of the groups of $n||h$ -type by furnishing a family of moduli problems for which the corresponding moduli spaces may be realized as quotients of the form $\mathcal{Y}_\Gamma = \Gamma \backslash \mathbb{H}$ where Γ is a group of $n||h$ -type.

Consider pairs (E, C) where E is an elliptic curve over $\mathbb{C}$ and C is an oriented subgroup of E isomorphic to S^1 . We call such a pair a *solid torus*. For (E, C) a solid torus, we call E the *underlying elliptic curve*, and we call C the *underlying primitive cycle*. Note that for (E, C) a solid torus, the group C determines a primitive element of the first homology group of (the smooth real surface underlying) the elliptic curve E .

Given $z \in \mathbb{H}$ let us write Λ_z for the lattice $\mathbb{Z}z + \mathbb{Z} \subset \mathbb{C}$. Observe that the lattices Λ_z and $\Lambda_{z'}$ coincide if and only if $z - z' \in \mathbb{Z}$, so the assignment $z \mapsto \Lambda_z$ descends naturally to the orbit space $B(\mathbb{Z}) \backslash \mathbb{H}$. If we agree to write $\llbracket z$ as a shorthand for the $B(\mathbb{Z})$ orbit determined by $z \in \mathbb{H}$, then we may unambiguously write $\Lambda_{\llbracket z}$ for the lattice $\mathbb{Z}z + \mathbb{Z}$. We write $E_{\llbracket z}$ for the corresponding elliptic curve $E_{\llbracket z} = \mathbb{C}/\Lambda_{\llbracket z} = \mathbb{C}/\mathbb{Z}z + \mathbb{Z}$. Then the pair $(E_{\llbracket z}, C_{\llbracket z})$ is a solid torus when we take the subgroup $C_{\llbracket z}$ to be the image of $\mathbb{R}$ in $E_{\llbracket z}$ under the composition $\mathbb{R} \rightarrow \mathbb{C} \rightarrow E_{\llbracket z}$. Observe that for any solid torus (E, C) there exists $z \in \mathbb{H}$ and a unique isomorphism of elliptic curves $E \rightarrow E_{\llbracket z}$ inducing an orientation preserving isomorphism $C \rightarrow C_{\llbracket z}$. This observation is the content of the following proposition.

Proposition 6.2.1. *The quotient space $B(\mathbb{Z}) \backslash \mathbb{H}$ is a fine moduli space for solid tori.*

For (E, C) a solid torus and $n \in \mathbb{Z}^+$ we write $E[n]$ for the group of n -division points of E , and we write $C[n]$ for the intersection $C \cap E[n]$. Since C is oriented, each group $C[n]$ comes equipped with a distinguished generator; viz., the point corresponding to $1/n + \Lambda_{\llbracket z}$ under an isomorphism $E \rightarrow E_{\llbracket z}$ inducing an orientation preserving isomorphism $C \rightarrow C_{\llbracket z}$. If K is a subgroup of $E[n]$ for some n then the quotient $\bar{E} = E/K$ is again an elliptic curve, and the image $\bar{C}$ say, of C under the natural map $E \rightarrow \bar{E}$ defines a primitive cycle of $\bar{E}$, so that the pair $(\bar{E}, \bar{C})$ is again a solid torus. We call this the *quotient of (E, C) by K* and denote it $(E, C)/K$.

For $n \in \mathbb{Z}^+$ define an *n -isogeny of solid tori* $(E', C') \rightarrow (E, C)$ to be an isogeny $E' \rightarrow E$ of elliptic curves that maps $C'[n]$ to a subgroup of $C[n]$. Then a 1-isogeny of solid tori is just an isogeny of the underlying elliptic curves. We define an *isogeny of solid tori* $(E', C') \rightarrow (E, C)$ to be an isogeny $E' \rightarrow E$ of elliptic curves that restricts to an orientation preserving map $C' \rightarrow C$.

on the underlying primitive cycles. For (E, C) a solid torus and $n \in \mathbb{Z}^+$, we may now interpret the canonical map $E \rightarrow E/C[n]$ as defining an isogeny $(E, C) \rightarrow (E, C)/C[n]$ of solid tori. We call this isogeny the *canonical n -fold quotient* of (E, C) .

Observe that the elliptic curve $E/C[n]$ underlying the quotient $(E, C)/C[n]$ both receives a natural map from E and maps naturally to E , for the quotient $E/E[n]$ is naturally isomorphic to E , and for the map $E/C[n] \rightarrow E$ we may take the composition $E/C[n] \rightarrow E/E[n] \xrightarrow{\sim} E$ where the first map is the natural projection and the second map is the isomorphism just mentioned. This map $E/C[n] \rightarrow E$ defines an isogeny $(E, C)/C[n] \rightarrow (E, C)$ of solid tori; indeed, it restricts to an isomorphism on the underlying primitive cycles. We call the isogeny $(E, C)/C[n] \rightarrow (E, C)$ of solid tori the *canonical n -fold cover* of (E, C) .

Say an n -isogeny of solid tori $(E', C') \rightarrow (E, C)$, for $n \in \mathbb{Z}^+$, is an *n -isomorphism of solid tori* in case it admits an inverse n -isogeny; i.e. an isogeny $E \rightarrow E'$ mapping $C[n]$ to a subgroup of $C'[n]$ such that the compositions $E' \rightarrow E \rightarrow E'$ and $E \rightarrow E' \rightarrow E$ are the identity maps on E' and E , respectively. Simply put, an n -isomorphism of solid tori $(E', C') \rightarrow (E, C)$ is an isomorphism of the underlying elliptic curves $E' \rightarrow E$ that induces an isomorphism of groups $C'[n] \rightarrow C[n]$. An *isomorphism of solid tori* $(E', C') \rightarrow (E, C)$ is an isomorphism $E' \rightarrow E$ of elliptic curves that restricts to an orientation preserving isomorphism $C' \rightarrow C$ on the underlying primitive cycles.

Suppose (E, C) and (E', C') are solid tori. For n a positive integer and e an exact divisor of n , say (E, C) and (E', C') are $n + e$ -related if there is an n -isogeny $(E', C') \rightarrow (E, C)$ that induces n/e -isomorphisms $(E', C')/C'[e] \rightarrow (E, C)$ and $(E', C') \rightarrow (E, C)/C[e]$. More precisely, we require that the n -isogeny $(E', C') \rightarrow (E, C)$ factor through the canonical e -fold maps $(E', C') \rightarrow (E', C')/C'[e]$ and $(E, C)/C[e] \rightarrow (E, C)$, defining n/e -isomorphisms of the specified type.

$$\begin{array}{ccc}
 (E', C') & \xrightarrow[n/e]{\sim} & (E, C)/C[e] \\
 \downarrow & & \downarrow \\
 (E', C')/C'[e] & \xrightarrow[n/e]{\sim} & (E, C)
 \end{array} \tag{6.2.1}$$

Then solid tori (E, C) and (E', C') are $n + 1$ -related just if they are n -isomorphic, which is the case just if there is an isomorphism of elliptic curves $E' \rightarrow E$ mapping $C'[n]$ onto $C[n]$. In particular, the notion of being $n + 1$ -related is an equivalence relation on solid tori. For S a subset of the set of exact divisors of n , say solid tori (E, C) and (E', C') are $n + S$ -related if (E, C) and (E', C') are $n + e$ -related for some $e \in S$.

Recall from §2.2 that the set of exact divisors of n , denoted $\text{Ex}(n)$, admits a naturally defined group structure.

Lemma 6.2.2. *Let $n \in \mathbb{Z}^+$ and let $S \subset \text{Ex}(n)$. Then the $n + S$ -relation is an equivalence relation on solid tori if and only if S is a subgroup of $\text{Ex}(n)$.*

In light of Lemma 6.2.2, we say that solid tori (E, C) and (E', C') are $n + S$ -equivalent if they are $n + S$ -related and S is a subgroup of the group of exact divisors of n .

In a slight modification of the notation of [CN79], we write $\Gamma_0(n) + S$ for the group formed by taking the union of the Hecke congruence group $\Gamma_0(n)$ and the Atkin–Lehner involutions $W_e(n)$ for e in S , when S is a subgroup of $\text{Ex}(n)$.

$$W_e(n) = \left\{ \begin{bmatrix} ae & b \\ cn & de \end{bmatrix} \in G(\mathbb{Q}) \mid a, b, c, d \in \mathbb{Z}, ade - bcn/e = 1 \right\} \quad (6.2.2)$$

Let n be a positive integer and let h be a divisor of n . Then an n -isomorphism $\phi : (E', C') \rightarrow (E, C)$ naturally determines an nh -isogeny $\hat{\phi} : (E', C') \rightarrow (E, C)$ which factors through an n/h -isomorphism $\bar{\phi} : (E', C')/C'[h] \rightarrow (E, C)/C[h]$ on the canonical h -fold quotients, via the canonical maps $(E', C') \rightarrow (E', C')/C'[h]$ and $(E, C)/C[h] \rightarrow (E, C)$, as depicted in (6.2.3).

$$\begin{array}{ccc} (E', C') & \xrightarrow[nh]{\hat{\phi}} & (E, C) \\ \searrow & & \nearrow \\ (E', C')/C'[h] & \xrightarrow[n/h]{\bar{\phi}} & (E, C)/C[h] \end{array} \quad (6.2.3)$$

We may take this nh -isogeny $\hat{\phi}$ to be the composition $E' \rightarrow E \rightarrow E/E[h] \rightarrow E$ where the first map is the given map ϕ , the second map is the natural projection, and the third map is the natural isomorphism. Since ϕ is an n -isomorphism the composition $E' \rightarrow E \rightarrow E/E[h]$ factors through the natural projection $E/C[h] \rightarrow E/E[h]$, and the kernel of the induced map $E' \rightarrow E/C[h]$ is just $C'[h]$, and so we arrive at an isomorphism $E'/C'[h] \rightarrow E/C[h]$ of elliptic curves, which defines the required n/h -isomorphism $\bar{\phi} : (E', C')/C'[h] \rightarrow (E, C)/C[h]$ of the corresponding solid tori.

$$\begin{array}{ccccccc} (E', C') & \xrightarrow[n]{\phi} & (E, C) & \longrightarrow & (E, C)/E[h] & \xrightarrow{\sim} & (E, C) \\ & \searrow & & & \nearrow & & \nearrow \\ & & (E', C')/C'[h] & \xrightarrow[n/h]{\bar{\phi}} & (E, C)/C[h] & & \end{array} \quad (6.2.4)$$

More generally, and by the same argument, an n -isogeny $(E', C') \rightarrow (E, C)$ naturally defines an nh -isogeny $(E', C') \rightarrow (E, C)$ which factors through an n/h -isogeny $(E', C')/C'[h] \rightarrow$

$(E, C)/C[h]$, via the canonical maps $(E', C') \rightarrow (E', C')/C'[h]$ and $(E, C)/C[h] \rightarrow (E, C)$.

Not every n/h -isogeny $\bar{\phi} : (E', C')/C'[h] \rightarrow (E, C)/C[h]$ arises in this way from an n -isogeny $\phi : (E', C') \rightarrow (E, C)$, but, given an n/h -isogeny $\bar{\phi} : (E', C')/C'[h] \rightarrow (E, C)/C[h]$, we always have the nh -isogeny $\hat{\phi} : (E', C') \rightarrow (E, C)$ obtained as the composition

$$(E', C') \rightarrow (E', C')/C'[h] \rightarrow (E, C)/C[h] \rightarrow (E, C) \quad (6.2.5)$$

where the first map is the canonical h -fold quotient, and the third map is the canonical h -fold cover. We call this composition $\hat{\phi}$ the *nh -isogeny induced from $\bar{\phi}$* .

For $n \in \mathbb{Z}^+$ and h a divisor of n say solid tori (E, C) and (E', C') are $n|h$ -related if the canonical h -fold quotients $(E, C)/C[h]$ and $(E', C')/C'[h]$ are n/h -isomorphic. For e an exact divisor of n/h , say (E, C) and (E', C') are $n|h + e$ -related if the quotients $(E, C)/C[h]$ and $(E', C')/C'[h]$ are $n/h + e$ -related. For S a subset of $\text{Ex}(n/h)$ say (E, C) and (E', C') are $n|h + S$ -related if they are $n|h + e$ -related for some $e \in S$.

Lemma 6.2.3. *Let $n \in \mathbb{Z}^+$, let h be a divisor of n , and let $S \subset \text{Ex}(n/h)$. Then the $n|h + S$ -relation is an equivalence relation on solid tori if and only if S is a subgroup of $\text{Ex}(n/h)$.*

Let N be a positive integer, set $h = ((N, 24))$ (cf. §2.2), and set $n = N/h$. Then for S a subgroup of the group of exact divisors of n/h we arrive at the notion of $n||h + S$ -equivalence of solid tori in the following way. Suppose (E, C) and (E', C') are $\Gamma_0(n|h) + S$ -equivalent. Then we have isomorphisms $(E, C) \rightarrow (E|_z, C|_z)$ and $(E', C') \rightarrow (E'|_{z'}, C'|_{z'})$ for some $z, z' \in \mathbb{H}$. Let S' be the image of S under the natural injection $\text{Ex}(n/h) \rightarrow \text{Ex}(N)$ (cf. §2.2), and define elements x, y and $w_{e'}$ for $e' \in S'$ by setting

$$x = \begin{bmatrix} 1 & 1/h \\ 0 & 1 \end{bmatrix}, \quad y = \begin{bmatrix} 1 & 0 \\ n & 1 \end{bmatrix}, \quad w_{e'} = \begin{bmatrix} ae' & b \\ cN & de' \end{bmatrix}, \quad (6.2.6)$$

where the a, b, c and d in the definition of $w_{e'}$ are arbitrary integers for which we have $ade' - bcN/e' = 1$. Then there is a sequence $(\gamma_0, \gamma_1, \dots, \gamma_k)$ with $\gamma_0 \in \Gamma_0(N)$ and $\gamma_i \in \{x, y, w_{e'} \mid e' \in S'\}$ for $1 \leq i \leq k$ for which we have $z' = \gamma_k \cdots \gamma_1 \gamma_0 \cdot z$. We say now that (E, C) and (E', C') are $\Gamma_0(n||h) + S$ -equivalent in the case that the sum $\sum_{i=1}^k \lambda(\gamma_i)$ vanishes in $\mathbb{Z}/h$, where $\lambda(x) = 1$, and $\lambda(w_{e'}) = 0$ for all $e' \in S'$, and $\lambda(y)$ is -1 or 1 according as $\Gamma_0(n|h) + S$ is Fricke or not.

Theorem 6.2.4. *Let $N \in \mathbb{Z}^+$, set $h = ((N, 24))$ and set $n = N/h$. Let S be a subgroup of $\text{Ex}(n/h)$, and set $\Gamma = \Gamma_0(n||h) + S$. Then the quotient $\Gamma \backslash \mathbb{H}$ is a moduli space for solid tori under $n||h + S$ -equivalence.*

6.3 Exponents

The third of the four conditions of the main theorem of [CMS04] states that if Γ is of the form $\Gamma_0(n|h)+S$ then the quotient group $\Gamma/\Gamma_0(nh)$ should be a group of exponent two. In the present section we investigate the relationship between this exponent two condition and properties of scaling cosets.

Lemma 6.3.1. *Suppose that Δ is a group commensurable with $G(\mathbb{Z})$ that has width one at infinity and let $r \in P_\Delta$ be a cusp of Δ . Let $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$ and let $m \in \mathbb{Z}^+$. If Σ_r is a scaling coset for Δ at r with the property that $\Sigma_r^2 = \Delta$ then*

$$Q_{\Delta,r}^{\kappa(m)}(z) = Q_{\Delta|r}^{\kappa(m)}(z), \quad (6.3.1)$$

and Σ_r is contained in the normalizer of Δ .

Proof. Set $U = \llbracket \Sigma_r^{-1}$ and set $U' = \llbracket \Sigma_r$. Then, under the assumption that Δ has width one at infinity, we have $Q_{\Delta,r}^{\kappa(m)}(z) = Q_U^{\kappa(m)}(z)$ and $Q_{\Delta|r}^{\kappa(m)}(z) = Q_{U'}^{\kappa(m)}(z)$. Let $\sigma \in \Sigma_r$. Then $\Sigma_r^2 = \Delta$ implies $\sigma\Delta\sigma = \Delta$, which is equivalent to the identity $\Delta\sigma = \sigma^{-1}\Delta$. We conclude that $\Sigma_r^{-1} = \Sigma_r$, so that $U = U'$, so that $Q_{\Delta,r}^{\kappa(m)}(z) = Q_{\Delta|r}^{\kappa(m)}(z)$. The identity $\sigma\Delta\sigma = \Delta$ also implies $\sigma^2 \in \Delta$, so that $\sigma\Delta\sigma = \sigma^{-1}\Delta\sigma$. We conclude that σ normalizes Δ , so that $\Sigma_r = \Delta\sigma \in \Delta \backslash N(\Delta)$. This completes the proof. $\square$

Lemma 6.3.1 states that if a scaling coset at a cusp r of a group Δ can be chosen so that its square is the trivial coset then we have a kind of commutativity for the Rademacher sums associated to Δ ; viz. the expansion at infinity of the modified Rademacher sum associated to Δ and the cusp r coincides with the expansion at r of the modified Rademacher sum associated to Δ and the cusp at infinity.

We also have the following converse to Lemma 6.3.1.

Proposition 6.3.2. *Suppose that Δ is a group commensurable with $G(\mathbb{Z})$ that has width one at infinity and let $r \in P_\Delta$ be a cusp of Δ . Let $\kappa \in \mathbb{Z}$ such that $1 - \kappa \in \mathbb{Z}^+$ and let $m \in \mathbb{Z}^+$. If Σ_r is a scaling coset for Δ at r that is contained in the normalizer of Δ and*

$$Q_{\Delta,r}^{\kappa(m)}(z) = Q_{\Delta|r}^{\kappa(m)}(z) \quad (6.3.2)$$

then $\Sigma_r^2 = \Delta$.

Proof. As in the proof of Lemma 6.3.1 we set $U = \llbracket \Sigma_r^{-1}$ and $U' = \llbracket \Sigma_r$. Then since Δ is assumed to have width one at infinity we have $Q_{\Delta,r}^{\kappa(m)}(z) = Q_U^{\kappa(m)}(z)$ and $Q_{\Delta|r}^{\kappa(m)}(z) = Q_{U'}^{\kappa(m)}(z)$. Let $\sigma \in \Sigma_r$. Then we have $U = \llbracket \sigma^{-1}\Delta$ and $U' = \llbracket \Delta\sigma = \llbracket \sigma\Delta$. Set $r' = \Delta\sigma^{-1} \cdot \infty$. Then

$Q_{U'}^{\kappa(m)}(z) = Q_{\Delta, r'}^{\kappa(m)}(z)$, so the coincidence $Q_U^{\kappa(m)}(z) = Q_{U'}^{\kappa(m)}(z)$ implies that $r = r'$. This in turn implies $\sigma^2 \in \Delta$, so that $\Sigma_r^2 = \Delta$, as we required to show. $\square$

6.4 Cusps

The fourth condition of the main theorem of [CMS04] is the following. Supposing that a group Γ is of the form $\Gamma = \Gamma_0(n|h) + S$ for some subgroup S of the group of exact divisors of n/h , for each $p \in \hat{\mathbb{Q}}$ there should exist an element $\tilde{\sigma}_p \in G(\mathbb{R})$ such that

$$\infty = \tilde{\sigma}_p \cdot p, \quad (\tilde{\sigma}_p \Gamma \tilde{\sigma}_p^{-1})_\infty = B(\mathbb{Z}), \quad \tilde{\sigma}_p \Gamma \tilde{\sigma}_p^{-1} \supset \Gamma_0(nh). \quad (6.4.1)$$

Observe that if we set $\sigma_p = \tilde{\sigma}_p^{-1}$, then the conditions of (6.4.1) translate into

$$\sigma_p \cdot \infty = p, \quad (\sigma_p^{-1} \Gamma \sigma_p)_\infty = B(\mathbb{Z}), \quad \sigma_p^{-1} \Gamma \sigma_p \supset \Gamma_0(nh), \quad (6.4.2)$$

the first two of which are just the conditions (2.6.1) of Lemma 2.6.1 except that we allow σ_p to lie in $G(\mathbb{R})$ in (6.4.2) but insist that σ_p belong to $G(\mathbb{Q})$ in (2.6.1). Actually, an element $\sigma_p \in G(\mathbb{R})$ satisfying (6.4.2) must lie in $G(\mathbb{Q})$, as the following result demonstrates.

Lemma 6.4.1. *Let Δ and Γ be groups commensurable with $G(\mathbb{Z})$ and suppose that Δ has width one at infinity. Let $p \in \hat{\mathbb{Q}}$ and suppose that $\sigma \in G(\mathbb{R})$ satisfies $\sigma \cdot \infty = p$ and $\sigma^{-1} \Gamma \sigma \supset \Delta$. Then $\sigma \in G(\mathbb{Q})$.*

Proof. Suppose that $\sigma \in G(\mathbb{R})$ satisfies the hypotheses of the lemma. Then $(\sigma^{-1} \Gamma \sigma)_\infty$ is an infinite cyclic group containing Δ_∞ and $\Delta_\infty = B(\mathbb{Z})$ since Δ has width one at infinity. It must be then that $(\sigma^{-1} \Gamma \sigma)_\infty$ is generated by $T^{1/n}$ for some $n \in \mathbb{Z}^+$, so that $([n]\sigma^{-1} \Gamma \sigma [1/n])_\infty = B(\mathbb{Z})$. According to Lemma 2.6.1 there exists $\sigma' \in G(\mathbb{Q})$ such that $\sigma' \cdot \infty = p$ and $((\sigma')^{-1} \Gamma \sigma')_\infty = B(\mathbb{Z})$. Now $[n]\sigma^{-1} \sigma'$ fixes ∞ and so lies in $B(\mathbb{R})$. Since σ' and $\sigma[1/n]$ both conjugate Γ to a group with width one at infinity we must in fact have $[n]\sigma^{-1} \sigma' \in B_u(\mathbb{R})$, where $B_u(\mathbb{R})$ consists of all the elements T^α (cf. (2.3.4)) with $\alpha \in \mathbb{R}$. We conclude that $\sigma = \sigma'' T^\alpha$ for some $\sigma'' \in G(\mathbb{Q})$ and $\alpha \in \mathbb{R}$. Let $\gamma \in \Gamma$ such that $\sigma^{-1} \gamma \sigma \in \Delta^\times$, so that $\sigma^{-1} \gamma \sigma \cdot \infty = q$ for some $q \in \mathbb{Q}$. Then $q = q'' - \alpha$ for $q'' = (\sigma'')^{-1} \gamma \sigma'' \cdot \infty$. Since both q and q'' lie in $\mathbb{Q}$ we conclude that α also lies in $\mathbb{Q}$. Then the identity $\sigma = \sigma'' T^\alpha$ implies that $\sigma \in G(\mathbb{Q})$, as we required to show. $\square$

Recall from §2.6 that for Γ a group commensurable with $G(\mathbb{Z})$, for $\mathfrak{p} \in \mathcal{P}_\Gamma$ a cusp of Γ , and for $\Sigma_\mathfrak{p} \in \Gamma \backslash G(\mathbb{Q})$ a scaling coset for Γ at $\mathfrak{p}$, we write $\Gamma^\mathfrak{p}$ as a shorthand for the group $\Sigma_\mathfrak{p}^{-1} \Sigma_\mathfrak{p}$. In light of Lemma 6.4.1 we may reformulate the fourth condition of the main theorem of [CMS04] as follows.

Lemma 6.4.2. *Let Γ be a group of $n|h$ -type, so that $\Gamma = \Gamma_0(n|h) + S$ for some positive integers n and h , and some subgroup S of the group of exact divisors of n/h . Then the fourth condition*

of the main theorem of [CMS04] is satisfied if and only if for each cusp $\mathfrak{p} \in \mathcal{P}_\Gamma$ of Γ there exists a scaling coset $\Sigma_{\mathfrak{p}} \in \Gamma \backslash G(\mathbb{Q})$ for Γ at $\mathfrak{p}$ such that $\Gamma^{\mathfrak{p}}$ contains $\Gamma_0(nh)$.

We conclude this section by relating the reformulation of Lemma 6.4.2 to the normalized Rademacher sums associated to genus zero groups of $n||h$ -type.

Proposition 6.4.3. *Let Γ be a group of $n||h$ -type, so that $\Gamma = \Gamma_0(n||h) + S$ for some positive integers n and h , and some subgroup S of the group of exact divisors of n/h , and suppose that Γ has genus zero. Let $\mathfrak{p}, \mathfrak{q} \in \mathcal{P}_\Gamma$ be cusps of Γ and let $\Sigma_{\mathfrak{p}}$ and $\Sigma_{\mathfrak{q}}$ be scaling cosets for Γ at $\mathfrak{p}$ and $\mathfrak{q}$, respectively. Then the normalized Rademacher sum $T_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(1)}(z)$ is $\Gamma_0(nh)$ -invariant if and only if $\Gamma^{\mathfrak{q}}$ contains $\Gamma_0(nh)$.*

Proof. Let us set $\Delta = \Gamma_0(nh)$. Observe that if we set $\mathfrak{p}^{\mathfrak{q}} = \Sigma_{\mathfrak{q}}^{-1} \Sigma_{\mathfrak{p}} \cdot \infty$ then $\mathfrak{p}^{\mathfrak{q}}$ is a cusp of $\Gamma^{\mathfrak{q}} = \Sigma_{\mathfrak{q}}^{-1} \Sigma_{\mathfrak{p}}$, and $\Sigma_{\mathfrak{q}}^{-1} \Sigma_{\mathfrak{p}}$ is a scaling coset for $\Gamma^{\mathfrak{q}}$ at $\mathfrak{p}^{\mathfrak{q}}$, so the function $T_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(1)}(z)$ may be identified with the normalized Rademacher sum $T_{\Gamma^{\mathfrak{q}}, \mathfrak{p}^{\mathfrak{q}}}^{(1)}(z)$ associated to $\Gamma^{\mathfrak{q}}$ at the cusp $\mathfrak{p}^{\mathfrak{q}}$, for $\Sigma_{\mathfrak{p}^{\mathfrak{q}}} = \Sigma_{\mathfrak{q}}^{-1} \Sigma_{\mathfrak{p}}$. Under the assumption that Γ has genus zero, $\Gamma^{\mathfrak{q}}$ also has genus zero, and has width one at infinity by the defining properties (cf. Lemma 2.6.2) of $\Sigma_{\mathfrak{q}}$. By Theorem 6.1.3 then the normalized Rademacher sum $T_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(1)}(z)$ is, up to a constant function, the expansion at $\mathfrak{p}^{\mathfrak{q}}$ of the normalized hauptmodul of the group $\Gamma^{\mathfrak{q}}$. In particular, $T_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(1)}(z)$ defines an isomorphism of Riemann surfaces $X_{\Gamma^{\mathfrak{q}}} \rightarrow \hat{\mathbb{C}}$, and so we have $T_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(1)}(\gamma \cdot z) = T_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(1)}(z)$ for all $z \in \mathbb{H}$ if and only if $\gamma \in \Gamma^{\mathfrak{q}}$. It follows then that $T_{\Gamma, \mathfrak{p}|\mathfrak{q}}^{(1)}(z)$ is Δ -invariant if and only if Δ is contained in $\Gamma^{\mathfrak{q}}$, which is what we required to show. $\square$

6.5 Rademacher sums and the Monster

We will now use Rademacher sums to reformulate the characterization of the groups of monstrous moonshine due to Conway–McKay–Sebbar (cf. [CMS04]).

Theorem 6.5.1. *Let Γ be a subgroup of $G(\mathbb{R})$. Then we have $\Gamma = \Gamma_g$ for some $g \in \mathbb{M}$ if and only if the following conditions are satisfied:*

- Γ is the group defining $n||h + S$ -equivalence of solid tori for some positive integers n and h , and some subgroup $S < \text{Ex}(n/h)$;
- the normalized Rademacher sum $T_{\Gamma}^{(1)}(z)$ is Γ -invariant;
- there exists a system $\{\Sigma_{\mathfrak{p}} \mid \mathfrak{p} \in \mathcal{P}_\Gamma\}$ of scaling cosets for Γ such that $T_{\Gamma|\mathfrak{p}}^{(1)}(z)$ is Δ -invariant for every cusp $\mathfrak{p} \in \mathcal{P}_\Gamma$ of Γ , where $\Delta = \Gamma_0(nh)$;
- for every cusp $\mathfrak{r} \in \mathcal{P}_\Delta$ of $\Delta = \Gamma_0(nh)$ that is contained in $\Gamma \cdot \infty$ we have $T_{\Delta, \mathfrak{r}}^{(1)}(z) = T_{\Delta|\mathfrak{r}}^{(1)}(z)$ when the scaling coset $\Sigma_{\mathfrak{r}}$ for Δ at $\mathfrak{r}$ is taken to lie in $\Delta \backslash \Gamma$.

Proof. According to the main theorem of [CMS04] it suffices to show that a group Γ satisfies each of the four conditions of Theorem 6.5.1 if and only if it satisfies the four conditions of the

main theorem of [CMS04]. As a first step in establishing this equivalence, observe that the first condition of Theorem 6.5.1 is exactly the same as the second condition of the main theorem of [CMS04].

Suppose that Γ satisfies the four conditions of the main theorem of [CMS04]. Then, in particular, it satisfies the first condition of Theorem 6.5.1, and has genus zero. We may suppose then that $\Gamma = \Gamma_0(n|h) + S$ for some $n, h \in \mathbb{Z}^+$, and $S < \text{Ex}(n/h)$, so that Γ is commensurable with $G(\mathbb{Z})$, has width one at infinity, and contains and normalizes $\Delta = \Gamma_0(nh)$. According to Theorem 6.1.1 the normalized Rademacher sum $T_\Gamma^{(1)}(z)$ is Γ -invariant, so the second condition of Theorem 6.5.1 is satisfied. According to Lemma 6.4.2 there exists a system $\{\Sigma_p \mid p \in P_\Gamma\}$ of scaling cosets for Γ such that Γ^p contains Δ for each cusp $p \in P_\Gamma$. Applying Proposition 6.4.3 we see that the normalized Rademacher sum $T_{\Gamma|p}^{(1)}(z)$ is Δ -invariant when the scaling coset Σ_p of the above scaling coset system is chosen for the definition of $T_{\Gamma|p}^{(1)}(z)$. This confirms that the third condition of Theorem 6.5.1 is satisfied. The fourth condition of the main theorem of [CMS04] is that the quotient Γ/Δ have exponent two. Let $r \in P_\Delta$ be a cusp of Δ that is contained in $\Gamma \cdot \infty$, and let Σ_r be the unique right coset of Δ in Γ such that $r = \Sigma_r \cdot \infty$. Then the exponent two condition implies $\Sigma_r^2 = \Delta$, so that the fourth condition of Theorem 6.5.1 follows from an application of Lemma 6.3.1.

Suppose now that Γ satisfies the four conditions of Theorem 6.5.1. Then Γ satisfies the second condition of [CMS04], and so we have $\Gamma = \Gamma_0(n|h) + S$ for some $n, h \in \mathbb{Z}^+$, and $S < \text{Ex}(n/h)$, and Γ is a group commensurable with $G(\mathbb{Z})$ that has width one at infinity. Applying Theorem 6.1.1 to the second condition of Theorem 6.5.1 we conclude that Γ has genus zero, and so satisfies the first condition of [CMS04]. Set $\Delta = \Gamma_0(nh)$, so that Γ contains and normalizes Δ . Applying Proposition 6.4.3 to the third condition of Theorem 6.5.1 we see that scaling cosets $\{\Sigma_p \mid p \in P_\Gamma\}$ can be chosen for Γ so that Γ^p contains Δ for each cusp $p \in P_\Gamma$. Applying Lemma 6.4.2 to this fact we conclude that Γ satisfies the fourth condition of the main theorem of [CMS04]. It remains to check that the quotient Γ/Δ has exponent two. Let $\gamma \in \Gamma$ and set $r = \Delta\gamma \cdot \infty$. If we take $\Sigma_r = \Delta\gamma$ then the fourth condition of Theorem 6.5.1 states that $T_{\Delta|r}^{(1)}(z) = T_{\Delta,r}^{(1)}(z)$. Proposition 6.3.2 now implies that $\Sigma_r^2 = \Delta$, which in turn implies $\gamma^2 \in \Delta$. This argument applies to arbitrary $\gamma \in \Gamma$ so we conclude that the quotient Γ/Δ has exponent two. This completes the proof. $\square$

Perhaps the most technical condition of Theorem 6.5.1 is the last one. Beyond the monstrous moonshine conjectures, there are the generalized moonshine conjectures of Norton (cf. [Mas87]) which associate genus zero groups to commuting pairs of elements in the Monster. A number of the groups appearing in generalized moonshine do not satisfy the last condition of Theorem 6.5.1, but they all satisfy the second property, and a slight weakening of the first (cf. [Fer93]), and we do not know of any examples that fail to satisfy the third condition of Theorem 6.5.1. It is an interesting question then to determine if the first three conditions of Theorem 6.5.1 furnish

a characterization of the groups of generalized moonshine.

7 Gravity

In this section we consider applications of the normalized Rademacher sums to chiral 3d quantum gravity.

7.1 First conjecture

It is clear from our results that the Rademacher sums are particularly convenient for understanding the special characteristics of the McKay–Thompson series; in particular, their crucial genus zero property. In order to fully explain the moonshine phenomena, one has to relate these sums to the structure of the vertex operator algebra $V^\natural$. In search for the new relation, one might look again for hints from physics. Very recently, Witten, revisiting 3d quantum gravity, has formulated a number of results and observations [Wit07] including a conjecture about the existence of the 3d quantum gravities with central charges $c_L = c_R$ proportional to 24. In particular, he asserted that the simplest in his list of 3d quantum gravities should be equivalent to the 2d conformal field theory $V^\natural \otimes (V^\natural)^*$. Manschot [Man07] then suggested to consider chiral 3d quantum gravities, the simplest of which, with $c_L = 24$ and $c_R = 0$, should be equivalent exactly to $V^\natural$. Furthermore, Li, Song and Strominger [LSS08] have argued that the chiral gravity possesses stability and consistency, the necessary properties of a sound physical theory. A very recent work [MSS09] by Maloney, Song and Strominger provided further support for the existence of chiral gravities with c_L proportional to 24. These results and observations from physics taken together suggest that there exists an alternative construction of the vertex operator algebra $V^\natural$ that may be viewed as a rigorous version of the simplest chiral 3d quantum gravity with $c = 24$, in the same way as the original construction of $V^\natural$ was interpreted as a rigorous version of the chiral 2d conformal field theory with the partition function $J(z)$. In 3d quantum gravity one expects to obtain the partition function as a sum over minimum points of a 3d quantum gravity action; i.e. over all 3d hyperbolic structures on a solid torus with genus one boundary, whose conformal structure corresponds to the point z on the moduli space. Since all such structures are naturally parameterized by $\Gamma_\infty \backslash PSL_2(\mathbb{Z})$ (cf. [DMMV00]), the partition function should be a kind of Rademacher sum. In fact Manschot and Moore [MM07] argued that the subtraction of constants in (1.2.1) can be explained by a regularization of the partition function of 3d gravity. Our continued Rademacher sums of §4, leading naturally to the modified and normalized Rademacher sums, provide a good candidate for such a regularization.

To obtain the McKay–Thompson series (1.1.3) from the chiral 3d quantum gravity is a more challenging problem even at the heuristic level. However, one may look at these series from a slightly different point of view. It has been shown in [DLM00] that for any $g \in \mathbb{M}$ there exists a

unique simple g -twisted $V^\natural$ -module $V_g^\natural$, whose partition function is equal to $c(g)T_g(-1/z)$ where $c(g)$ is a constant depending on $g \in \mathbb{M}$. The heuristic analysis of the twisted sector $V_g^\natural$ in [Tui95] strongly supports the general assumption that $c(g) = 1$ for all $g \in \mathbb{M}$. This assumption was proven recently by Carnahan in [Car09].

We now state our first conjecture.

Conjecture 7.1.1. *There exists a family of twisted chiral 3d quantum gravities at central charge $c = 24$ associated with elements of the Monster $g \in \mathbb{M}$ whose partition functions are naturally given by sums over geometries parameterized by $\Gamma_\infty \backslash \Gamma_g$, and these partition functions coincide with the normalized Rademacher sum $T_\Gamma^{(1)}(-1/z)$, for $\Gamma = \Gamma_g$. Moreover, the untwisted ($g = e$) chiral 3d quantum gravity has a VOA structure isomorphic to $V^\natural$, and the twisted 3d quantum gravity corresponding to $g \in \mathbb{M}$ has a structure of g -twisted $V^\natural$ -module isomorphic to that of $V_g^\natural$.*

The special case $g = e$ of Conjecture 7.1.1, corresponding to the untwisted chiral 3d quantum gravity, is strongly supported by the physics literature mentioned above. We will now make a few remarks about the general twisted case. The principle of the 3d quantum gravity/2d CFT correspondence suggests that all the structures in either theory should have appropriate counterparts in the other. The g -twisted sector $V_g^\natural$ is an intrinsic part of the extended theory of the chiral 2d CFT associated to $V^\natural$, and, according to the physical principle, should therefore have analogues in chiral 3d quantum gravity. Basic information about these twisted chiral 3d quantum gravities can be extracted from the Rademacher sums $J_g(z) = T_g(-1/z)$. We'll illustrate the general case with the level two examples, which are the cases that $\Gamma = \Gamma_0(2)$ or $\Gamma_0(2)+$. (In the notation of §2.5 the symbols $\Gamma_0(2)+$ are a shorthand for $\Gamma_0(2) + \text{Ex}(2)$.) As in the case that $g = e$ one expects that each term of the Rademacher sum comes from a classical solution of the corresponding twisted chiral quantum gravity. Every Rademacher sum $J_g(z)$ contains the term $e(1/z)$, which corresponds to the BTZ black hole solution (cf. [BTZ92]). The other terms arising for $\Gamma = \Gamma_0(2)$ can be characterized among all those arising for $\Gamma = G(\mathbb{Z})$ as the solutions that have the same spin structure as the BTZ black hole on the boundary. The Rademacher sum $J_g(z)$ in this case coincides with the partition function of the Ramond sector in supergravity (cf. [MW07, §7]). The group $\Gamma = \Gamma_0(2)+$ is no longer a subgroup of the modular group $G(\mathbb{Z})$. In this case we have to consider orbifold solutions in addition to smooth solutions; we consider orbifold solutions with a codimension 2 singularity along the defining circle of the solid torus which looks locally like $\mathbb{C}/\langle e(1/2) \rangle$ (cf. [MW07, §2]), and we still impose the same spin structure boundary condition. Note that on any space X , choices of spin structure on X (when they exist) are in bijective correspondence with a particular family of double covers of X (cf. e.g. [LM89]). Such a double cover smooths the $\mathbb{Z}/2$ -orbifold singularity. One may view the spin structure as a $\mathbb{Z}/2$ -structure, and for a general level N group Γ , spin structures are replaced with $\mathbb{Z}/N$ -structures. One can also allow singularities along the defining circle of the solid torus that look locally like $\mathbb{C}/\langle e(1/N) \rangle$. In general, a choice of twisted gravity with $\mathbb{Z}/N$ -structure

group should impose an $n||h + S$ equivalence for solid tori, for some n and h with $N = nh$ and h a divisor of 24 (cf. Theorem 6.2.4), which yields the Rademacher sum $J_g(z)$, for a corresponding $g \in \mathbb{M}$, as the saddle point approximation of the twisted quantum gravity partition function.

We expect that the conjectural twisted chiral 3d quantum gravity construction of $V_g^{\frac{1}{2}}$ will imply the Γ_g -invariance of its partition function, in analogy with the way in which 2d CFTs are found to have modular invariant partition functions (cf. [Wit87]). Then our conjecture, in combination with our results on Rademacher sums, will naturally imply the genus zero conjecture of Conway–Norton (cf. [CN79]), and most importantly, will reveal the geometric nature of moonshine. In particular, the solution to Conjecture 7.1.1 will answer the genus zero question: why the discrete groups attached to the Monster via monstrous moonshine all have genus zero.

We also expect that the conditions on Rademacher sums in our reformulation Theorem 6.5.1, of the group theoretic characterization of the groups of monstrous moonshine due to Conway–McKay–Seiberg, will also find an interpretation as consistency conditions for twisted chiral 3d quantum gravities. In particular, the completeness of the family of twisted chiral 3d quantum gravities at central charge 24 associated with elements of the Monster group $\mathbb{M}$ should have a deep meaning in quantum gravity.

Our Conjecture 7.1.1 should have significance for the future development of analytic number theory also. We already know from 2d CFT that various constructions yield remarkable number theoretic identities. Our conjecture implies that the development of the theory of 3d quantum gravity might encompass a whole new family of number theoretic results such as the theory of Rademacher sums studied in this work. In §7.3 we’ll show that the Hecke operators also admit an interpretation in terms of quantum gravity. In preparation for this we will recall in the next section some results about a certain class of generalized Kac–Moody algebras and their representations.

7.2 Monstrous Lie algebras

In his paper [Bor92], Borchers constructed a remarkable generalized Kac–Moody algebra (*GKM algebra*) $\mathfrak{m}$, called the *Monster Lie algebra*, which plays a key rôle in his approach to monstrous moonshine. He also defined GKM superalgebras $\mathfrak{m}'_g$, for each $g \in \mathbb{M}$, by using the McKay–Thompson series $T_g(z)$ to specify the simple roots. In this section we’ll consider another family of GKM algebras $\mathfrak{m}_g$, parameterized by elements $g \in \mathbb{M}$, with simple roots specified by the functions $J_g(z) = T_g(-1/z)$. Since the Fourier coefficients of the $J_g(z)$ are non-negative integers, the algebras $\mathfrak{m}_g$ are purely even. We will also study Verma modules for the GKM algebras $\mathfrak{m}_g$.

In the following section §7.3, we will explain how these algebraic structures arise from the Rademacher sums, and the conjectural twisted chiral 3d quantum gravities. This will lead us to a further extension of our first conjecture.

The algebras $\mathfrak{m}_g$ were introduced and studied by Carnahan in [Car07] and were discovered

independently by the second author. We will now recall the relevant results of [Car07].

Let $L = II_{1,1}$ denote a copy of the unique even self-dual Lorentzian lattice of rank 2. We identify L with the group $\mathbb{Z} \times \mathbb{Z}$ and set the norm of the pair (m, n) to be $-2mn$. For any positive integer N we denote the sublattice $\mathbb{Z} \times N\mathbb{Z}$ by $L(N)$, and we let $L(N)^\vee$ denote the dual lattice $\frac{1}{N}\mathbb{Z} \times \mathbb{Z}$. Let V_L denote the vertex operator algebra (VOA) associated to the lattice L (cf. [FLM88]),

$$V_L = \bigoplus_{(m,n) \in L} V_L^{(m,n)}, \quad (7.2.1)$$

and let h_N denote the automorphism of V_L which acts as multiplication by $e(n/N)$ on the subspace $V_L^{(m,n)}$. Then $V_{L(N)}$ is a vertex operator subalgebra of $V_{L(N)}$ fixed by h_N . Following [DL93] each coset $L + (k/N, 0)$ of L in $L(N)^\vee$ defines an h_N^k -twisted module for V_L . We denote this twisted module by $V_{L+(k/N,0)}$, and regard it as graded in the natural way by $L + (k/N, 0)$.

For $g \in \mathbb{M}$ of order N we define $W_g^\natural$ to be the following $L(N)^\vee$ -graded space invariant under $\langle g \rangle$.

$$W_g^\natural = \bigoplus_{k=0}^{N-1} \left(V_{g^k}^\natural \otimes V_{L+(k/N,0)} \right)^{\langle g \rangle} \quad (7.2.2)$$

The summand corresponding to $k = 0$ has a natural VOA structure, while the other summands are naturally modules for this VOA. There also exists a unique (up to scalar) intertwining operator between the product of the k -th and l -th summands and the $(k + l)$ -th summand. Carnahan asserts that there is a consistent choice of these constants such that one has

Theorem 7.2.1. *The space $W_g^\natural$ naturally admits a VOA structure.*

By construction $W_g^\natural$ has rank 26 and one can define the semi-infinite cohomology of the Virasoro algebra with coefficients in $W_g^\natural$. The first semi-infinite cohomology group acquires a Lie algebra structure according to [LZ95]. In this way we obtain a *monstrous Lie algebra* $\mathfrak{m}_g$ for each $g \in \mathbb{M}$.

$$\mathfrak{m}_g = H^{\infty/2+1}(W_g^\natural) \quad (7.2.3)$$

An alternative construction of $\mathfrak{m}_g$ may be based on the no-ghost theorem of the 26-dimensional bosonic string (cf. [Fre85], [Bor92]). For $g = e$, the Lie algebra $\mathfrak{m}_e$ is the original Monster Lie algebra of Borcherds [Bor92].

The Lie algebra $\mathfrak{m}_g$ inherits an $L(N)^\vee$ -grading

$$\mathfrak{m}_g = \bigoplus_{m,n \in \mathbb{Z}} \mathfrak{m}_g^{(m/N, n)}. \quad (7.2.4)$$

The same argument as in the case $g = e$ applied to the semi-infinite cohomology or no-ghost theorem construction of $\mathfrak{m}_g$ yields natural isomorphisms

$$\mathfrak{m}_g^{(m/N, n)} \cong V_{g^{-m}, n/N}^{\natural}(mn/N) \quad (7.2.5)$$

where $V_{g^{-m}, n/N}^{\natural}$ is the subspace of $V_{g^{-m}}^{\natural}$ upon which g acts by $e(n/N)$ and $V_{g^{-m}, n/N}^{\natural}(mn/N)$ is the L_0 -eigenspace of $V_{g^{-m}, n/N}^{\natural}$ with eigenvalue $mn/N + 1$.

Remark 7.2.2. Let us write V^{Λ} for the vertex operator algebra associated to the Leech lattice. In his paper [Tui95] Tuite gave strong evidence that the genus zero property of monstrous moonshine is equivalent to the following duality isomorphisms,

$$V_{g^m, n/N}^{\natural} \cong V_{g^n, m/N}^{\natural}, \quad (7.2.6)$$

$$V_{g^m, n/N}^{\natural} \cong V_{h^n, m/N}^{\Lambda}, \quad (7.2.7)$$

the first holding in the case that g is Fricke, and the second in the case that g is non-Fricke. The h in (7.2.7) denotes an automorphism of Λ lifting to the automorphism g in $\mathbb{M}$. Thus the Tuite duality in the Fricke case implies the symmetry

$$\mathfrak{m}_g^{(m/N, n)} \cong \mathfrak{m}_g^{(n/N, m)} \quad (7.2.8)$$

of the monstrous Lie algebra $\mathfrak{m}_g$, as is known to hold in the case $g = e$ studied in [Bor92]. In the non-Fricke case the Tuite duality suggests an alternative construction of the VOA $W_g^{\natural}$ as

$$W_g^{\natural} = \bigoplus_{k=0}^{N-1} \left(V_{h^k}^{\Lambda} \otimes V_{\tilde{L}+(0, k)} \right)^{\langle h \rangle} \quad (7.2.9)$$

where $\tilde{L} = \frac{1}{N}\mathbb{Z} \times N\mathbb{Z}$.

We now re-scale the grading in $W_g^{\natural}$ and $\mathfrak{m}_g$ by interchanging $(m/N, n)$ with $(m, n/N)$, and we set

$$\mathfrak{m}_g^m = \bigoplus_{n \in \frac{1}{N}\mathbb{Z}} \mathfrak{m}_g^{(m, n/N)} \quad (7.2.10)$$

for $m \in \mathbb{Z}$. Then (7.2.5) implies isomorphisms of graded spaces

$$\mathfrak{m}_g^1 \cong V_{g^{-1}}^{\natural}, \quad \mathfrak{m}_g^{-1} \cong V_g^{\natural}, \quad (7.2.11)$$

and we also have

$$\mathfrak{m}_g^0 = \mathfrak{m}_g^{(0,0)} = \mathbb{C}c \oplus \mathbb{C}d, \quad (7.2.12)$$

where c and d are the degree operators for the re-scaled grading.

In [Car07] Carnahan shows that $\mathfrak{m}_g$ is a GKM algebra, thus generalizing a result of [Bor92]. When g is Fricke, $\mathfrak{m}_g$ has a structure very similar to that of the original Borchers Monster Lie algebra $\mathfrak{m} = \mathfrak{m}_e$. In particular, it has one real simple root and all simple roots correspond to a basis for $V_g^\natural$. This implies that $\mathfrak{m}_g$ can be reconstructed from the subalgebra

$$\mathfrak{m}_g^1 \oplus \mathfrak{m}_g^0 \oplus \mathfrak{m}_g^{-1} \quad (7.2.13)$$

which is called the *local subalgebra of $\mathfrak{m}_g$* (cf. [FF83]).

Remark 7.2.3. When g is non-Fricke $\mathfrak{m}_g$ does not have real simple roots, and in general it cannot be reconstructed from its local subalgebra (7.2.13). However the Tuite duality suggests that if one considers the second grading of $\mathfrak{m}_g$ then the local subalgebra consists of V_h^Λ and $V_{h^{-1}}^\Lambda$ and the h -twisted Heisenberg algebra. This local subalgebra is expected to generate $\mathfrak{m}_g$ in the case that g is non-Fricke.

In his thesis [Car07] Carnahan also obtained remarkable generalizations of the Borchers identities for each of the monstrous Lie algebras $\mathfrak{m}_g$; viz.,

$$p(T_g(w) - J_g(z)) = \prod_{m \in \mathbb{Z}^+, n \in \mathbb{Z}} (1 - p^m q^{n/N})^{c_{g^m, n/N}(mn/N)} \quad (7.2.14)$$

where we have set

$$c_{g^m, n/N}(mn/N) = \dim V_{g^m, n/N}^\natural(mn/N). \quad (7.2.15)$$

When g is of the Fricke type the invariance of $T_g(w)$ under the Fricke involution and a re-scaling $w \mapsto Nw$ yields

$$p(J_g(Nw) - J_g(Nz)) = \prod_{m \in \mathbb{Z}^+, n \in \mathbb{Z}} (1 - p^m q^n)^{c_{g^m, n/N}(mn/N)}. \quad (7.2.16)$$

Dividing both sides of (7.2.16) by $(1 - pq^{-1})$ we obtain expressions which are invariant under the transposition of p with q . This entails the identity

$$c_{g^m, n/N}(mn/N) = c_{g^m, m/N}(mn/N) \quad (7.2.17)$$

which is in agreement with the Tuite duality of (7.2.6). The identity (7.2.14) implies an un-

expected formula for the graded dimension of the Verma module with trivial character $\mathcal{V}_g$ associated to g .

$$\mathcal{V}_g = \mathcal{U}(\mathfrak{m}_g^-) \quad (7.2.18)$$

In fact the standard product formula

$$\text{gdim } \mathcal{V}_g = \prod_{m \in \mathbb{Z}^+, n \in \mathbb{Z}} (1 - p^m q^{n/N})^{-c_{g^{m, n/N}}(mn/N)} \quad (7.2.19)$$

for the graded dimension of $\mathcal{V}_g$ follows from (7.2.5) and (7.2.15). The identity (7.2.14) yields the alternative expression

$$\text{gdim } \mathcal{V}_g = \frac{1}{p(T_g(w) - J_g(z))}. \quad (7.2.20)$$

This expression allows us to view the bi-graded dimension as a meromorphic function on $\mathbb{H} \times \mathbb{H}$. Thanks to the properties of principal moduli for curves of genus zero, the expression (7.2.20) is singular at the point (w, z) precisely when $\Gamma_g \cdot w = \Gamma_g \cdot (-1/z)$. In §7.3 we will give an interpretation of this fact in the setting of three dimensional quantum gravity.

With the goal of making such an interpretation in mind we rewrite the first expression (7.2.19) for the bi-graded dimension of the Verma module $\mathcal{V}_g$ using the generalized Hecke operators of §5.5.

$$\text{gdim } \mathcal{V}_g = \exp \left(\sum_{m \in \mathbb{Z}^+} \hat{T}(m) J_g(z) p^m \right) \quad (7.2.21)$$

Comparing with the generating functions

$$Z_g(p, q) = \sum_{m \in \mathbb{Z}^+} m \hat{T}(m) J_g(z) p^m \quad (7.2.22)$$

of the higher order Rademacher sums $J_g^{(m)}(z)$, we obtain the following expression.

$$Z_g(p, q) = p \partial_p \log(\text{gdim } \mathcal{V}_g) \quad (7.2.23)$$

An identity of this form is well-known in the theory of symmetric functions (cf. [Mac95]), where it serves to relate the generating function of the power symmetric functions

$$p_n = \sum_{1 \leq i} x_i^n \quad (7.2.24)$$

with that of the complete symmetric functions

$$h_n = \sum_{1 \leq i_1 \leq \dots \leq i_n} \prod_{k=1}^n x_{i_k}. \quad (7.2.25)$$

Thus the graded dimensions of the Verma modules $\mathcal{V}_g$ may be viewed as *complete Rademacher sums*, where the rôle of the variables x_i , for $i \in \mathbb{Z}^+$, in (7.2.24) and (7.2.25) is taken up by the exponential expressions $e(\llbracket \gamma \cdot z' \rrbracket)$ for $z' = -1/z$ and $\llbracket \gamma \rrbracket \in \llbracket \Gamma_g \rrbracket$.

Since all the coefficients appearing in the bi-graded dimension $Z_g(p, q)$ are non-negative integers, one may expect to naturally find a bi-graded vector space, and possibly even an $\mathfrak{m}_g$ -module, with bi-graded dimension given by $Z_g(p, q)$. The next result suggests that one can expect to find such a space within $\mathcal{V}_g$.

Given two elements $F = \sum F_{m,n} p^m q^n$ and $G = \sum G_{m,n} p^m q^n$ in $\mathbb{Z}((q))[[p]]$, let us write $F \leq G$ in the case that $F_{m,n} \leq G_{m,n}$ for all $m, n \in \mathbb{Z}$.

Proposition 7.2.4. *Let $g \in \mathbb{M}$. If g is Fricke then we have $Z_g(p, q) \leq \text{gdim } \mathcal{V}_g$.*

Proof. Dividing both sides of (7.2.16) by $(1 - pq^{-1})$ we obtain

$$1 - \sum_{m,n \in \mathbb{Z}^+} c_{g, \frac{m+n-1}{N}} \left(\frac{m+n-1}{N} \right) p^m q^n = \prod_{m,n \in \mathbb{Z}^+} (1 - p^m q^n)^{c_{g,m,n/N}(mn/N)}. \quad (7.2.26)$$

We denote the sum in the left hand side by Σ^+ and the product in the right hand side by Π^+ . Then we have

$$Z_g(p, q) = -p \partial_p \log(1 - pq^{-1}) \Pi^+ = \frac{pq^{-1}}{1 - pq^{-1}} + \frac{p \partial_p \Sigma^+}{1 - \Sigma^+}. \quad (7.2.27)$$

On the other hand

$$\text{gdim } \mathcal{V}_g = \frac{1}{1 - pq^{-1}} \frac{1}{1 - \Sigma^+} = \frac{1}{1 - pq^{-1}} + \frac{\Sigma^+}{(1 - pq^{-1})(1 - \Sigma^+)}. \quad (7.2.28)$$

Thus it is sufficient to show that

$$p \partial_p (\Sigma^+) \leq \frac{1}{1 - pq^{-1}} \Sigma^+, \quad (7.2.29)$$

but this follows from the observation that both sides of (7.2.29) have nonnegative coefficients and the left hand side may be obtained from the right hand side by deleting the terms which are singular or constant with respect to q . $\square$

To formulate a conjecture on the nature of the subspaces of $\mathcal{V}_g$ with bi-graded dimensions $Z_g(p, q)$, as well as the origin of the higher order Rademacher sums, we turn again, in the next

section, to the structures of twisted chiral 3d quantum gravity.

7.3 Second conjecture

We have shown in §5.5 that the higher order Rademacher sums recover the action of the Hecke operators on the first order Rademacher sums. In view of the conjectural relation between first order Rademacher sums and chiral 3d quantum gravities at central charge $c = 24$, we may consider an analogous interpretation for the higher order Rademacher sums given by $m(\hat{T}(m)J_g)(z)$ for $g \in \mathbb{M}$ and $m \in \mathbb{Z}^+$. Since all the Fourier coefficients of these expressions are non-negative integers, one might guess that there exist families of twisted chiral 3d quantum gravities for all central charges $c = 24m$, for $m \in \mathbb{Z}^+$. In particular, the untwisted ($g = e$) chiral 3d quantum gravities possess the structure of certain extremal vertex algebras with the Monster symmetry.

The aforementioned class of extremal vertex algebras, with the addition of a Virasoro element, has been originally conjectured by Witten [Wit07]. However, it was shown in [Gai08] and [Hoe08] that the addition of a Virasoro element precludes a non-trivial action of the Monster group. The higher order Rademacher sums also point to the partition functions without elements of spin 2. We may modify Witten's conjecture by asking for the existence of extremal vertex algebras with partition functions given by the function $J^{(m)}(z)$, related to the higher order Rademacher sums via $J^{(m)}(z) = Q_\Gamma^{(m)}(z) - c_\Gamma(m, 0)$, with $\Gamma = PSL_2(\mathbb{Z})$. The validity of this conjecture remains open.

Note that the Virasoro algebra, even when not represented by any actual state, may still act on a vertex algebra with partition function $J^{(m)}(z)$. Regardless of whether or not spaces $V^{(m)}$ with partition function $J^{(m)}(z)$ admit natural vertex algebra structures, they certainly naturally inherit actions of the Monster group, since the actions of Hecke operators $m\hat{T}(m)$ on the Fourier coefficients of $J(z)$ may be interpreted as actions on representations of $\mathbb{M}$ (cf. [JLW95]).

It is natural to enquire as to the meaning of the spaces $V_g^{(m)}$, for $m \in \mathbb{Z}^+$ and $g \in \mathbb{M}$, with graded dimension $J_g^{(m)}(z)$. In light of our First Conjecture, it is unlikely that the spaces $V^{(m)}$ can represent states of a three dimensional quantum gravity with $c = 24m$, since a Virasoro element is generally not present. Thus we have to conclude that the three dimensional quantum gravity $V^{(1)} = V^\natural$ is the only viable candidate. Then one might view the spaces $V^{(m)}$ for $m > 1$ as some higher overtones of the basic $m = 1$ theory.

In fact, we have shown in §5.5 that the action of the classical Hecke operator $\hat{T}(m)$ on the Rademacher sum (1.2.1) yields a sum over $\Gamma_\infty \backslash M(m)$, where $M(m)$ denotes the (image in $G(\mathbb{Q})$ of the) set of 2×2 matrices with integral entries and determinant m . This sum may be viewed as the *m-instanton correction* of the partition function in three dimensional quantum gravity. By *n-instanton* in this context one understands an elliptic curve which admits a holomorphic map of degree n into a given elliptic curve $E_{\mathbb{Z}}$ (cf. [DMVV97]). Thus the sum over $\Gamma_\infty \backslash M(n)$ becomes a sum over all three dimensional hyperbolic structures on a solid torus with genus one boundary

whose conformal structure corresponds to an n -instanton on $E_{\mathbb{Z}}$. The same phenomena is expected for all the twisted 3d quantum gravities corresponding to elements $g \in \mathbb{M}$.

Collecting all the m -instanton contributions in one generating function we obtain a partition function

$$Z_g(p, q) = \sum_{m \in \mathbb{Z}^+} m(\hat{T}(m)J_g)(z)p^m \quad (7.3.1)$$

depending on two variables, for each $g \in \mathbb{M}$. This partition function $Z_g(p, q)$ may be viewed as a part of the full partition function

$$\tilde{Z}_g(p, q) = \exp \left(\sum_{m \in \mathbb{Z}^+} (\hat{T}(m)J_g)(z)p^m \right) \quad (7.3.2)$$

of the stringy second quantization of $V_g^{\natural}$, introduced in [DMVV97] in the untwisted case $g = e$. (See also [Tui08].) We will now generalize their construction to an arbitrary twisted module $V_g^{\natural}$.

Let V be an arbitrary vertex operator algebra; in our case $V = V^{\natural}$. For n a positive integer, let $V^{\otimes n}$ denote the tensor product of n copies of V , and let Z_n be the group of cyclic permutations of the factors generated by an n -cycle $\sigma_n = (1, 2, \dots, n)$. Let g be an automorphism of V of order N , then $g \times \sigma_n$ is an automorphism of $V^{\otimes n}$ of order nN/d where $d = (n, N)$. To the pair $(V^{\otimes n}, g \times \sigma_n)$ is canonically associated a twisted module (cf. [BDM02]), which we denote $V_{(g,n)}$. The group $Z_{g,n} = \langle g \times \sigma_n \rangle$ acts naturally on $V_{(g,n)}$. We write $V_{(g,n)}^{Z_{g,n}}$ for the $Z_{g,n}$ -invariant subspace. The *stringy second quantization of the twisted module* V_g , to be denoted $\mathbb{S}V_g$, is, by definition, the space

$$\mathbb{S}V_g = \bigoplus_{\lambda \in \mathcal{P}} \bigotimes_{r > 0} S^{m_r} V_{(g,r)}^{Z_{g,r}}, \quad (7.3.3)$$

where the sum is taken over all partitions λ with m_r parts of length $r > 0$. The space $\mathbb{S}V_g$ is doubly graded: by the degrees of products of twisted sectors and by the value of $|\lambda|$. For $g = e$ our definition of the stringy second quantization coincides with that of [DMVV97].

Next we establish a relation between the second quantization $\mathbb{S}V_g^{\natural}$ and the Verma module $\mathcal{V}_g$. First note the canonical isomorphism

$$\mathcal{U}(\mathfrak{m}_g^-) \cong S(\mathfrak{m}_g^-). \quad (7.3.4)$$

Then we obtain the following result.

Theorem 7.3.1. *There is a canonical isomorphism of bi-graded vector spaces*

$$\mathbb{S} V_g^{\natural} \cong S(\mathfrak{m}_g^{-}). \quad (7.3.5)$$

Proof. The isomorphism (7.3.5) follows from the existence of isomorphisms

$$(V_{(g,n)}^{\natural})^{Z_{g,n}} \cong \mathfrak{m}_g^{-n} \quad (7.3.6)$$

of bi-graded vector spaces for each $n \in \mathbb{Z}^+$. To establish the existence of these we will use the isomorphism

$$V_{(g,n)}^{\natural} \cong V_{g^n}^{\natural} \quad (7.3.7)$$

of [BDM02].

We wish to show that the subspace of $V_{g^n}^{\natural}$ fixed by the operator

$$g \times e((L_0 - 1)/n) \quad (7.3.8)$$

is naturally isomorphic to $\mathfrak{m}_g^n$. Consider the invariant subspace of $V_{g^n}^{\natural}$ with respect to the N -th power of the operator (7.3.8), which may be expressed as $e((L_0 - 1)N/n)$. Its action on the graded subspace $V_{g^n}^{\natural}(kd/N)$ for $k \in \mathbb{Z}$ is multiplication by $e(kd/n)$, and this scalar is 1 if and only if $k = (n/d)m$ for some $m \in \mathbb{Z}$. Thus we are only concerned with the graded subspaces $V_{g^n}^{\natural}(nm/N)$ where $m \in \mathbb{Z}$. To find the invariant subspaces with respect to the operator (7.3.8) we consider its action on the subspaces of the form $V_{g^n, m'/N}^{\natural}(mn/N)$. The action is scalar multiplication by $e(-m'/N)e(m/N)$ and is therefore trivial if and only if $m \equiv m' \pmod{N}$. Thus the full subspace of $V_{g^n}^{\natural}$ invariant under (7.3.8) is given by

$$\bigoplus_{m \in \mathbb{Z}} V_{g^n, m/N}^{\natural}(mn/N) \cong \mathfrak{m}_g^{-n}. \quad (7.3.9)$$

To complete the proof we note that the group generated by the action of the operator (7.3.8) has order nN/d , and under the isomorphism (7.3.7) its action recovers that of $Z_{g,n}$. $\square$

The isomorphism of Theorem 7.3.1 suggests that the stringy second quantization $\mathbb{S} V_g^{\natural}$ admits an action by the Lie algebra $\mathfrak{m}_g$ and thus provides a Fock space realization of the Verma module $\mathcal{V}_g$. When $\mathfrak{m}_g$ is generated by its local subalgebra (7.2.13), as happens in the Fricke case, it is sufficient to describe the action of brackets $[\mathfrak{m}_g^{\pm 1}, \mathfrak{m}_g^{-n}] \subset \mathfrak{m}_g^{-n \pm 1}$ via the action of $V_{g^{\pm 1}}^{\natural}$ on the twisted sector for $((V^{\natural})^{\otimes n}, g \times \sigma_n)$. Note also that the removal of the subspaces corresponding to the terms which are singular or constant with respect to q , as in the proof of Proposition 7.2.4, will yield an embedding of the instanton subspace $\mathcal{I}_g$ inside the stringy second quantization of

$V_g^\natural$.

Our interpretation of the higher order Rademacher sums and the complete Rademacher sums, and their relation to the second quantization of the moonshine vertex operator algebra and its twisted modules, as well as the monstrous Lie algebras and their Verma modules, suggests the following extension of our first conjecture, Conjecture 7.1.1.

Conjecture 7.3.2. *Assuming the existence of a family of g -twisted chiral 3d quantum gravities for $g \in \mathbb{M}$, having the properties stated in Conjecture 7.1.1, there also exists a natural geometric interpretation of the family of twisted monstrous Lie algebras $\mathfrak{m}_g$, the instanton spaces $\mathcal{I}_g$, and the denominator formulas for all $g \in \mathbb{M}$ via the second quantization of the corresponding g -twisted chiral three 3d quantum gravities.*

First of all, the partition function of the second quantized g -twisted chiral 3d quantum gravity should depend on two modular parameters w and z , and should respect the symmetry that interchanges w with $-1/z$. Second, there should be a quantum gravity theoretic explanation for why these partition functions are singular precisely when w and $-1/z$ belong to the same orbit of Γ_g . This will explain the remarkable denominator formulas (7.2.14) of Carnahan, and ultimately, the fundamental rôle of the principal moduli in the moonshine Conjectures.

Where can we find an appropriate setting for all the structures that appear in our Second Conjecture? The first answer that might come to mind is the 26 dimensional bosonic string theory. It was known for a long time that the physical space has a Lie algebra structure (cf. [Fre85], [GO85]), which can also be recovered from the semi-infinite cohomology (cf. [FGZ86], [LZ95]). However, in this way one can only get a fake Monster Lie algebra (cf. [Bor92]) and there is no (straight forward) geometric way to turn it into the real Monster Lie algebra that appears in the three dimensional quantum gravity approach. Also, the 26 dimensions of the bosonic string have nothing to do with the 3 dimensions of the quantum gravity. It is still possible that some constructions of string theory can be applied in the 3d quantum gravity setting. In particular, one can expect to identify the negative part $\mathfrak{m}^-$ of the Monster Lie algebra in the space of the second quantization of $V^\natural$ with a certain version of BPS states (cf. [HM96]).

Clearly, our results on Rademacher sums, their relation to the moonshine module, monstrous Lie algebras, and 3d quantum gravity, admit a super-counterpart [DF09]. In this case the promise of relationships with the structures of the 10 dimensional superstring is even more tempting, since the latter has played such a prominent rôle in physics over the past 25 years. However, in spite of the of the remarkable mathematical similarity, the 10 of the ten dimensional superstring and the 3 of the 3d quantum supergravity emerging from the super-counterpart of the Rademacher sums have different geometric meaning and are not related by a compactification of any kind.

The most fascinating fact about the 3d quantum supergravity is that while it is similar, but not directly related, to the initial 3d quantum gravity, the corresponding second quantized theo-

ries are directly related. The higher order Rademacher sums point to a larger second quantized space in the super-case; a space which contains the second quantized space we considered above.

Then what can these second quantized 3d quantum gravities mean in physics? Naïvely, any quantization can be interpreted as a categorification, which lifts a given theory one dimension up. One can then wonder if the second quantization in our case might point to an extreme sector of certain four dimensional quantum gravities, such as the extreme Kerr black hole, which was recently found in [GHSS08] also to be dual to a chiral 2d CFT. In this case the Monster (or moonshine) might be the answer to the perpetual question of what is behind the letter M in the theory that has not yet revealed its true name.

References

- [BDM02] Katrina Barron, Chongying Dong, and Geoffrey Mason. Twisted sectors for tensor product vertex operator algebras associated to permutation groups. *Comm. Math. Phys.*, 227(2):349–384, 2002.
- [Bor92] Richard E. Borcherds. Monstrous moonshine and monstrous Lie superalgebras. *Invent. Math.*, 109(2):405–444, 1992.
- [BTZ92] Máximo Bañados, Claudio Teitelboim, and Jorge Zanelli. Black hole in three-dimensional spacetime. *Phys. Rev. Lett.*, 69(13):1849–1851, 1992.
- [Car07] Scott Carnahan. *Monstrous Lie algebras and generalized moonshine*. PhD thesis, University of California, Berkeley, 2007.
- [Car09] Scott Carnahan. Generalized moonshine II: Borcherds products. arXiv:0908.4223v2, October 2009.
- [CMS04] John Conway, John McKay, and Abdellah Sebbar. On the discrete groups of Moonshine. *Proc. Amer. Math. Soc.*, 132:2233–2240, 2004.
- [CN79] J. H. Conway and S. P. Norton. Monstrous Moonshine. *Bull. London Math. Soc.*, 11(3):308–339, 1979.
- [DF09] John F. R. Duncan and Igor B. Frenkel. Rademacher sums and super-moonshine. In preparation, 2009.
- [DL93] Chongying Dong and James Lepowsky. *Generalized vertex algebras and relative vertex operators*, volume 112 of *Progress in Mathematics*. Birkhäuser Boston Inc., Boston, MA, 1993.
- [DLM00] Chongying Dong, Haisheng Li, and Geoffrey Mason. Modular-invariance of trace functions in orbifold theory and generalized Moonshine. *Communications in Mathematical Physics*, 214:1–56, 2000.

-
- [DMMV00] Robbert Dijkgraaf, Juan Maldacena, Gregory Moore, and Erik Verlinde. A black hole Farey tail. arXiv:hep-th/0005003v3, April 2000.
- [DMVV97] Robbert Dijkgraaf, Gregory Moore, Erik Verlinde, and Herman Verlinde. Elliptic genera of symmetric products and second quantized strings. *Comm. Math. Phys.*, 185(1):197–209, 1997.
- [Fer93] Charles R. Ferenbaugh. The genus-zero problem for $n|h$ -type groups. *Duke Math. J.*, 72(1):31–63, 1993.
- [FF83] Alex J. Feingold and Igor B. Frenkel. A hyperbolic Kac-Moody algebra and the theory of Siegel modular forms of genus 2. *Math. Ann.*, 263(1):87–144, 1983.
- [FGZ86] I. B. Frenkel, H. Garland, and G. J. Zuckerman. Semi-infinite cohomology and string theory. *Proc. Nat. Acad. Sci. U.S.A.*, 83(22):8442–8446, 1986.
- [FLM88] Igor Frenkel, James Lepowsky, and Arne Meurman. *Vertex operator algebras and the Monster*, volume 134 of *Pure and Applied Mathematics*. Academic Press Inc., Boston, MA, 1988.
- [Fre85] I. B. Frenkel. Representations of Kac-Moody algebras and dual resonance models. In *Applications of group theory in physics and mathematical physics (Chicago, 1982)*, volume 21 of *Lectures in Appl. Math.*, pages 325–353. Amer. Math. Soc., Providence, RI, 1985.
- [Gai08] Davide Gaiotto. Monster symmetry and extremal CFTs. arXiv:0801.0988v1, January 2008.
- [GHSS08] Monica Guica, Thomas Hartman, Wei Song, and Andrew Strominger. The Kerr/CFT correspondence. arXiv:0809.4266v1, September 2008.
- [GO85] P. Goddard and D. Olive. Algebras, lattices and strings. In *Vertex operators in mathematics and physics (Berkeley, Calif., 1983)*, volume 3 of *Math. Sci. Res. Inst. Publ.*, pages 51–96. Springer, New York, 1985.
- [GS83] D. Goldfeld and P. Sarnak. Sums of Kloosterman sums. *Invent. Math.*, 71(2):243–250, 1983.
- [Hej83] Dennis A. Hejhal. *The Selberg trace formula for $\mathrm{PSL}(2, \mathbf{R})$. Vol. 2*, volume 1001 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1983.
- [HM96] Jeffrey A. Harvey and Gregory Moore. Algebras, BPS states, and strings. *Nuclear Phys. B*, 463(2-3):315–368, 1996.
- [Hoe08] G. Hoehn. Self-dual vertex operator superalgebras of large minimal weight. arXiv:0801.1822v1, January 2008.
- [Iwa97] Henryk Iwaniec. *Topics in classical automorphic forms*, volume 17 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 1997.

-
- [Iwa02] Henryk Iwaniec. *Spectral methods of automorphic forms*, volume 53 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, second edition, 2002.
 - [JLW95] E. Jurisich, J. Lepowsky, and R. L. Wilson. Realizations of the Monster Lie algebra. *Selecta Math. (N.S.)*, 1(1):129–161, 1995.
 - [Kno61a] Marvin Isadore Knopp. Construction of a class of modular functions and forms. *Pacific J. Math.*, 11:275–293, 1961.
 - [Kno61b] Marvin Isadore Knopp. Construction of a class of modular functions and forms. II. *Pacific J. Math.*, 11:661–678, 1961.
 - [Kno62a] Marvin Isadore Knopp. Construction of automorphic forms on H -groups and supplementary Fourier series. *Trans. Amer. Math. Soc.*, 103:168–188, 1962.
 - [Kno62b] Marvin Isadore Knopp. On abelian integrals of the second kind and modular functions. *Amer. J. Math.*, 84:615–628, 1962.
 - [Kno90] Marvin I. Knopp. Rademacher on $J(\tau)$, Poincaré series of nonpositive weights and the Eichler cohomology. *Notices Amer. Math. Soc.*, 37(4):385–393, 1990.
 - [KR01] Marvin Knopp and Sinai Robins. Easy proofs of Riemann’s functional equation for $\zeta(s)$ and of Lipschitz summation. *Proc. Amer. Math. Soc.*, 129(7):1915–1922 (electronic), 2001.
 - [LM89] H. Blaine Lawson, Jr. and Marie-Louise Michelsohn. *Spin geometry*, volume 38 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 1989.
 - [LSS08] Wei Li, Wei Song, and Andrew Strominger. Chiral gravity in three dimensions. *J. High Energy Phys.*, (4):082, 15, 2008.
 - [LZ95] Bong H. Lian and Gregg J. Zuckerman. Moonshine Cohomology. arXiv:q-alg/9501015v1, January 1995.
 - [Mac95] I. G. Macdonald. *Symmetric functions and Hall polynomials*. Oxford Mathematical Monographs. The Clarendon Press Oxford University Press, New York, second edition, 1995. With contributions by A. Zelevinsky, Oxford Science Publications.
 - [Man07] Jan Manschot. AdS_3 partition functions reconstructed. *J. High Energy Phys.*, (10):103, 7, 2007.
 - [Mas87] Geoffrey Mason. Finite groups and modular functions. In *The Arcata Conference on Representations of Finite Groups (Arcata, Calif., 1986)*, volume 47 of *Proc. Sympos. Pure Math.*, pages 181–210. Amer. Math. Soc., Providence, RI, 1987. With an appendix by S. P. Norton.
 - [MM07] Jan Manschot and Gregory Moore. A modern Farey tail. arXiv:0712.0573v1, December 2007.

-
- [MSS09] Alexander Maloney, Wei Song, and Andrew Strominger. Chiral gravity, log gravity and extremal CFT. arXiv:0903.4573v1, March 2009.
- [MW07] A. Maloney and E. Witten. Quantum gravity partition functions in three dimensions. arXiv:0712.0155v1, December 2007.
- [Nie73] Douglas Niebur. A class of nonanalytic automorphic functions. *Nagoya Math. J.*, 52:133–145, 1973.
- [Nie74] Douglas Niebur. Construction of automorphic forms and integrals. *Trans. Amer. Math. Soc.*, 191:373–385, 1974.
- [Nor84] S. P. Norton. More on moonshine. In *Computational group theory (Durham, 1982)*, pages 185–193. Academic Press, London, 1984.
- [Rad38] Hans Rademacher. The Fourier Coefficients of the Modular Invariant $J(\tau)$. *Amer. J. Math.*, 60(2):501–512, 1938.
- [Rad39] Hans Rademacher. The Fourier Series and the Functional Equation of the Absolute Modular Invariant $J(\tau)$. *Amer. J. Math.*, 61(1):237–248, 1939.
- [Tho79a] J. G. Thompson. Finite groups and modular functions. *Bull. London Math. Soc.*, 11(3):347–351, 1979.
- [Tho79b] J. G. Thompson. Some numerology between the Fischer-Griess Monster and the elliptic modular function. *Bull. London Math. Soc.*, 11(3):352–353, 1979.
- [Tui95] Michael P. Tuite. On the relationship between Monstrous Moonshine and the uniqueness of the Moonshine module. *Comm. Math. Phys.*, 166(3):495–532, 1995.
- [Tui08] Michael Tuite. Monstrous and Generalized Moonshine and permutation orbifolds. arXiv:0811.4525v1, November 2008.
- [Wit87] Edward Witten. Physics and geometry. In *Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Berkeley, Calif., 1986)*, pages 267–303, Providence, RI, 1987. Amer. Math. Soc.
- [Wit07] Edward Witten. Three-dimensional gravity revisited. arXiv:0706.3359v1, June 2007.